

Unified cosmic history in modified gravity: from $F(R)$ theory to Lorentz non-invariant models

Shin'ichi Nojiri^{1,2} and Sergei D. Odintsov^{3*}

¹ *Department of Physics, Nagoya University, Nagoya 464-8602, Japan*

² *Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Nagoya 464-8602, Japan*

³ *Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Ciències de l'Espai (IEEC-CSIC), Barcelona, Spain*

The classical generalization of general relativity is considered as the gravitational alternative for a unified description of the early-time inflation with late-time cosmic acceleration. The structure and cosmological properties of a number of modified theories, including traditional $F(R)$ and Hořava-Lifshitz $F(R)$ gravity, scalar-tensor theory, string-inspired and Gauss-Bonnet theory, non-local gravity, non-minimally coupled models, and power-counting renormalizable covariant gravity are discussed. Different representations of and relations between such theories are investigated. It is shown that some versions of the above theories may be consistent with local tests and may provide a qualitatively reasonable unified description of inflation with the dark energy epoch. The cosmological reconstruction of different modified gravities is provided in great detail. It is demonstrated that eventually any given universe evolution may be reconstructed for the theories under consideration, and the explicit reconstruction is applied to an accelerating spatially flat Friedmann-Robertson-Walker (FRW) universe. Special attention is paid to Lagrange multiplier constrained and conventional $F(R)$ gravities, for latter $F(R)$ theory, the effective Λ CDM era and phantom divide crossing acceleration are obtained. The occurrences of the Big Rip and other finite-time future singularities in modified gravity are reviewed along with their solutions via the addition of higher-derivative gravitational invariants.

PACS numbers: 95.36.+x, 98.80.Cq

* also at Tomsk State Pedagogical University, Tomsk.

Contents

I. Introduction	4
II. Modified gravity unifying the early-time inflation with late-time acceleration	6
A. $F(R)$ gravity	6
1. General properties	6
2. Scalar-tensor description	8
3. Matter instability	11
4. Unified theories without a flat space solution	12
5. Viable modified gravities	14
B. $f(\mathcal{G})$ gravity	21
1. General properties	21
2. $F(\mathcal{G}, R)$ gravity	26
C. String-inspired model and scalar-Einstein-Gauss-Bonnet gravity	26
D. Non-local gravity	28
1. Non-local $F(R)$ gravity	28
2. Non-local Gauss-Bonnet gravity	30
3. Other classical non-local Gauss-Bonnet models and their de Sitter solutions	33
E. Gravity non-minimally coupled with the matter Lagrangian	34
1. Non-minimally coupled scalar theory	34
2. Non-minimally coupled vector model	35
3. Non-minimal Yang-Mills theory	36
F. Modified $F(R)$ Hořava-Lifshitz gravity	37
1. General formulation	37
2. FRW cosmology	40
3. Unified inflation and dark energy in modified Hořava-Lifshitz gravity	41
G. Covariant power-counting renormalizable gravity	42
1. Perfect fluid formulation	42
2. Covariant field theory of gravity with a Lagrange multiplier	43
3. FRW cosmology	45
H. Dark fluid with an inhomogeneous equation of state	46
III. Cosmological reconstruction of modified gravity	49
A. Scalar-tensor gravity	49
1. The reconstruction using the cosmological time	49
2. The reconstruction of the scalar-tensor models using e-folding	52
3. Two-scalar model	52
4. Brans-Dicke gravity	56
B. The k -essence model	57
1. The reconstruction using the cosmological time	57
2. The reconstruction using e-folding	58
C. Modified $F(R)$ gravity	60
1. The reconstruction using the cosmological time	60
2. Reconstructed model realizing the phantom divide crossing	62
3. The reconstruction using e-folding	65
4. Stability of the cosmological solution	66
5. FRW cosmology in $F(R)$ gravity with a Lagrange multiplier	67
D. $F(\mathcal{G})$ gravity	70
E. Scalar-Einstein-Gauss-Bonnet gravity	71
F. $F(R)$ Hořava-Lifshitz gravity	72
G. Non-minimal Yang-Mills theory	75
IV. Finite-time future singularities in modified gravity	76
A. Future singularities in a dark, fluid universe: dark matter versus modified gravity	76
B. Finite-time singularities in $F(R)$ gravity	80
C. Future singularities in $F(\mathcal{G})$ gravity	85
D. Late-time dynamics in Hořava-Lifshitz gravity	88

V. Discussion	90
Acknowledgments	91
References	91

I. INTRODUCTION

Current theoretical cosmology is faced with two fundamental problems: what is inflation and what is dark energy? In other words, why and how did both the very early and the very late universe expand with acceleration? What is the reason for the similarity of the universe evolution at small as well as at large curvature?

The scenarios to describe these early-time and late-time accelerations are usually very similar, which is why it is quite natural to expect that the same theory/principle lies behind both accelerating cosmological epochs. Indeed, there are various proposals to construct an acceptable dark energy model, such as: scalar, spinor, (non-)abelian vector theory, cosmological constant, fluid with a complicated equation of state, and higher dimensions. Remarkably, the same proposals are also intended to describe inflationary era. As a result, we have a number of competing scenarios with which to describe inflation and late-time acceleration. This situation is very reasonable due to the fact that the evolution of the cosmological parameters is not defined with precise accuracy. Even their current values are defined with at least 3-5% error. Moreover, working in framework of one of the above proposals, we are forced to introduce the following extra cosmological components: inflaton, a dark component and dark matter. Even if such a scenario seems to be partially successful, it immediately introduce a new set of problems: such as the following: coupling with usual matter, (anti-)screening of dark components during the evolution of the universe, compatibility with standard elementary particle theories, and consistency of formulation. Thus, another natural proposal is to not introduce extra fields to resolve the cosmological problems, which is the most economical solution in the spirit of the Occam's razor principle.

The least understood and most fundamental force of nature is the gravitational interaction. It is expected that general relativity is just an approximation that is valid at small curvatures and that is predicted by the Theory of Everything (strings/M-theory). In the very early universe, some unknown gravitational theory dictated the evolution of the universe. Thus, a very reasonable assumption is the gravitational alternative for a unified description of the inflation and dark energy. In other words, the modified gravity, which represents a classical generalization of general relativity, should consistently describe the early-time inflation and late-time acceleration, without the introduction of any other dark component. A sector of modified gravity that contains the gravitational terms relevant at very high energies produced the inflationary epoch. In the course of evolution, the curvature decreases and general relativity gives a sufficient approximation in the intermediate universe. With a further decrease of the curvature, when subdominant terms quickly grow, one sees the change from deceleration to cosmic acceleration. Thus, the early-time as well as the late-time cosmic speed-up is caused simply by the fact that some subdominant terms of gravitational action may become essential at large or small curvatures. Moreover, such an approach may also be considered as dynamic solution of the cosmological constant problem. Of course, the complete gravitational action should be defined by a fundamental theory, which remains to be the open problem of modern high-energy physics. In the absence of fundamental quantum gravity, the modified gravity approach is a phenomenological model that is constructed by complying with observational data and data from local tests.

In addition, just as by-product to be a cosmological model, the modified gravity may provide the explanation for dark matter. It may resolve the coincidence problem simply by the fact of the universe expansion. It also may describe the transition from deceleration to acceleration of the universe, and it may be useful for high-energy physics problems (i.e., unification of all interactions, and hierarchy problem resolution). Even if the current universe is entering the phantom phase, modified gravity effectively describes the transition from the non-phantom to phantom era without the need to introduce the exotic matter (phantom) with extremely strange properties.

In this work, we present a detailed review of a number of popular models of modified gravity. Their properties and different representations are discussed. Assuming the spatially flat FRW cosmology, we investigate the modified gravity background evolution with the aim of achieving a unified description of the universe's inflation with dark energy epoch. Using sufficient freedom in the choice of the action as some function of curvature invariants, we develop the cosmological reconstruction scheme, which is explicitly applied to obtain the (effective quintessence/cosmological constant/phantom) accelerating expansion for the models under discussion. Late-time dynamics is analyzed and it is shown that the future universe may end in one of the four known finite-time singularities. For instance, the effective phantom superacceleration brings the universe to the famous Big Rip (or Type I) singularity [1]. In order to avoid the future singularity one has to add the extra higher-derivative gravitational terms, which are relevant at the early universe and help in the unification of early-time inflation with late-time acceleration because such terms induce inflation.

It should be noted that, as a rule, modified gravity equations of motion are higher-derivative differential equations. Thus, we consider only the background evolution of such theories in this report; we do not discuss the cosmological perturbation theory. The reason for this is very simple. Actually, in almost all existing approaches to cosmological perturbations in modified gravity, the corresponding equations are reduced to second-order differential equations. Clearly, this is not a well-justified approach, and the results obtained in this way are questionable. The new higher-derivative covariant perturbation theory should be developed to obtain correct answers to the related circle of questions.

The work is organized as follows. The second chapter gives the introduction to several popular models of modified gravity: $F(R)$ -theory, modified Gauss-Bonnet and string-inspired gravity, non-local gravity, non-minimally coupled theories, $F(R)$ Hořava-Lifshitz gravity, and power-counting renormalizable covariant gravity. The general properties of $F(R)$ -theory are discussed in section II A. Its different representations, namely Jordan frame, scalar-tensor form (Einstein frame) and fluid representation, are given. The thermodynamics of Schwarzschild-(anti)-de Sitter space are briefly discussed. Some local tests, in particular, the Newtonian regime and matter instability, are investigated. Several specific models (i.e., theory with negative and positive curvature power terms, $\ln R$ -model, and exponential theory), are presented. The spatially flat FRW cosmology is investigated. Viable modified gravities that allow the unification of the early-time inflation with late-time acceleration are given (for the first work where such a unification was proposed, see [2]). Note that the literature on $F(R)$ -gravity is vast, and the earlier reviews on this theory are given in refs. [3, 4]. Different aspects of $F(R)$ -gravity are discussed in refs. [5–7], and various exact solutions are presented in refs. [8]. The comparison with observational data and local tests is made in refs. [3, 4, 9]. As is indicated in this work, there are a number of viable $F(R)$ models that successfully pass local tests and are in accordance with observational data.

Modified Gauss-Bonnet gravity is considered in section II B, and its spatially flat FRW equations are given. The de Sitter solution and its thermodynamics are also studied, and the possibility of the unification of inflation with dark energy is mentioned. Section II C is devoted to a brief study of string-inspired (scalar-Gauss-Bonnet) gravity. The emergence of an accelerating dark energy epoch in such a model is proposed. The non-local generalization of $F(R)$ -gravity and Gauss-Bonnet gravity is discussed in section II D. We find the scalar-tensor representation for such non-local gravities and investigate their spatially flat FRW cosmology. The possibility of early-time or late-time acceleration is again demonstrated. Non-minimally coupled theories are presented in section II E. We consider the theories where scalar, vector or Yang-Mills Lagrangian is multiplied to function $F(R)$. FRW equations are given and possibility of cosmic acceleration is mentioned. In section II F we propose modified $F(R)$ Hořava-Lifshitz gravity. The original Hořava-Lifshitz theory, based on general relativity, has been introduced as a candidate for renormalizable quantum gravity at the price of the violation of the Lorentz invariance. The $F(R)$ generalization of Hořava-Lifshitz gravity yields consistent FRW equations that coincide with equations associated with the conventional theory by choosing the parameters to be specific values. We study several explicit models that are introduced in section II A in this framework and show that they may again predict the unified description of early-time inflation with dark energy. Motivated by the Hořava-Lifshitz proposal, the covariant power-counting renormalizable gravity is introduced in section II G. Two formulations of the theory: one utilizing an extra perfect fluid and the other using the Lagrange multiplier constraint, are given. Accelerating FRW cosmology is briefly discussed. In section II D, it is shown that arbitrary modified gravity may be interpreted as an effective fluid with an inhomogeneous equation of state. The late-time evolution of a specific dark fluid with a power-law equation of state is investigated. It is demonstrated that an effective quintessence/phantom fluid may drive the universe to a finite-time future singularity.

The background evolution of modified gravity is studied in the third chapter. Usually, in theories like general relativity, we start from a theory that is defined by the action and solve the corresponding equations of motion to define the background dynamics. However, an approach is already too complicated for simple models of modified gravity. Thus, using the fact that modified gravity is defined up to some arbitrary function of curvature invariants, the inverse problem is analyzed. The corresponding cosmological reconstruction scheme is proposed. Within the framework of this scheme, we show how the complicated background cosmology, which complies with observational data, may be reconstructed. This provides way to define (at least partially) the reconstructed form of modified gravity, which has a particular cosmological solution. Section III A is devoted to a formulation of cosmological reconstruction (in terms of cosmological time or in terms of e-folding N) for scalar-tensor gravity. Several examples of a (super)accelerating epoch are used for the reconstruction of scalar potentials. This formulation is extended for the theory with two scalars where the reconstruction of the transition from matter dominance to late-time acceleration as well as a reconstruction of the Λ CDM epoch is done. In section III B, we study the reconstruction in the k-essence model, using both formulations. Cosmological reconstruction for the $F(R)$ -model, using its presentation with an auxiliary scalar, is developed in section III C. The differential equation that provides the form of function $F(R)$ for a given spatially flat FRW cosmology is obtained. The Λ CDM epoch and dark era with a phantom divide crossing are reconstructed in $F(R)$ -gravity with/without matter in great detail. The general condition for the stability of the cosmological solution is derived. $F(R)$ -gravity with a Lagrange multiplier constraint is formulated, and its FRW cosmology is studied. In sections III D and III E, we propose the reconstruction of modified Gauss-Bonnet and scalar-Einstein-Gauss-Bonnet gravities, respectively. In section III F, we show that the reconstruction is easily generalized for the case of $F(R)$ Hořava-Lifshitz gravity. The explicit reconstruction of the Λ CDM epoch and of phantom superacceleration is fulfilled. The cosmological reconstruction of the power-law FRW solution for non-minimal Yang-Mills theory is given in section III G.

The fourth chapter is devoted to an investigation of late-time dynamics for effective quintessence/phantom dark energy. It is demonstrated that, regardless of whether the dark energy model utilizes fluid, scalar or alternative

gravity, the future universe may evolve to a finite-time singularity of one of four known types. The prescription to resolve such a future singularity via the addition of a higher-derivative gravitational term is proposed. In section IV A, we consider dark fluid coupled with dark matter. For a power-law equation of state, it is shown that the future dark era may be singular, depending on the parameter choice. The coupling with dark matter may prevent some of the soft future singularities. However, such coupling does not resolve the future singularity in a general case. Only the addition of the R^2 -term may cure all types of future singularities in the model under consideration. In section IV B, using the reconstruction method, we show that $F(R)$ gravity may have late-time accelerating cosmological solutions with all four types of finite-time future singularities. To avoid this, one should additionally modify the theory by the R^2 term. It is interesting that the use of such a term may be motivated by conformal anomaly considerations. The same problem with qualitatively similar results is studied for modified Gauss-Bonnet and $F(R)$ Hořava-Lifshitz gravities in section IV C and IV D. It is remarkable that, in the last case, one may need other higher-derivative gravitational terms to resolve all types of future singularities.

A summary and outlook are given in the Discussion section.

II. MODIFIED GRAVITY UNIFYING THE EARLY-TIME INFLATION WITH LATE-TIME ACCELERATION

This chapter is devoted to the general review of many popular models of modified gravity such as $F(R)$ gravity, modified Gauss-Bonnet gravity, string-inspired theory, non-local gravity and non-minimal theories. Their general properties, different representations and background spatially flat FRW cosmologies are investigated. Special attention is paid to possible unification of early-time inflation with late-time acceleration within such theories. The extension of such a study to power-counting renormalizable gravity with explicit or apparent Lorentz symmetry breaking is also made.

A. $F(R)$ gravity

In this section, we give a review of general properties of $F(R)$ gravity. Its scalar-tensor description is presented, and some constant curvature solutions are discussed. A number of popular models are introduced, and their applications for a unified description of inflation with dark energy are studied.

1. General properties

Let us start with the general introduction of the model. In $F(R)$ gravity [3], the scalar curvature R in the Einstein-Hilbert action¹

$$S_{\text{EH}} = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} + \mathcal{L}_{\text{matter}} \right), \quad (\text{II.1})$$

is replaced by an appropriate function of the scalar curvature:

$$S_{F(R)} = \int d^4x \sqrt{-g} \left(\frac{F(R)}{2\kappa^2} + \mathcal{L}_{\text{matter}} \right). \quad (\text{II.2})$$

Let us review the general properties of $F(R)$ gravity. For $F(R)$ theory, as for any other modified gravity theory, one can define an effective EoS parameter using its fluid representation. The FRW equations in the Einstein gravity coupled with perfect fluid are:

$$\rho_{\text{matter}} = \frac{3}{\kappa^2} H^2, \quad p_{\text{matter}} = -\frac{1}{\kappa^2} (3H^2 + 2\dot{H}). \quad (\text{II.3})$$

¹ We use the following convention for the curvatures:

$$R = g^{\mu\nu} R_{\mu\nu}, \quad R_{\mu\nu} = R^\lambda_{\mu\lambda\nu}, \quad R^\lambda_{\mu\rho\nu} = -\Gamma^\lambda_{\mu\rho,\nu} + \Gamma^\lambda_{\mu\nu,\rho} - \Gamma^\eta_{\mu\rho} \Gamma^\lambda_{\nu\eta} + \Gamma^\eta_{\mu\nu} \Gamma^\lambda_{\rho\eta}, \quad \Gamma^\eta_{\mu\lambda} = \frac{1}{2} g^{\eta\nu} (g_{\mu\nu,\lambda} + g_{\lambda\nu,\mu} - g_{\mu\lambda,\nu}).$$

For modified gravity, one may define an effective equation of state (EoS) parameter as follows:

$$w_{\text{eff}} = -1 - \frac{2\dot{H}}{3H^2}. \quad (\text{II.4})$$

This relation is very useful for a generalized fluid description of modified gravity.

The equation of motion for modified gravity is given by

$$\frac{1}{2}g_{\mu\nu}F(R) - R_{\mu\nu}F'(R) - g_{\mu\nu}\square F'(R) + \nabla_\mu \nabla_\nu F'(R) = -\frac{\kappa^2}{2}T_{\text{matter}\mu\nu}. \quad (\text{II.5})$$

Assuming a spatially flat FRW universe,

$$ds^2 = -dt^2 + a(t)^2 \sum_{i=1,2,3} (dx^i)^2, \quad (\text{II.6})$$

the FRW like equations are as follows:

$$0 = -\frac{F(R)}{2} + 3(H^2 + \dot{H})F'(R) - 18(4H^2\dot{H} + H\ddot{H})F''(R) + \kappa^2\rho_{\text{matter}}, \quad (\text{II.7})$$

$$0 = \frac{F(R)}{2} - (\dot{H} + 3H^2)F'(R) + 6(8H^2\dot{H} + 4\dot{H}^2 + 6H\ddot{H} + \ddot{H})F''(R) + 36(4H\dot{H} + \ddot{H})^2F'''(R) + \kappa^2 p_{\text{matter}}. \quad (\text{II.8})$$

Here, the Hubble rate H is defined by $H = \dot{a}/a$ and the scalar curvature R is given by $R = 12H^2 + 6\dot{H}$.

Several (often exact) solutions of (II.7) maybe found. However, due to the complicated nature of field equations, the number of exact solutions is much less than, for instance, that in general relativity. Without any matter, assuming that the Ricci tensor is covariantly constant, that is, $R_{\mu\nu} \propto g_{\mu\nu}$, Eq. (II.5) reduces to the algebraic equation:

$$0 = 2F(R) - RF'(R). \quad (\text{II.9})$$

If Eq. (II.9) has a solution, the (anti-)de Sitter and/or Schwarzschild- (anti-)de Sitter space

$$ds^2 = -\left(1 - \frac{2MG}{r} \mp \frac{r^2}{L^2}\right) dt^2 + \left(1 - \frac{2MG}{r} \mp \frac{r^2}{L^2}\right)^{-1} dr^2 + r^2 d\Omega^2, \quad (\text{II.10})$$

or Kerr - (anti-)de Sitter space is an exact vacuum solution. In (II.10), the minus and plus signs in $\pm$ correspond to the de Sitter and anti-de Sitter space, respectively. M is the mass of the black hole, $G = \frac{\kappa^2}{8\pi}$, and L is the length parameter of (anti-)de Sitter space, which is related to the curvature $R = \pm \frac{12}{L^2}$ (the plus sign for the de Sitter space and the minus sign for the anti-de Sitter space). For example, if we consider the model

$$F(R) = R - \frac{\mu^{2m+2}}{R^m}, \quad (\text{II.11})$$

with a constant μ with a mass dimension, the solution of (II.9) is given by

$$R = (2 + m)^{\frac{1}{m+1}} \mu^2. \quad (\text{II.12})$$

As another example, for a logarithmic model

$$F(R) = \mu^2 \ln \frac{R}{R_0}, \quad (\text{II.13})$$

with a constant R_0 , one gets

$$R = R_0 e^{\frac{1}{\mu^2}}. \quad (\text{II.14})$$

The entropy of the Schwarzschild - (anti-)de Sitter space can be evaluated by the WKB approximation for the Euclidean partition function Z [10]:

$$Z \sim e^{S_{\text{Sds}}}. \quad (\text{II.15})$$

Here, S_{Sds} is the value of the classical action calculated in the Schwarzschild - (anti-)de Sitter space. Because the value of S_{Sds} diverges, we regularize the action by subtracting the value when the solution of the (anti-)de Sitter space without a black hole is substituted [10]. The free energy $\mathcal{F}$ is given by

$$\mathcal{F} = -\frac{1}{\beta} \ln Z. \quad (\text{II.16})$$

And, using the standard thermodynamics relations,

$$\mathcal{S} = -\frac{d\mathcal{F}}{dT_H}, \quad (\text{II.17})$$

one can evaluate the entropy $\mathcal{S}$. Here, T_H is the Hawking temperature. Then, it follows that

$$\mathcal{S} = \frac{F'(R)A_H}{4G}. \quad (\text{II.18})$$

Here, A_H expresses the area of the horizon.

Let us now assume that $F(R)$ behaves as $F(R) \propto f_0 R^m$. Eq. (II.7) gives

$$0 = f_0 \left\{ -\frac{1}{2} \left(6\dot{H} + 12H^2 \right)^m + 3m \left(\dot{H} + H^2 \right) \left(6\dot{H} + 12H^2 \right)^{m-1} - 3mH \frac{d}{dt} \left\{ \left(6\dot{H} + 12H^2 \right)^{m-1} \right\} \right\} + \kappa^2 \rho_0 a^{-3(1+w)}. \quad (\text{II.19})$$

Eq. (II.8) is irrelevant because it can be derived from (II.19). When the contribution from the matter can be neglected ($\rho_0 = 0$), the following solution appears:

$$H \sim \frac{-\frac{(m-1)(2m-1)}{m-2}}{t}, \quad (\text{II.20})$$

with a corresponding EoS parameter (II.4):

$$w_{\text{eff}} = -\frac{6m^2 - 7m - 1}{3(m-1)(2m-1)}. \quad (\text{II.21})$$

On the other hand, when the matter contribution is include with a constant EoS parameter w , an exact solution of (II.19) is given by

$$a = a_0 t^{h_0}, \quad h_0 \equiv \frac{2m}{3(1+w)},$$

$$a_0 \equiv \left[-\frac{3f_0 h_0}{\kappa^2 \rho_0} (-6h_0 + 12h_0^2)^{m-1} \{ (1-2m)(1-m) - (2-m)h_0 \} \right]^{-\frac{1}{3(1+w)}}, \quad (\text{II.22})$$

and the effective EoS parameter (II.4) is

$$w_{\text{eff}} = -1 + \frac{w+1}{m}. \quad (\text{II.23})$$

These solutions (II.20) and (II.22) show that modified gravity may describe early/late-time universe acceleration. Moreover, it is very natural to propose that a more complicated modified gravity from the above class may give the unified description for inflation with late-time acceleration.

2. Scalar-tensor description

Let us now introduce perfect fluid and scalar-tensor representations for the modified gravity under consideration. These mathematically equivalent representations may simplify the investigation of a number of problems.

Eqs. (II.7) and (II.8) show that the effective energy density ρ_{eff} and the effective pressure p_{eff} including the contribution from $f(R)$ are given by (see, for instance, [11])

$$\rho_{\text{eff}} = \frac{1}{\kappa^2} \left(-\frac{1}{2} f(R) + 3 \left(H^2 + \dot{H} \right) f'(R) - 18 \left(4H^2 \dot{H} + H \ddot{H} \right) f''(R) \right) + \rho_{\text{matter}}, \quad (\text{II.24})$$

$$p_{\text{eff}} = \frac{1}{\kappa^2} \left(\frac{1}{2} f(R) - \left(3H^2 + \dot{H} \right) f'(R) + 6 \left(8H^2 \dot{H} + 4\dot{H}^2 + 6H \ddot{H} + \ddot{H} \right) f''(R) + 36 \left(4H \dot{H} + \ddot{H} \right)^2 f'''(R) \right) + p_{\text{matter}}. \quad (\text{II.25})$$

The equations (II.7) and (II.8) can be rewritten as in the Einstein gravity case (II.3):

$$\rho_{\text{eff}} = \frac{3}{\kappa^2} H^2, \quad p_{\text{eff}} = -\frac{1}{\kappa^2} (3H^2 + 2\dot{H}). \quad (\text{II.26})$$

Such a mathematically equivalent fluid representation for the FRW equations in modified gravity may lead to a number of assumptions that are not well justified. Indeed, the generalized gravitational fluid contains higher-derivative curvature invariants. This fact is often ignored in the study of cosmological perturbations in modified gravity when such a theory is developed in strict similarity with general relativity, such that corresponding differential equations become of second order (see, for instance, [12]). Indeed, first of all, the field equations of the theory under investigation are of fourth order. It is evident that cosmological perturbation equations should be of fourth order as well. Moreover, such equations should be covariant. The corresponding covariant and higher-derivative cosmological perturbation theory (for review, see [13, 14]) is developed, but it is extremely complicated.

One can also rewrite $F(R)$ gravity in the scalar-tensor form. By introducing the auxiliary field A , the action (II.2) of the $F(R)$ gravity is rewritten in the following form:

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \{ F'(A) (R - A) + F(A) \}. \quad (\text{II.27})$$

By the variation of A , one obtains $A = R$. Substituting $A = R$ into the action (II.27), one can reproduce the action in (II.2). Furthermore, we rescale the metric in the following way (conformal transformation):

$$g_{\mu\nu} \rightarrow e^\sigma g_{\mu\nu}, \quad \sigma = -\ln F'(A). \quad (\text{II.28})$$

Thus, the Einstein frame action is obtained:

$$S_E = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \left(R - \frac{3}{2} g^{\rho\sigma} \partial_\rho \sigma \partial_\sigma \sigma - V(\sigma) \right),$$

$$V(\sigma) = e^\sigma g(e^{-\sigma}) - e^{2\sigma} f(g(e^{-\sigma})) = \frac{A}{F'(A)} - \frac{F(A)}{F'(A)^2}. \quad (\text{II.29})$$

Here $g(e^{-\sigma})$ is given by solving the equation $\sigma = -\ln(1 + f'(A)) = -\ln F'(A)$ as $A = g(e^{-\sigma})$. Due to the scale transformation (II.28), a coupling of the scalar field σ with usual matter arises. The mass of σ is given by

$$m_\sigma^2 \equiv \frac{3}{2} \frac{d^2 V(\sigma)}{d\sigma^2} = \frac{3}{2} \left\{ \frac{A}{F'(A)} - \frac{4F(A)}{(F'(A))^2} + \frac{1}{F''(A)} \right\}. \quad (\text{II.30})$$

Unless m_σ is very large, the large correction to Newton's law appears. Naively, one expects the order of the mass m_σ to be that of the Hubble rate, that is, $m_\sigma \sim H \sim 10^{-33} \text{ eV}$, which is very light and could make the correction very large.

In [15], a “realistic” $F(R)$ model was proposed. It has been found, however, that the model has an instability where the large curvature can be easily produced (manifestation of a possible future singularity). In the model of [15], a parameter $m \sim 10^{-33} \text{ eV}$ with a mass dimension is included. The parameter m plays a role of the effective cosmological constant. When the curvature R is large enough compared with m^2 , $R \gg m^2$, $F(R)$ [15] looks as follows:

$$F(R) = R - c_1 m^2 + \frac{c_2 m^{2n+2}}{R^n} + \mathcal{O}(R^{-2n}). \quad (\text{II.31})$$

Here c_1 , c_2 , and n are positive dimensionless constants. The potential $V(\sigma)$ (II.29) has the following asymptotic form:

$$V(\sigma) \sim \frac{c_1 m^2}{A^2}. \quad (\text{II.32})$$

Then, the infinite curvature $R = A \rightarrow \infty$ corresponds to a small value of the potential and, therefore, the large curvature can be easily produced.

Let us assume that when R is large, $F(R)$ behaves as

$$F(R) \sim F_0 R^\epsilon. \quad (\text{II.33})$$

Here, F_0 and ϵ are positive constants. We also assume $\epsilon > 1$ so that this term dominates compared with general relativity. Then, the potential $V(\sigma)$ (II.29) behaves as

$$V(\sigma) \sim \frac{\epsilon - 1}{\epsilon^2 F_0 R^{\epsilon-2}}. \quad (\text{II.34})$$

Therefore, if $1 < \epsilon < 2$, the potential $V(\sigma)$ diverges when $R \rightarrow \infty$ and, therefore, the large curvature is not realized so easily. When $\epsilon = 2$, $V(\sigma)$ takes a finite value $1/F_0$ when $R \rightarrow \infty$. As long as $1/F_0$ is large enough, the large curvature can be prevented.

Note that the anti-gravity regime appears when $F'(R)$ is negative, which follows from Eq. (II.27) [2]. Then, we need to require

$$F'(R) > 0. \quad (\text{II.35})$$

We should also note that

$$\frac{dV(\sigma)}{dA} = \frac{F''(A)}{F'(A)^3} (-AF'(A) + 2F(A)). \quad (\text{II.36})$$

Therefore, if

$$0 = -AF'(A) + 2F(A), \quad (\text{II.37})$$

the scalar field σ is on the local maximum or local minimum of the potential and, therefore, σ can be a constant. Note that the condition (II.37) is nothing but the condition (II.9) for the existence of the de Sitter solution. When the condition (II.37) is satisfied, the mass (II.30) can be rewritten as

$$m_\sigma^2 = \frac{3}{2F'(A)} \left(-A + \frac{F'(A)}{F''(A)} \right). \quad (\text{II.38})$$

Then, when the condition (II.35) for the non-existence of the anti-gravity is satisfied, the mass squared m_σ^2 is positive and, therefore the, scalar field is on the local minimum if

$$-A + \frac{F'(A)}{F''(A)} > 0. \quad (\text{II.39})$$

On the other hand, if

$$-A + \frac{F'(A)}{F''(A)} < 0, \quad (\text{II.40})$$

the scalar field is on the local maximum of the potential and the mass squared m_σ^2 is negative. As we will see later, the condition (II.39) is nothing but the condition for stability of the de Sitter space.

We have rewritten the action (II.2) of $F(R)$ gravity in a scalar-tensor form (II.29). Inversely, it is always possible to rewrite the action of the scalar-tensor theory

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} R - \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - \tilde{V}(\varphi) \right\}, \quad (\text{II.41})$$

as the action of $F(R)$ gravity [16]. By comparing (II.41) with (II.29), we find

$$\varphi = \frac{\sqrt{3}}{\kappa} \sigma, \quad \tilde{V}(\varphi) = V\left(\frac{\kappa}{\sqrt{3}} \varphi\right). \quad (\text{II.42})$$

One now uses the conformal transformation

$$g_{\mu\nu} \rightarrow e^{\pm \kappa \varphi \sqrt{\frac{2}{3}}} g_{\mu\nu}, \quad (\text{II.43})$$

to make the kinetic term of φ vanish. Then, we obtain

$$S = \int d^4x \sqrt{-g} \left\{ \frac{e^{\pm \kappa \varphi \sqrt{\frac{2}{3}}}}{2\kappa^2} R - e^{\pm 2\kappa \varphi \sqrt{\frac{2}{3}}} \tilde{V}(\varphi) \right\}. \quad (\text{II.44})$$

The action (II.44) is often called the ‘‘Jordan frame action’’, whereas the action (II.41) is the ‘‘Einstein frame action’’ due to either the non-minimal coupling or the standard coupling in front of the scalar curvature. Because φ now

becomes an auxiliary field, which does not have a kinetic term, one may delete φ by using the corresponding equation of motion:

$$R = e^{\pm\kappa\varphi\sqrt{\frac{2}{3}}} \left(4\kappa^2 \tilde{V}(\varphi) \pm 2\kappa\sqrt{\frac{3}{2}} \tilde{V}'(\varphi) \right), \quad (\text{II.45})$$

which can be solved with respect to φ as a function of R as $\varphi = \varphi(R)$. Thus, we can rewrite the action (II.44) in the form of $F(R)$ gravity:

$$S = \int d^4x \sqrt{-g} F(R), \quad F(R) \equiv \frac{e^{\pm\kappa\varphi(R)\sqrt{\frac{2}{3}}}}{2\kappa^2} R - e^{\pm 2\kappa\varphi(R)\sqrt{\frac{2}{3}}} \tilde{V}(\varphi(R)). \quad (\text{II.46})$$

Note that one can rewrite the scalar-tensor theory (II.41) only when the sign in front of the kinetic term is negative, that is, when the scalar field is canonical. One cannot rewrite the action if the sign is positive, as

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} R + \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - \tilde{V}(\varphi) \right\}, \quad (\text{II.47})$$

which corresponds to the phantom scalar [17].

Note that, for the metric transformed as in (II.43), even if the Einstein frame universe is in a non-phantom phase, where the effective EoS w_{eff} in (II.4) is larger than -1 , the Jordan frame universe can be, in general, in a phantom phase. In other words, despite the mathematical equivalence between two frames there occurs some kind of physical non-equivalence [18]. More precisely, if physics in one frame is matched with observations and describes the observable accelerating universe, then another frame of physics becomes extremely unconventional. For instance, instead of the acceleration, the observer sees the deceleration, and matter couples with scalar, etc.

3. Matter instability

Let us discuss the matter instability issue [19] in modified gravity. It is related to the fact that the spherical body solution in general relativity may not be the solution in modified gravity theory. The matter instability may appear when the energy density or the curvature is large compared with the average density or curvature in the universe, as is the case inside of a planet. Multiplying $g^{\mu\nu}$ with Eq. (II.5), one obtains

$$\square R + \frac{F^{(3)}(R)}{F^{(2)}(R)} \nabla_\rho R \nabla^\rho R + \frac{F'(R)R}{3F^{(2)}(R)} - \frac{2F(R)}{3F^{(2)}(R)} = \frac{\kappa^2}{6F^{(2)}(R)} T_{\text{matter}}. \quad (\text{II.48})$$

Here, T_{matter} is the trace of the matter energy-momentum tensor: $T_{\text{matter}} \equiv T_{\text{matter}\rho}{}^\rho$. We also denote $d^n F(R)/dR^n$ by $F^{(n)}(R)$. Let us now consider the perturbation from the solution of general relativity. We denote the scalar curvature solution given by the matter density in the Einstein gravity by $R_b \sim (\kappa^2/2)\rho_{\text{matter}} > 0$ and separate the scalar curvature R into the sum of R_b and the perturbed part R_p as $R = R_b + R_p$ ($|R_p| \ll |R_b|$). Then, Eq. (II.48) leads to the following perturbed equation:

$$\begin{aligned} 0 = & \square R_b + \frac{F^{(3)}(R_b)}{F^{(2)}(R_b)} \nabla_\rho R_b \nabla^\rho R_b + \frac{F'(R_b)R_b}{3F^{(2)}(R_b)} \\ & - \frac{2F(R_b)}{3F^{(2)}(R_b)} - \frac{R_b}{3F^{(2)}(R_b)} + \square R_p + 2 \frac{F^{(3)}(R_b)}{F^{(2)}(R_b)} \nabla_\rho R_b \nabla^\rho R_p + U(R_b)R_p. \end{aligned} \quad (\text{II.49})$$

Here, $U(R_b)$ is given by

$$\begin{aligned} U(R_b) \equiv & \left(\frac{F^{(4)}(R_b)}{F^{(2)}(R_b)} - \frac{F^{(3)}(R_b)^2}{F^{(2)}(R_b)^2} \right) \nabla_\rho R_b \nabla^\rho R_b + \frac{R_b}{3} \\ & - \frac{F^{(1)}(R_b)F^{(3)}(R_b)R_b}{3F^{(2)}(R_b)^2} - \frac{F^{(1)}(R_b)}{3F^{(2)}(R_b)} + \frac{2F(R_b)F^{(3)}(R_b)}{3F^{(2)}(R_b)^2} - \frac{F^{(3)}(R_b)R_b}{3F^{(2)}(R_b)^2}. \end{aligned} \quad (\text{II.50})$$

It is convenient to consider the case that R_b and R_p are uniform; that is, they do not depend on the spatial coordinates. Thus, the d'Alembertian can be replaced with the second derivative with respect to the time coordinate: $\square R_p \rightarrow -\partial_t^2 R_p$ and Eq. (II.50) has the following structure:

$$0 = -\partial_t^2 R_p + U(R_b)R_p + \text{const.} \quad (\text{II.51})$$

Then, if $U(R_b) > 0$, R_p becomes exponentially large with time t : $R_p \sim e^{\sqrt{U(R_b)}t}$ and the system is unstable. In the $1/R$ model [20–22], because the order of the mass parameter m_μ is

$$\mu^{-1} \sim 10^{18} \text{sec} \sim (10^{-33} \text{eV})^{-1}, \quad (\text{II.52})$$

one finds

$$U(R_b) = -R_b + \frac{R_b^3}{6\mu^4} \sim \frac{R_0^3}{\mu^4} \sim (10^{-26} \text{sec})^{-2} \left(\frac{\rho_{\text{matter}}}{\text{g cm}^{-3}} \right)^3, \quad R_b \sim (10^3 \text{sec})^{-2} \left(\frac{\rho_{\text{matter}}}{\text{g cm}^{-3}} \right). \quad (\text{II.53})$$

Thus, the model is unstable and it presumably decays in 10^{-26} sec (for a planet-sized object). On the other hand, in a $1/R + R^2$ model [2], one gets

$$U(R_0) \sim \frac{R_0}{3} > 0. \quad (\text{II.54})$$

Then, the system could be unstable again, but the decay time is $\sim 1,000$ sec, that is, macroscopic. For the model [15], $U(R_b)$ is negative

$$U(R_0) \sim -\frac{(n+2)m^2c_2^2}{c_1n(n+1)} < 0. \quad (\text{II.55})$$

Therefore, there is no matter instability. Note that the models considered in the following sections successfully pass the matter instability test.

4. Unified theories without a flat space solution

We now review the $F(R)$ models which do not have a flat space solution because of the non-analytical behavior at zero curvature. Some of these such models may induce large corrections to Newton's law.

Modified gravity with negative and positive powers of the curvature

In [2], the following action was proposed:

$$F(R) = R - \frac{c}{(R - \Lambda_1)^n} + b(R - \Lambda_2)^m. \quad (\text{II.56})$$

Here n , m , c , and b are constants. In general, n and m may be fractional. Let us show that the above model leads to an acceptable cosmic speed-up and is consistent with some of the solar system tests.

For the action (II.56), Eq. (II.9) has the following form:

$$0 = R - \frac{c\{(n+2)R - 2\Lambda_1\}}{(R - \Lambda_1)^{n+1}} + b\{(2-m)R - 2\Lambda_2\}(R - \Lambda_2)^m. \quad (\text{II.57})$$

Especially when $n = 1$, $m = 2$, and $\Lambda_1 = 0$, one gets

$$R = R_\pm = \frac{b\Lambda_2 \pm \sqrt{b^2\Lambda_2^2 + 3c(1 + 2b\Lambda_2)}}{1 + 2b\Lambda_2}. \quad (\text{II.58})$$

If $b\Lambda_2 > 0$, $c < 0$, and $b^2\Lambda_2^2 + 3c(1 + 2b\Lambda_2) > 0$, both of the solutions express the de Sitter space. Thus, the natural possibility for the unification of early-time inflation with late-time acceleration appears. Here, higher-derivative terms act in favor of early-time inflation, and the $1/R$ term supports a cosmic speed-up.

We now neglect the contribution from matter. When $n = 1$, $m = 2$, and $\Lambda_1 = \Lambda_2 = 0$ in (II.56) and the curvature is small, we obtain the solution of (II.7): $\hat{a} \propto t^2$, which is consistent with the result in [20–22].

If the present universe expands with the above power-law, the curvature of the present universe should be small compared with that of the de Sitter universe solution in (II.58). As the Hubble rate in the present universe is 10^{-33} eV, the magnitude of the parameter c , which corresponds to μ^4 in [20–22], is on the order of $(10^{-33} \text{eV})^4$.

Let us consider the more general case that $F(R)$ is given by (II.56) when the curvature is small and $\Lambda_1 = 0$. Neglecting the contribution from the matter again, solving (II.7), we obtain $\hat{a} \propto t^{\frac{(n+1)(2n+1)}{n+2}}$ (see (II.20)). It is quite

remarkable that actually any negative power of the curvature supports the cosmic acceleration. This gives the freedom in modification of the model to achieve the better consistency with experimental tests of Newtonian gravity.

On the other hand, when the scalar curvature R is large, one obtains $\hat{a} \propto t^{-\frac{(m-1)(2m-1)}{m-2}}$ (see (II.20)). When $\frac{(m-1)(2m-1)}{m-2} > 0$, the universe is shrinking but with a change in the arrow of time by $t \rightarrow t_s - t$, the inflation occurs with the inverse power-law. At $t = t_s$, the size of the universe diverges. It is remarkable that when m is fractional (or irrational) and $1 < m < 2$, the expression of $\hat{a}$ is still valid and the power becomes positive, the universe evolves with the (fractional) power-law expansion.

It is interesting that the model in (II.56) is more consistent than refs. [20–22] (for a discussion of the Newtonian limit in $1/R$ theory, see [23]) as it may pass (at least, some) solar system tests. An example of the tests is matter instability as discussed in the previous subsection. Indeed, in [19], small gravitational object like the earth or the sun in the model [20–22] is considered. It has been shown that the system becomes unstable. This may cause an unacceptable force between distant galaxies [24, 25]. As shown in the previous subsection (see ref. [2]), however, by adding the positive power (higher than 1) of the scalar curvature to the action, the instability could be significantly improved. Furthermore, the account of quantum effects in modified gravity also acts against instability [26].

It has also been mentioned in [27] that the $1/R$ model [20–22], which is equivalent to scalar-tensor gravity, is ruled out as a realistic theory due to the constraints to such Brans-Dicke-type theories. In [2], it was shown that by adding the scalar curvature squared term to the action, the mass of the scalar field can be adjusted to be very large and the scalar field can decouple. Thus, the modified gravity theory (II.56) (after some fine-tuning [2]) passes most of the solar system tests. In addition, at precisely the same parameter values, the above modified gravity has Newtonian limit that does not deviate significantly from that in general relativity. Nevertheless, as has been mentioned at the very beginning, such theory has non-analytical behavior at zero (or some constant) curvature which may be phenomenologically unacceptable.

ln R gravity

Other gravitational alternatives for dark energy may be suggested along this same line. As an extension of the theory (II.56), one may consider the model [28] containing the logarithm of the scalar curvature R :

$$F(R) = R + \alpha' \ln \frac{R}{\mu^2} + \beta R^m. \quad (\text{II.59})$$

We should note that the $m = 2$ choice simplifies the model. Assuming the scalar curvature is constant and the Ricci tensor is also covariantly constant, the equations (II.9) are as follows:

$$0 = 2F(R) - RF'(R) = \tilde{F}(R) \equiv R + 2\alpha' \ln \frac{R}{\mu^2} - \alpha'. \quad (\text{II.60})$$

If $\alpha' > 0$, $\tilde{F}(R)$ is a monotonically increasing function and $\lim_{R \rightarrow 0} \tilde{F}(R) = -\infty$ and $\lim_{R \rightarrow +\infty} \tilde{F}(R) = +\infty$. There is one and only one solution of (II.60). This solution may correspond to the inflation. On the other hand, if $\alpha' < 0$, $\lim_{R \rightarrow 0} \tilde{F}(R) = \infty$ and $\lim_{R \rightarrow +\infty} \tilde{F}(R) = +\infty$. Because $\tilde{F}'(R) = 1 + \frac{\alpha'}{R}$, the minimum of $\tilde{F}(R)$, where $\tilde{F}'(R) = 0$, is given by $R = -2\alpha'$. If $\tilde{f}(-2\alpha') > 0$, there is no solution of (II.60). If $\tilde{f}(-2\alpha') = 0$, there is only one solution and if $\tilde{f}(-2\alpha') < 0$, there are two solutions. Because $\tilde{f}(-2\alpha') = -2\alpha' \left(1 - \ln \frac{-2\alpha'}{\mu^2}\right)$, there are two solutions if $-\frac{2\alpha'}{\mu^2} > e$. Because the square root of the curvature R corresponds to the rate of the expansion of the universe, the larger of the two solutions might correspond to the inflation in the early universe and the smaller one to the present accelerating universe.

We can consider the late FRW cosmology when the scalar curvature R is small. Solving (II.7), it follows that the power-law inflation could occur: $\hat{a} \propto t^{\frac{1}{2}}$.

One may discuss further generalizations like [28] (see also [29])

$$F(R) = R + \gamma R^{-n} \left(\ln \frac{R}{\mu^2} \right)^m. \quad (\text{II.61})$$

Here, n is restricted by $n > -1$ (m is an arbitrary integer) in order that the second term could be more dominant than the Einstein term when R is small. For this model, we find

$$\hat{a} \sim t^{\frac{(n+1)(2n+1)}{n+2}}. \quad (\text{II.62})$$

This does not depend on m . The logarithmic factor is almost irrelevant. The effective w_{eff} (II.4) is given by

$$w_{\text{eff}} = -\frac{6n^2 + 7n - 1}{3(n+1)(2n+1)}. \quad (\text{II.63})$$

Then, w_{eff} can be negative if

$$-1 < n < -\frac{1}{2} \text{ or } n > \frac{-7 + \sqrt{73}}{12} = 0.1287 \dots \quad (\text{II.64})$$

From (II.62), the condition that the universe could accelerate is given by $\frac{(n+1)(2n+1)}{n+2} > 1$, that is:

$$n > \frac{-1 + \sqrt{3}}{2} = 0.366 \dots \quad (\text{II.65})$$

Clearly, the effective EoS parameter w may be within the existing bounds. Thus, the logarithmic model may also propose the unified description of inflation with dark energy era.

5. Viable modified gravities

Using the previous arguments, let us review the viable models of gravity that could unify the accelerating expansion of the present universe and the inflation of the early universe.

The realistic $F(R)$ models unifying inflation with dark energy

In refs. [30–32], viable models of $F(R)$ gravity unifying the late-time acceleration and the inflation were proposed. To construct these models, we have required several conditions:

1. To generate the inflation, one requires

$$\lim_{R \rightarrow \infty} f(R) = -\Lambda_i. \quad (\text{II.66})$$

Here, Λ_i is an effective cosmological constant at the early universe; therefore, it is natural to assume $\Lambda_i \gg (10^{-33} \text{eV})^2$. For instance, it is natural to have $\Lambda_i \sim 10^{20 \sim 38} (\text{eV})^2$.

2. To enable the current cosmic acceleration to be generated, the current $f(R)$ gravity is considered to be a small constant, that is,

$$f(R_0) = -2\tilde{R}_0, \quad f'(R_0) \sim 0. \quad (\text{II.67})$$

Here, R_0 is the current curvature $R_0 \sim (10^{-33} \text{eV})^2$. Note that $R_0 > \tilde{R}_0$ due to the contribution from matter. In fact, if one can regard $f(R_0)$ as an effective cosmological constant, the effective Einstein equation gives $R_0 = \tilde{R}_0 - \kappa^2 T_{\text{matter}}$. Here, T_{matter} is the trace of the matter energy-momentum tensor. Note that $f'(R_0)$ need not vanish completely. Because we consider the time scale from one billion years to ten billion years, we only require $|f'(R_0)| \ll (10^{-33} \text{eV})^4$.

Instead of the model corresponding to (II.66), one may consider a model that satisfies

$$\lim_{R \rightarrow \infty} f(R) = \alpha R^m, \quad (\text{II.68})$$

with a positive integer $m > 1$ and a constant α . To avoid the anti-gravity $f'(R) > -1$, $\alpha > 0$ and, therefore, $f(R)$ should be positive at the early universe. On the other hand, Eq. (II.67) shows that $f(R)$ is negative at the present universe. Therefore, $f(R)$ should cross zero in the past.

3. The last condition is

$$\lim_{R \rightarrow 0} f(R) = 0, \quad (\text{II.69})$$

which means that there is a flat space-time solution.

One typical model proposed in ref. [30] that satisfies the above conditions is

$$f(R) = \frac{\alpha R^{2n} - \beta R^n}{1 + \gamma R^n}. \quad (\text{II.70})$$

Here, α , β , and γ are positive constants and n is a positive integer. Eq. (II.70) gives [30]

$$R_0 = \left\{ \left(\frac{1}{\gamma} \right) \left(1 + \sqrt{1 + \frac{\beta\gamma}{\alpha}} \right) \right\}^{1/n}, \quad (\text{II.71})$$

and, therefore,

$$f(R_0) \sim -2\tilde{R}_0 = \frac{\alpha}{\gamma^2} \left(1 + \frac{\left(1 - \frac{\beta\gamma}{\alpha} \right) \sqrt{1 + \frac{\beta\gamma}{\alpha}}}{2 + \sqrt{1 + \frac{\beta\gamma}{\alpha}}} \right). \quad (\text{II.72})$$

Then, it follows that

$$\alpha \sim 2\tilde{R}_0 R_0^{-2n}, \quad \beta \sim 4\tilde{R}_0^2 R_0^{-2n} R_I^{n-1}, \quad \gamma \sim 2\tilde{R}_0 R_0^{-2n} R_I^{n-1}. \quad (\text{II.73})$$

In the model (II.70), the correction to Newton's law is small because the mass m_σ (II.30) is large and is given by $m_\sigma^2 \sim 10^{-160+109n} \text{ eV}^2$ in the solar system and $m_\sigma^2 \sim 10^{-144+98n} \text{ eV}^2$ in the air on the earth [30]. In both cases, the mass m_σ is very large if $n \geq 2$.

Another model corresponding to (II.66) was proposed in [32]

$$f(R) = -\alpha_0 \left(\tanh \left(\frac{b_0(R-R_0)}{2} \right) + \tanh \left(\frac{b_0 R_0}{2} \right) \right) - \alpha_I \left(\tanh \left(\frac{b_I(R-R_I)}{2} \right) + \tanh \left(\frac{b_I R_I}{2} \right) \right). \quad (\text{II.74})$$

One now assumes

$$R_I \gg R_0, \quad \alpha_I \gg \alpha_0, \quad b_I \ll b_0, \quad (\text{II.75})$$

and

$$b_I R_I \gg 1. \quad (\text{II.76})$$

When $R \rightarrow 0$ or $R \ll R_0$, R_I , $f(R)$ behaves as

$$f(R) \rightarrow - \left(\frac{\alpha_0 b_0}{2 \cosh^2 \left(\frac{b_0 R_0}{2} \right)} + \frac{\alpha_I b_I}{2 \cosh^2 \left(\frac{b_I R_I}{2} \right)} \right) R. \quad (\text{II.77})$$

and $f(0) = 0$ again. When $R \gg R_I$, it follows that

$$f(R) \rightarrow -2\Lambda_I \equiv -\alpha_0 \left(1 + \tanh \left(\frac{b_0 R_0}{2} \right) \right) - \alpha_I \left(1 + \tanh \left(\frac{b_I R_I}{2} \right) \right) \sim -\alpha_I \left(1 + \tanh \left(\frac{b_I R_I}{2} \right) \right). \quad (\text{II.78})$$

On the other hand, when $R_0 \ll R \ll R_I$, one gets

$$f(R) \rightarrow -\alpha_0 \left[1 + \tanh \left(\frac{b_0 R_0}{2} \right) \right] - \frac{\alpha_I b_I R}{2 \cosh^2 \left(\frac{b_I R_I}{2} \right)} \sim -2\Lambda_0 \equiv -\alpha_0 \left[1 + \tanh \left(\frac{b_0 R_0}{2} \right) \right]. \quad (\text{II.79})$$

Here, we have assumed the condition (II.76). One also finds

$$f'(R) = -\frac{\alpha_0 b_0}{2 \cosh^2 \left(\frac{b_0(R-R_0)}{2} \right)} - \frac{\alpha_I b_I}{2 \cosh^2 \left(\frac{b_I(R-R_I)}{2} \right)}, \quad (\text{II.80})$$

which has two valleys when $R \sim R_0$ or $R \sim R_I$. When $R = R_0$,

$$f'(R_0) = -\alpha_0 b_0 - \frac{\alpha_I b_I}{2 \cosh^2 \left(\frac{b_I(R_0-R_I)}{2} \right)} > -\alpha_I b_I - \alpha_0 b_0. \quad (\text{II.81})$$

On the other hand, when $R = R_I$, it follows that

$$f'(R_I) = -\alpha_I b_I - \frac{\alpha_0 b_0}{2 \cosh^2 \left(\frac{b_0(R_0-R_I)}{2} \right)} > -\alpha_I b_I - \alpha_0 b_0. \quad (\text{II.82})$$

Due to the condition (II.35) required to avoid the anti-gravity regime, one obtains

$$\alpha_I b_I + \alpha_0 b_0 < 2. \quad (\text{II.83})$$

In the solar system domain, on or inside the earth, where $R \gg R_0$, $f(R)$ can be approximated by

$$f(R) \sim -2\Lambda_{\text{eff}} + 2\alpha e^{-b(R-R_0)}. \quad (\text{II.84})$$

On the other hand, because $R_0 \ll R \ll R_I$, by assuming Eq. (II.76), $f(R)$ (II.74) could also be approximated by

$$f(R) \sim -2\Lambda_0 + 2\alpha e^{-b_0(R-R_0)}, \quad (\text{II.85})$$

which has the same expression, after having identified $\Lambda_0 = \Lambda_{\text{eff}}$ and $b_0 = b$. Thus, one may check the case (II.84) only. The effective mass has the following form

$$m_\sigma^2 \sim \frac{e^{b(R-R_0)}}{4\alpha b^2}, \quad (\text{II.86})$$

which could be very large, that is, $m_\sigma^2 \sim 10^{1,000} \text{ eV}^2$ in the solar system and $m_\sigma^2 \sim 10^{10,000,000,000} \text{ eV}^2$ in the air surrounding the earth. Thus, the correction to Newton's law becomes negligible.

One may consider another model [30]:

$$f(R) = -\frac{(R-R_0)^{2k+1} + R_0^{2k+1}}{f_0 + f_1 \left\{ (R-R_0)^{2k+1} + R_0^{2k+1} \right\}}. \quad (\text{II.87})$$

It has been shown [30] that for $k \geq 10$, such modified gravity passes the local tests. It also unifies the early-time inflation with the dark energy epoch. In (II.87), R_0 is the current curvature $R_0 \sim (10^{-33} \text{ eV})^2$. We also require

$$f_0 \sim \frac{R_0^{2n}}{2}, \quad f_1 = \frac{1}{\Lambda_i}. \quad (\text{II.88})$$

Here Λ_i is the effective cosmological constant in the inflation epoch. When $R \gg \Lambda_i$, $f(R)$ (II.87) behaves as

$$f(R) \sim -\frac{1}{f_1} + \frac{f_0}{f_1^2 R^{2n+1}}. \quad (\text{II.89})$$

The trace equation, which is the trace part of (II.5), is as follows:

$$3\Box f'(R) = R + 2f(R) - Rf'(R) - \kappa^2 T_{\text{matter}}. \quad (\text{II.90})$$

Here, T_{matter} is the trace part of the matter energy-momentum tensor. For the FRW metric (II.6), one finds

$$R \sim (t_s - t)^{-2/(2n+3)}, \quad (\text{II.91})$$

which diverges at a finite future time $t = t_s$. By a similar analysis, we can show that if $f(R)$ behaves as $f(R) \sim R^\epsilon$ for large R with a constant ϵ , a future singularity appears if $\epsilon > 2$ or $\epsilon < 0$, which is consistent with the analysis (II.34) via the scalar-tensor form of the action. Conversely if $2 \geq \epsilon \geq 0$, the singularity does not appear. Thus, adding the term $R^2 \tilde{f}(R)$, where $\lim_{R \rightarrow 0} \tilde{f}(R) = c_1$, $\lim_{R \rightarrow \infty} \tilde{f}(R) = c_2$, to $f(R)$ (II.87), the future singularity (II.91) disappears. More details about future singularities in modified gravity will be given in the fourth chapter.

Let us investigate the above situation in more detail. Assume

$$f(R) \sim F_0 + F_1 R^\epsilon, \quad (\text{II.92})$$

when R is large. Here F_0 and F_1 are constants where F_0 may vanish but $F_1 \neq 0$. In the case of (II.89)

$$F_0 = -\frac{1}{f_1}, \quad F_1 = \frac{f_0}{f_1^2}, \quad \epsilon = -(2n+1). \quad (\text{II.93})$$

Under the assumption (II.92), the trace equation (II.90) gives

$$3F_1 \Box R^{\epsilon-1} = \begin{cases} R & \text{when } \epsilon < 0 \text{ or } \epsilon = 2 \\ (2-\epsilon) F_1 R^\epsilon & \text{when } \epsilon > 1 \text{ or } \epsilon \neq 2 \end{cases}. \quad (\text{II.94})$$

In the FRW background (II.6), when the Hubble rate has a singularity as

$$H \sim \frac{h_0}{(t_s - t)^\beta}, \quad (\text{II.95})$$

with constants h_0 and β , the scalar curvature $R = 6\dot{H} + 12H^2$ behaves as

$$R \sim \begin{cases} \frac{12h_0^2}{(t_s - t)^{2\beta}} & \text{when } \beta > 1 \\ \frac{6h_0 + 12h_0^2}{(t_s - t)^2} & \text{when } \beta = 1 \\ \frac{6\beta h_0}{(t_s - t)^{\beta+1}} & \text{when } \beta < 1 \end{cases}. \quad (\text{II.96})$$

In (II.95) or (II.96), the $\beta \geq 1$ case corresponds to Type I (Big Rip) singularity in [1, 33], $1 > \beta > 0$ to Type III, $0 > \beta > -1$ to Type II, and $\beta < -1$ (but $\beta \neq \text{integer}$) to Type IV.

The classification of the finite-time future singularities used above is given in ref. [34]:

- Type I (“Big Rip”) : For $t \rightarrow t_s$, $a \rightarrow \infty$, $\rho_{\text{eff}} \rightarrow \infty$ and $|p_{\text{eff}}| \rightarrow \infty$. This also includes the case of ρ_{eff} , p_{eff} being finite at t_s . This singularity has been discovered in ref. [1]. Its manifestations in different models have been studied in refs. [34].
- Type II (“sudden”) [35]: For $t \rightarrow t_s$, $a \rightarrow a_s$, $\rho_{\text{eff}} \rightarrow \rho_s$ and $|p_{\text{eff}}| \rightarrow \infty$.
- Type III : For $t \rightarrow t_s$, $a \rightarrow a_s$, $\rho_{\text{eff}} \rightarrow \infty$ and $|p_{\text{eff}}| \rightarrow \infty$.
- Type IV : For $t \rightarrow t_s$, $a \rightarrow a_s$, $\rho_{\text{eff}} \rightarrow 0$, $|p_{\text{eff}}| \rightarrow 0$ and higher derivatives of H diverge. This also includes the case in which p_{eff} (ρ_{eff}) or both of p_{eff} and ρ_{eff} tend to some finite values, whereas higher derivatives of H diverge.

Here, ρ_{eff} and p_{eff} are defined by

$$\rho_{\text{eff}} \equiv \frac{3}{\kappa^2} H^2, \quad p_{\text{eff}} \equiv -\frac{1}{\kappa^2} (2\dot{H} + 3H^2). \quad (\text{II.97})$$

Substituting (II.96) into (II.94), one finds that there are two classes of consistent solutions. The first solution is specified by $\beta = 1$ and $\epsilon > 1$ (but $\epsilon \neq 2$) case, which corresponds to the Big Rip ($h_0 > 0$ and $t < t_s$) or Big Bang ($h_0 < 0$ and $t > t_s$) singularity at $t = t_s$. Another solution is $\epsilon < 1$, and $\beta = -\epsilon/(\epsilon - 2)$ ($-1 < \beta < 1$) case, which corresponds to (II.91) and to the Type II future singularity. In fact, we find that $\epsilon = -2n - 1$ and, therefore, $\beta + 1 = -2/(2n + 3)$. We should note that when $\epsilon = 2$, that is, $f(R) \sim R^2$, there is no singular solution. Therefore, if the above term $R^2 \tilde{f}(R)$, where $\lim_{R \rightarrow 0} \tilde{f}(R) = c_1$, $\lim_{R \rightarrow \infty} \tilde{f}(R) = c_2$, is added to $f(R)$ in (II.87), the added term dominates, and the modified $f(R)$ behaves as $f(R) \sim R^2$. The future singularity (II.91) disappears. We also note that if the R^n -term with $n = 3, 4, 5, \dots$ is added, the singularity becomes (in some sense) worse because this case corresponds to $\epsilon = n > 1$, that is the Big Rip case. Using the potential that appears when we transform $F(R)$ gravity to scalar-tensor theory [36], it has been found that the future singularity may not appear in the case where $0 < \epsilon < 2$.

Thus, to avoid the finite-time future singularity, one has to add the R^2 -term to the above viable unification models.

$$f(R) = \frac{\alpha R^{2n} - \beta R^n}{1 + \gamma R^n} + cR^2, \quad (\text{II.98})$$

$$f(R) = -\alpha_0 \left(\tanh \left(\frac{b_0(R - R_0)}{2} \right) + \tanh \left(\frac{b_0 R_0}{2} \right) \right) - \alpha_I \left(\tanh \left(\frac{b_I(R - R_I)}{2} \right) + \tanh \left(\frac{b_I R_I}{2} \right) \right) + cR^2, \quad (\text{II.99})$$

$$f(R) = -\frac{(R - R_0)^{2k+1} + R_0^{2k+1}}{f_0 + f_1 \left\{ (R - R_0)^{2k+1} + R_0^{2k+1} \right\}} + cR^2. \quad (\text{II.100})$$

The addition of this term was proposed first in ref. [37], where it was shown that, in this case, the Big Rip singularity disappears. Moreover, such a term, which effectively eliminates future singularity also supports the early-time inflation. In other words, adding such a R^2 term to the gravitational dark energy model (for instance, to viable dark energy models [15, 38]) may lead to the emergence of an inflationary phase in the model as was first observed in ref. [2]. In the case when a model already contains the inflationary era, its dynamics will be changed by the R^2 -term. The

investigation showing that the R^2 -term cures all types of future singularities was carried out in refs. [36, 39–41]. In fact, it has been recently realized that some phenomenological problems [42–46] of $F(R)$ dark energy (like achieving a consistent description of neutron stars) may be resolved in the presence of the R^2 term. Moreover, the traditional phantom/quintessence (fluid/scalar) dark energy models often bring the universe to a finite-time future singularity. It was demonstrated in ref. [47] that the natural method for eliminating the singularity in these models is again the addition of the R^2 term. In other words, resolving future singularities in specific dark energy models requires (at least partly) modified gravity. We discuss these questions in more detail in the fourth chapter. Note also that in the above models, there is no matter instability (Eq. (II.51)).

It should be noted that the exponential-type model may represent a very interesting proposal for the resolution of the dark energy problem. Some versions of such a theory are

$$F(R) = R + \alpha (e^{-bR} - 1) \quad \text{or} \quad F(R) = R - \alpha \left(\frac{e^{bR} - 1}{e^{bR} + e^{bR_0}} \right), \quad (\text{II.101})$$

with constants α , b , and R_0 . This model, which was introduced in ref. [32] provides the accelerating cosmological solutions without a future singularity. As a rule, the accelerating late-time cosmology leads asymptotically to a de Sitter universe. It was shown [48] that such a model describes a realistic dark energy epoch that is compatible with local and observational tests. Moreover, the number of free parameters is not big. Of course, one can consider different modifications of the above model.

A more complicated but viable $F(R)$ model

The above analysis shows that, to obtain a realistic and viable model, $F(R)$ gravity should satisfy the following conditions:

1. When $R \rightarrow 0$, the Einstein gravity is recovered, that is,

$$F(R) \rightarrow R \quad \text{that is,} \quad \frac{F(R)}{R^2} \rightarrow \frac{1}{R}. \quad (\text{II.102})$$

This also means that there is a flat space solution as in (II.69).

2. As will be discussed after Eq. (III.154), there appears a stable de Sitter solution, which corresponds to the late-time acceleration and, therefore, the curvature is small $R \sim R_L \sim (10^{-33} \text{ eV})^2$. This requires, when $R \sim R_L$,

$$\frac{F(R)}{R^2} = f_{0L} - f_{1L} (R - R_L)^{2n+2} + o\left((R - R_L)^{2n+2}\right). \quad (\text{II.103})$$

Here, f_{0L} and f_{1L} are positive constants and n is a positive integer. Of course, in some cases this condition may not be strictly necessary. One can use it, for instance, to avoid a future singularity.

3. As will also be discussed after Eq. (III.154), there appears a quasi-stable de Sitter solution that corresponds to the inflation of the early universe and, therefore, the curvature is large $R \sim R_I \sim (10^{16 \sim 19} \text{ GeV})^2$. The de Sitter space should not be exactly stable so that the curvature decreases very slowly. It requires

$$\frac{F(R)}{R^2} = f_{0I} - f_{1I} (R - R_I)^{2m+1} + o\left((R - R_I)^{2m+1}\right). \quad (\text{II.104})$$

Here, f_{0I} and f_{1I} are positive constants and m is a positive integer.

4. Following the discussion after (II.33), when $R \rightarrow \infty$, to avoid the curvature singularity, it is proposed that

$$F(R) \rightarrow f_{\infty} R^2 \quad \text{that is} \quad \frac{F(R)}{R^2} \rightarrow f_{\infty}. \quad (\text{II.105})$$

Here, f_{∞} is a positive and sufficiently small constant. Instead of (II.105), we may take

$$F(R) \rightarrow f_{\infty} R^{2-\epsilon} \quad \text{that is} \quad \frac{F(R)}{R^2} \rightarrow \frac{f_{\infty}}{R^{\epsilon}}. \quad (\text{II.106})$$

Here, f_{∞} is a positive constant and $0 < \epsilon < 1$. The above condition (II.105) or (II.106) prevents both the future singularity and the singularity due to large density of matter.




FIG. 1: The qualitative behavior of $\frac{F(R)}{R^2}$ versus R in a viable model.

5. As in (II.35), to avoid the anti-gravity, we require

$$F'(R) > 0, \quad (\text{II.107})$$

which is rewritten as

$$\frac{d}{dR} \left(\ln \left(\frac{F(R)}{R^2} \right) \right) > -\frac{2}{R}. \quad (\text{II.108})$$

6. Combining conditions (II.102) and (II.107), one finds

$$F(R) > 0. \quad (\text{II.109})$$

7. To avoid the matter instability in (II.51) [19], we require

$$U(R_b) \equiv \frac{R_b}{3} - \frac{F^{(1)}(R_b)F^{(3)}(R_b)R_b}{3F^{(2)}(R_b)^2} - \frac{F^{(1)}(R_b)}{3F^{(2)}(R_b)} + \frac{2F(R_b)F^{(3)}(R_b)}{3F^{(2)}(R_b)^2} - \frac{F^{(3)}(R_b)R_b}{3F^{(2)}(R_b)^2} < 0. \quad (\text{II.110})$$

The conditions 1 and 2 show that an extra, unstable de Sitter solution must appear at $R = R_e$ ($0 < R_e < R_L$) (see Fig.1). The universe evolution will stop at $R = R_L$ because the de Sitter solution $R = R_L$ is stable; the curvature never becomes smaller than R_L and, therefore, the extra de Sitter solution is not realized. The behavior of $\frac{F(R)}{R^2}$, which satisfies the conditions (II.102), (II.103), (II.104), (II.105), and (II.109) is given in Fig.1.

An example of such $F(R)$ gravity is given in [49]

$$\begin{aligned} \frac{F(R)}{R^2} &= \left\{ (X_m(R_I; R) - X_m(R_I; R_1)) (X_m(R_I; R) - X_m(R_I; R_L))^{2n+2} \right. \\ &\quad \left. + X_m(R_I; R_1) X_m(R_I; R_L)^{2n+2} + f_\infty^{2n+3} \right\}^{\frac{1}{2n+3}}, \\ X_m(R_I; R) &\equiv \frac{(2m+1) R_I^{2m}}{(R - R_I)^{2m+1} + R_I^{2m+1}}. \end{aligned} \quad (\text{II.111})$$

Here, n and m are integers greater or equal to unity, and $n, m \geq 1$ and R_1 is a parameter related with R_e by

$$X(R_I; R_e) = \frac{(2n+2) X(R_I; R_1) X(R_I; R_1) + X(R_I; R_L)}{2n+3}. \quad (\text{II.112})$$

It is assumed that

$$0 < R_1 < R_L \ll R_I. \quad (\text{II.113})$$

$X(R_I; R)$ is a monotonically decreasing function of R and, in the limit $R \rightarrow 0$, $X(R_I; R)$ behaves as

$$X(R_I; R) \rightarrow \frac{1}{R}, \quad (\text{II.114})$$

which shows that in the limit $R \rightarrow 0$, $F(R)$ (II.111) reproduces Eq. (II.102) and, therefore, the condition 1 is satisfied. On the other hand, when $R \rightarrow \infty$,

$$X(R_I; R) \rightarrow \frac{(2m+1) R_I^{2m}}{R^{2m+1}} \rightarrow 0, \quad (\text{II.115})$$

and, therefore, $F(R)$ behaves as (II.105) and the condition 4 is satisfied.

When $R \sim R_L$, we find the behavior of (II.103), where

$$\begin{aligned} f_{0L} &= \left\{ X_m(R_I; R_1) X_m(R_I; R_L)^{2n+2} + f_\infty^{2n+3} \right\}^{\frac{1}{2n+3}}, \\ f_{1L} &= \frac{1}{2m+3} \left\{ X_m(R_I; R_1) X_m(R_I; R_L)^{2n+2} + f_\infty^{2n+3} \right\}^{-\frac{2(n+1)}{2n+3}} (X_m(R_I; R_1) - X_m(R_I; R_L)) \\ &\quad \times \frac{(2m+1)^{4(m+1)} \{R_I(R_L - R_I)\}^{4m(m+1)}}{\left\{ (R_L - R_I)^{2m+1} + R_I^{2m+1} \right\}^{4(m+1)}}. \end{aligned} \quad (\text{II.116})$$

Then, the condition 2 is satisfied.

On the other hand, when $R \sim R_I$, we also find the behavior of (II.104), where

$$\begin{aligned} f_{0I} &= \left\{ (X_m(R_I; R_I) - X_m(R_I; R_1)) (X_m(R_I; R_I) - X_m(R_I; R_L))^{2n+2} \right. \\ &\quad \left. + X_m(R_I; R_1) X_m(R_I; R_L)^{2n+2} + f_\infty^{2n+3} \right\}^{\frac{1}{2n+3}} \\ f_{1I} &= \frac{2m+1}{R_I^{2m+2}} \left\{ (X_m(R_I; R_I) - X_m(R_I; R_1)) (X_m(R_I; R_I) - X_m(R_I; R_L))^{2n+2} \right. \\ &\quad \left. + X_m(R_I; R_1) X_m(R_I; R_L)^{2n+2} + f_\infty^{2n+3} \right\}^{\frac{2(n+1)}{2n+3}} \left\{ (X_m(R_I; R_I) - X_m(R_I; R_L))^{2n+2} \right. \\ &\quad \left. + (2n+2) (X_m(R_I; R_I) - X_m(R_I; R_1)) (X_m(R_I; R_I) - X_m(R_I; R_L))^{2n+1} \right\}. \end{aligned} \quad (\text{II.117})$$

Thus, the condition 3 is satisfied.

Let us now investigate the mass of the scalar field σ to determine whether or not the Chameleon mechanism [50, 51] works or not. For this purpose, one investigates the region $R_1 < R_L \ll R \ll R_I$. In this region, $X_m(R_I; R)$ can be approximated as

$$X_m(R_I; R) \sim \frac{1}{R}, \quad X_m(R_I; R_1) \sim \frac{1}{R_1}, \quad X_m(R_I; R_L) \sim \frac{1}{R_L}. \quad (\text{II.118})$$

Then, $F(R)$ can be approximated as

$$\frac{F(R)}{R^2} \sim f_\infty + \frac{f_n}{R}, \quad f_n \equiv \frac{R_1 + (2n+2) R_L f_\infty^{-2n-2}}{(2n+3) R_1 R_L^{2n+2}} \sim \frac{f_\infty^{-2n-2}}{R_L^{2n+2}}. \quad (\text{II.119})$$

In the last equation, it was assumed that $R_1 \sim R_L$. If $f_\infty R \ll f_n$ or

$$\frac{1}{f_\infty^{2n+3}} \gg R R_L^{2n+2}, \quad (\text{II.120})$$

one gets

$$m_\sigma^2 \sim \frac{3}{4f_\infty}. \quad (\text{II.121})$$

If we consider the region inside the earth, because $1\text{ g} \sim 6 \times 10^{32}\text{ eV}$ and $1\text{ cm} \sim (2 \times 10^{-5}\text{ eV})^{-1}$, the density is about $\rho_{\text{matter}} \sim 1\text{ g/cm}^3 \sim 5 \times 10^{18}\text{ eV}^4$. This shows that the magnitude of the curvature is $R \sim \kappa^2 \rho_{\text{matter}} \sim (10^{-19}\text{ eV})^2$. In the air on the earth, one finds $\rho_{\text{matter}} \sim 10^{-6}\text{ g/cm}^3 \sim 10^{12}\text{ eV}^4$, which gives $R_0 \sim \kappa^2 \rho_{\text{matter}} \sim (10^{-25}\text{ eV})^2$. In the solar system, there could be interstellar gas. Typically, in the interstellar gas, there is one proton (or hydrogen atom) per 1 cm^3 , which shows $\rho_{\text{matter}} \sim 10^{-5}\text{ eV}^4$, $R_0 \sim 10^{-61}\text{ eV}^2$. Then, the condition (II.120) can be easily satisfied; for example, we can choose $\frac{1}{f_\infty} \sim \text{MeV}^2$. Then, the Compton length of the scalar field becomes very small, and the correction to Newton's law is negligible.

One can now check the matter (in)stability issue [19]. In the earth or sun, the condition $R_1 < R_L \ll R \ll R_I$ is satisfied and, therefore, $F(R)$ behaves as (II.119). Then, $U(R_b)$ in (II.110) is given by

$$U(R_b) \sim -\frac{f_n}{6f_\infty} < 0. \quad (\text{II.122})$$

Because $U(R_b)$ is negative, the matter instability does not occur in the model under consideration.

The condition 5 remain to be checked. Because $F(R)$ (II.111) satisfies Eqs. (II.102) and (II.105), it is clear that Eq. (II.107) or (II.108) and, therefore, the condition 5 are satisfied when $R \rightarrow 0$ or $R \rightarrow \infty$. Because $\frac{F(R)}{R^2}$ is a monotonically increasing function of R in the region $R_e < R < R_L$, Eq. (II.107) or (II.108) is trivially satisfied in the region. In the region $R_1 < R_L \ll R \ll R_I$, because $\frac{F(R)}{R^2}$ behaves as (II.119), Eq. (II.107) or (II.108) is again satisfied. Then, the condition 5 seems to be satisfied throughout the whole region.

After the inflation at $R = R_I$, the radiation and the matter are generated. If the matter and radiation energy densities dominate compared with the contribution from $f(R) = F(R) - R$, that is, only the first term in (II.24) dominates, a radiation/matter dominated-universe will be realized. The radiation and matter densities decrease rapidly compared with the contribution from the $f(R)$ term at late-time universe. When curvature arrives at $R \sim R_L$, the late-time acceleration occurs.

The condition (II.106) indicates that the R^2 term appears in the very high curvature region. The R^2 term will generate an inflation besides the inflation at $R = R_I$ if the universe started with a very high curvature $R \gg R_I$. Because the inflation due to the R^2 -term is unstable, the inflation at the very early stage would stop; however when the curvature decreases and reaches $R \sim R_I$, the inflationary phase will occur again.

Thus, we demonstrated that a number of viable $F(R)$ gravity models may explain the early-time inflation in addition to the dark energy epoch in a unified way. It will be shown in the third chapter that when such a unification is problematic, one can use cosmological (partial/complete) reconstruction to ensure it.

B. $f(\mathcal{G})$ gravity

1. General properties

Let us consider another class of modified gravity where the arbitrary function added to the action of the general relativity depends on topological Gauss-Bonnet invariant. It has been shown recently that such a class of modified gravity is closely related with (super)string theory. The starting action is given by [52–54]:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} R + f(\mathcal{G}) + \mathcal{L}_{\text{matter}} \right). \quad (\text{II.123})$$

Here, $\mathcal{L}_{\text{matter}}$ is the Lagrangian density of matter and $\mathcal{G}$ is the Gauss-Bonnet invariant:

$$\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\xi\sigma}R^{\mu\nu\xi\sigma}. \quad (\text{II.124})$$

By the variation of the metric $g_{\mu\nu}$, we obtain

$$\begin{aligned} 0 = & \frac{1}{2\kappa^2} \left(-R^{\mu\nu} + \frac{1}{2}g^{\mu\nu}R \right) + T_{\text{matter}}^{\mu\nu} + \frac{1}{2}g^{\mu\nu}f(\mathcal{G}) - 2f'(\mathcal{G})R R^{\mu\nu} \\ & + 4f'(\mathcal{G})R^\mu_{\rho}R^{\nu\rho} - 2f'(\mathcal{G})R^{\mu\rho\sigma\tau}R^\nu_{\rho\sigma\tau} - 4f'(\mathcal{G})R^{\mu\rho\sigma\nu}R_{\rho\sigma} + 2(\nabla^\mu\nabla^\nu f'(\mathcal{G}))R \\ & - 2g^{\mu\nu}(\nabla^2 f'(\mathcal{G}))R - 4(\nabla_\rho\nabla^\mu f'(\mathcal{G}))R^{\nu\rho} - 4(\nabla_\rho\nabla^\nu f'(\mathcal{G}))R^{\mu\rho} \\ & + 4(\nabla^2 f'(\mathcal{G}))R^{\mu\nu} + 4g^{\mu\nu}(\nabla_\rho\nabla_\sigma f'(\mathcal{G}))R^{\rho\sigma} - 4(\nabla_\rho\nabla_\sigma f'(\mathcal{G}))R^{\mu\rho\nu\sigma}. \end{aligned} \quad (\text{II.125})$$

It is interesting to note that some memory of the fact that the Gauss-Bonnet term is topological invariant is contained in the above equation of motion. Indeed, it will be seen that such an equation of motion does not contain the higher-derivative terms as, for instance, the theory in the previous section.

Choosing the spatially flat FRW universe metric (II.6), the equation corresponding to the first FRW equation has the following form:

$$0 = -\frac{3}{\kappa^2}H^2 - f(\mathcal{G}) + \mathcal{G}f'(\mathcal{G}) - 24\dot{\mathcal{G}}f''(\mathcal{G})H^3 + \rho_{\text{matter}}. \quad (\text{II.126})$$

In the FRW universe (II.6), $\mathcal{G}$ looks like:

$$\mathcal{G} = 24 \left(H^2 \dot{H} + H^4 \right). \quad (\text{II.127})$$

Then, from Eq. (II.126), the FRW like equations, as in the Einstein gravity case (II.3), we find

$$\rho_{\text{eff}}^{\mathcal{G}} = \frac{3}{\kappa^2}H^2, \quad p_{\text{eff}}^{\mathcal{G}} = -\frac{1}{\kappa^2} \left(3H^2 + 2\dot{H} \right). \quad (\text{II.128})$$

Here,

$$\begin{aligned} \rho_{\text{eff}}^{\mathcal{G}} &\equiv -f(\mathcal{G}) + \mathcal{G}f'(\mathcal{G}) - 24\dot{\mathcal{G}}f''(\mathcal{G})H^3 + \rho_{\text{matter}}, \\ p_{\text{eff}}^{\mathcal{G}} &\equiv f(\mathcal{G}) - \mathcal{G}f'(\mathcal{G}) + \frac{2\mathcal{G}\dot{\mathcal{G}}}{3H}f''(\mathcal{G}) + 8H^2\ddot{\mathcal{G}}f''(\mathcal{G}) + 8H^2\dot{\mathcal{G}}^2f'''(\mathcal{G}) + p_{\text{matter}}. \end{aligned} \quad (\text{II.129})$$

When $\rho_{\text{matter}} = 0$, Eq. (II.126) has a de Sitter universe solution where H , and therefore $\mathcal{G}$, are constant. For $H = H_0$, with a constant H_0 , Eq. (II.126) turns into

$$0 = -\frac{3}{\kappa^2}H_0^2 + 24H_0^4f'(24H_0^4) - f(24H_0^4). \quad (\text{II.130})$$

As an example, we consider the model

$$f(\mathcal{G}) = f_0 |\mathcal{G}|^\beta, \quad (\text{II.131})$$

with constants f_0 and β . Then, the solution of Eq. (II.130) is given by

$$H_0^4 = \frac{1}{24(8(n-1)\kappa^2f_0)^{\frac{1}{\beta-1}}}. \quad (\text{II.132})$$

For a large number of choices of the function $f(\mathcal{G})$, Eq. (II.130) has a non-trivial ($H_0 \neq 0$) real solution for H_0 (de Sitter universe). The late-time cosmology for above theory without matter has been discussed for a number of examples in refs. [52–54]. We will refer to these studies in the following.

The de Sitter universe can be expressed as an accelerating, expanding universe,

$$ds^2 = -dt^2 + e^{2H_0 t} \sum_{i=1,2,3} (dx^i)^2. \quad (\text{II.133})$$

Changing the coordinates, the de Sitter universe can be described by the static metric, which can be obtained by putting $M = 0$ in the metric (II.10) (and choosing the minus sign in $\mp$):

$$ds^2 = -\left(1 - \frac{r^2}{L^2}\right) dt^2 + \left(1 - \frac{r^2}{L^2}\right)^{-1} dr^2 + r^2 d\Omega^2. \quad (\text{II.134})$$

The length parameter L can be identified with $1/H_0$. Note that there is a cosmological horizon at $r = L$. When $r \sim L$, by the Wick-rotating time coordinate $t = i\tau$, the metric has the following form:

$$ds^2 \sim 2\left(1 - \frac{r}{L}\right) d\tau^2 + \frac{dr^2}{2\left(1 - \frac{r}{L}\right)} + l^2 d\Omega^2. \quad (\text{II.135})$$

Changing the radial coordinate r to ρ as

$$r = L \left(1 - \frac{\rho^2}{2L^2}\right), \quad (\text{II.136})$$

the metric (II.135) can be rewritten as

$$ds^2 \sim \frac{\rho^2}{L^2} d\tau^2 + d\rho^2 + l^2 d\Omega^2. \quad (\text{II.137})$$

To avoid the conical singularity at $\rho = 0$, the periodicity condition for the coordinate τ should be imposed as $\tau \sim \tau + 2\pi L$. Thus, the Hawking temperature T_H is defined as

$$T_H = \frac{1}{2\pi L}. \quad (\text{II.138})$$

Let us now calculate the entropy like in Eq. (II.18). The partition function Z (II.16) can be evaluated from the action (II.123) by neglecting the matter Lagrangian $\mathcal{L}_{\text{matter}}$ and substituting the metric of the de Sitter universe (II.134):

$$\begin{aligned} -\ln Z \sim S &= 4\pi \int_0^{2\pi L} dt \int_0^L dr r^2 \left(\frac{6}{\kappa^2 L^2} + f\left(\frac{24}{L^4}\right) \right) \\ &= \frac{8\pi^2 L^4}{3} \left(\frac{6}{\kappa^2 L^2} + f\left(\frac{24}{L^4}\right) \right) \\ &= \frac{1}{6\pi^2 T_H^4} \left(\frac{24\pi^2 T_H^2}{\kappa^2} + f(384\pi^4 T_H^4) \right). \end{aligned} \quad (\text{II.139})$$

Note that $R = \frac{12}{L^2}$ and $\mathcal{G} = \frac{24}{L^4}$. Using (II.16) and (III.148), the free energy $\mathcal{F}$ and the entropy $\mathcal{S}$ are

$$\mathcal{F} = \frac{1}{6\pi^2 T_H^3} \left(\frac{24\pi^2 T_H^2}{\kappa^2} + f(384\pi^4 T_H^4) \right), \quad (\text{II.140})$$

$$\begin{aligned} \mathcal{S} &= -\frac{4}{\kappa^2 T_H^2} - \frac{f(384\pi^4 T_H^4)}{2\pi^2 T_H^4} 256\pi^2 f'(384\pi^4 T_H^4) \\ &= -\frac{16\pi^2 L^2}{\kappa^2} - 8\pi^2 L^4 f\left(\frac{24}{L^4}\right) + 256\pi^2 f'\left(\frac{24}{L^4}\right). \end{aligned} \quad (\text{II.141})$$

Because $H_0 = \frac{1}{L}$, Eq. (II.130) can be rewritten as

$$0 = -\frac{3}{\kappa^2 L^2} + \frac{24}{L^4} f'\left(\frac{24}{L^4}\right) - f\left(\frac{24}{L^4}\right). \quad (\text{II.142})$$

By using (II.142), the entropy (II.141) has the following expression:

$$\mathcal{S} = \frac{A_H}{4G} + 64\pi^2 f'\left(\frac{24}{L^4}\right). \quad (\text{II.143})$$

Here, A_H is the area of the cosmological horizon: $A_H = 4\pi L^2$ and $G = \frac{\kappa^2}{8\pi}$. When $f(\mathcal{G})$ is a constant, which corresponds to the cosmological constant, Eq. (II.143) reproduces the standard result.

One now considers the case $\rho_{\text{matter}} \neq 0$. Assuming that the EoS parameter $w \equiv p_{\text{matter}}/\rho_{\text{matter}}$ is a constant, using the conservation of energy:

$$\dot{\rho}_{\text{matter}} + 3H(\rho_{\text{matter}} + p_{\text{matter}}) = 0, \quad (\text{II.144})$$

we find $\rho = \rho_0 a^{-3(1+w)}$. When the function $f(\mathcal{G})$ is chosen as in (II.131), again, if $\beta < 1/2$, the $f(\mathcal{G})$ term becomes dominant compared with the Einstein term when the curvature is small. If we neglect the contribution from the Einstein term in (II.126), the following solution may be found

$$\begin{aligned} a &= a_0 t^{h_0} \quad \text{when } h_0 > 0, \quad a = a_0 (t_s - t)^{h_0} \quad \text{when } h_0 < 0, \\ h_0 &= \frac{4\beta}{3(1+w)}, \quad a_0 = \left[-\frac{f_0(\beta-1)}{(h_0-1)\rho_0} \{24|h_0^3(-1+h_0)|\}^\beta (h_0-1+4\beta) \right]^{-\frac{1}{3(1+w)}}. \end{aligned} \quad (\text{II.145})$$

Thus, the effective EoS parameter w_{eff} (II.4) is less than -1 if $\beta < 0$, and for $w > -1$, it is

$$w_{\text{eff}} = -1 + \frac{2}{3h_0} = -1 + \frac{1+w}{2\beta}, \quad (\text{II.146})$$

which is again less than -1 for $\beta < 0$. Thus, if $\beta < 0$, we obtain an effective phantom with negative h_0 even when $w > -1$. In the phantom phase, the Big Rip might seem to occur at $t = t_s$. Near this Big Rip, however, the curvature becomes dominant and then the Einstein term dominates, so the $f(\mathcal{G})$ term can be neglected. Therefore, the universe behaves as $a = a_0 t^{2/3(w+1)}$ and as a consequence, the Big Rip does not occur. The phantom era is thus transient.

The case when $0 < \beta < 1/2$ may also be considered. As β is positive, the universe does not reach the phantom phase. When the curvature is large, the $f(\mathcal{G})$ term in the action (II.123) can be neglected and one can work with the Einstein gravity. Then, if w is positive, the matter energy density ρ_{matter} should behave as $\rho_{\text{matter}} \sim t^{-2}$, but $f(\mathcal{G})$ goes as $f(\mathcal{G}) \sim t^{-4\beta}$. For late times (large t), the $f(\mathcal{G})$ term may become dominant as compared with the Lagrangian density of matter. Neglecting the contribution from matter, Eq. (II.126) has a de Sitter universe solution where H , and, therefore, $\mathcal{G}$, are constant. If $H = H_0$ with a constant H_0 , Eq. (II.126) looks as (II.130). As a consequence, even if we start from the deceleration phase with $w > -1/3$, an asymptotically de Sitter universe emerges. Correspondingly, a transition also occurs here from deceleration to acceleration of the universe.

Now, we consider the case in which the contributions from the Einstein and matter terms can be neglected. Eq. (II.126) reduces to

$$0 = \mathcal{G}f'(\mathcal{G}) - f(\mathcal{G}) - 24\dot{\mathcal{G}}f''(\mathcal{G})H^3. \quad (\text{II.147})$$

If $f(\mathcal{G})$ behaves as (II.131), assuming

$$a = \begin{cases} a_0 t^{h_0} & \text{when } h_0 > 0 \text{ (quintessence)} \\ a_0 (t_s - t)^{h_0} & \text{when } h_0 < 0 \text{ (phantom)} \end{cases}, \quad (\text{II.148})$$

one obtains

$$0 = (\beta - 1) h_0^6 (h_0 - 1) (h_0 - 1 + 4\beta). \quad (\text{II.149})$$

As $h_0 = 1$ implies $\mathcal{G} = 0$, one may choose

$$h_0 = 1 - 4\beta, \quad (\text{II.150})$$

and Eq. (II.4) gives

$$w_{\text{eff}} = -1 + \frac{2}{3(1 - 4\beta)}. \quad (\text{II.151})$$

Therefore, if $\beta > 0$, the universe is accelerating ($w_{\text{eff}} < -1/3$), and if $\beta > 1/4$, the universe is in a phantom phase ($w_{\text{eff}} < -1$). Thus, we are led to consider the following model:

$$f(\mathcal{G}) = f_i |\mathcal{G}|^{\beta_i} + f_l |\mathcal{G}|^{\beta_l}, \quad (\text{II.152})$$

where it is assumed that

$$\beta_i > \frac{1}{2}, \quad \frac{1}{2} > \beta_l > \frac{1}{4}. \quad (\text{II.153})$$

Then, when the curvature is large, as in the primordial universe, the first term dominates, compared with the second term and the Einstein term, and it gives

$$-1 > w_{\text{eff}} = -1 + \frac{2}{3(1 - 4\beta_i)} > -\frac{5}{3}. \quad (\text{II.154})$$

On the other hand, when the curvature is small, as is the case in the present universe, the second term in (II.152) dominates compared with the first term and the Einstein term and yields

$$w_{\text{eff}} = -1 + \frac{2}{3(1 - 4\beta_l)} < -\frac{5}{3}. \quad (\text{II.155})$$

Therefore, theory (II.152) can produce a model that is able to describe inflation and the late-time acceleration of the universe in a unified manner.

Instead of (II.153), one may also choose β_l as

$$\frac{1}{4} > \beta_l > 0, \quad (\text{II.156})$$

which gives

$$-\frac{1}{3} > w_{\text{eff}} > -1. \quad (\text{II.157})$$

Then, we obtain an effective quintessence epoch. Moreover, by properly adjusting the couplings f_i and f_l in (II.152), one can obtain a period where the Einstein term dominates and the universe is in a deceleration phase. After that, a transition occurs from deceleration to acceleration when the Gauss-Bonnet term becomes the dominant one. More choices of $f(\mathcal{G})$ may be studied for the purpose of the construction of the current accelerating universe. Nevertheless, many non-linear choices for this function may be approximated by the above model. For instance, one can mention some realistic examples of $f(\mathcal{G})$ gravity:

$$f_1(\mathcal{G}) = \frac{a_1 \mathcal{G}^n + b_1}{a_2 \mathcal{G}^n + b_2}, \quad f_2(\mathcal{G}) = \frac{a_1 \mathcal{G}^{n+N} + b_1}{a_2 \mathcal{G}^n + b_2}. \quad (\text{II.158})$$

Let us address the issue of the correction to Newton's law. Let $g_{(0)}$ be a solution of (II.125) and represent the perturbation of the metric as $g_{\mu\nu} = g_{(0)\mu\nu} + h_{\mu\nu}$. First, we consider the perturbation around the de Sitter background. The de Sitter space metric is taken as $g_{(0)\mu\nu}$, which gives the following Riemann tensor:

$$R_{(0)\mu\nu\rho\sigma} = H_0^2 (g_{(0)\mu\rho}g_{(0)\nu\sigma} - g_{(0)\mu\sigma}g_{(0)\nu\rho}). \quad (\text{II.159})$$

The flat background corresponds to the limit of $H_0 \rightarrow 0$. For simplicity, the following gauge condition is chosen: $g_{(0)}^{\mu\nu}h_{\mu\nu} = \nabla_{(0)}^\mu h_{\mu\nu} = 0$. Then Eq. (II.125) gives

$$0 = \frac{1}{4\kappa^2} (\nabla^2 h_{\mu\nu} - 2H_0^2 h_{\mu\nu}) + T_{\text{matter}\mu\nu}. \quad (\text{II.160})$$

The Gauss-Bonnet term contribution does not appear except in the length parameter $1/H_0$ of the de Sitter space, which is determined by taking into account the Gauss-Bonnet term. This may occur due to the special structure of the Gauss-Bonnet invariant. Eq. (II.160) shows that there is no correction to Newton's law in de Sitter space and even in the flat background corresponding to $H_0 \rightarrow 0$, regardless of the form of f (at least, with the above gauge condition). In contrast, the study of the Newtonian limit in $1/R$ gravity indicates that significant corrections to Newton's law may appear as is shown in refs. [23]. For most $1/R$ models, the corrections to Newton's law do not comply with solar system tests.

By introducing two auxiliary fields, A and B , one can rewrite the action (II.123) as

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} R + B(\mathcal{G} - A) + f(A) + \mathcal{L}_{\text{matter}} \right). \quad (\text{II.161})$$

By the variation of the action (II.161) with respect to B , it follows that $A = \mathcal{G}$. Using this in (II.161), the action (II.123) is recovered. Alternatively, varying the action (II.161) with respect to A in (II.161), one gets $B = f'(A)$, and, thus,

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} R + f'(A)\mathcal{G} - A f'(A) + f(A) \right). \quad (\text{II.162})$$

By varying the action (II.162) with respect to A , the relation $A = \mathcal{G}$ is obtained again. The scalar is not dynamic as it has no kinetic term. One may add, however, a kinetic term to the action by hand

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} R - \frac{\epsilon}{2} \partial_\mu A \partial^\mu A + f'(A)\mathcal{G} - A f'(A) + f(A) \right). \quad (\text{II.163})$$

Then, one obtains a dynamical scalar theory coupled with the Gauss-Bonnet invariant and with a potential. It is known that, in general, a theory of this kind has no ghosts and is stable. Actually, it is related to string-inspired dilaton gravity, which was proposed as an alternative for dark energy [55–57]. Then, when the limit $\epsilon \rightarrow 0$ can be obtained smoothly, the corresponding $f(\mathcal{G})$ theory has no ghost and could actually be stable. It may be of interest to study the cosmology of such a theory in the limit $\epsilon \rightarrow 0$.

Thus, we investigated the modified Gauss-Bonnet gravity and demonstrated that it may naturally lead to the unified cosmic history. Further study of cosmological properties of such a theory as well as discussion of particular models of the modified Gauss-Bonnet gravity may be found in refs. [58].

2. $F(\mathcal{G}, R)$ gravity

More general models include the dependence from curvature as well as from the Gauss-Bonnet term [54]:

$$S = \int d^4x \sqrt{-g} (F(\mathcal{G}, R) + \mathcal{L}_{\text{matter}}) . \quad (\text{II.164})$$

An example is given by

$$F(\mathcal{G}, R) = R \tilde{f}\left(\frac{\mathcal{G}}{R^2}\right) , \quad \tilde{f}\left(\frac{\mathcal{G}}{R^2}\right) = \frac{1}{2\kappa^2} + f_0 \left(\frac{\mathcal{G}}{R^2}\right) , \quad (\text{II.165})$$

which gives a solution describing the FRW universe:

$$H = \frac{h_0}{t} , \quad h_0 = \frac{\frac{3}{\kappa^2} - 2f_0 \pm \sqrt{8f_0(f_0 - \frac{3}{8\kappa^2})}}{\frac{6}{\kappa^2} + 2f_0} . \quad (\text{II.166})$$

Then, for example, if $\kappa^2 f_0 < -3$, there is a solution describing a phantom with $h_0 < -1 - \sqrt{2}$ and a solution describing the effective matter with $h_0 > -1 + \sqrt{2}$. Late-time cosmology in other versions of such theory can be constructed. The unification with inflation in such model may be achieved via cosmological reconstruction, which is developed in the third chapter.

C. String-inspired model and scalar-Einstein-Gauss-Bonnet gravity

In string theories, the compactification from higher dimensions to four dimensions induces many scalar fields, such as moduli or dilaton fields. These scalars couple with curvature invariants. Neglecting the moduli fields associated with the radii of the internal space, we may consider the following action of the low-energy effective string theories [56, 57]:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2} + \mathcal{L}_\phi + \mathcal{L}_c + \dots \right] , \quad (\text{II.167})$$

where ϕ is the dilaton field, which is related to the string coupling, $\mathcal{L}_\phi$ is the Lagrangian of ϕ , and $\mathcal{L}_c$ expresses the string curvature correction terms to the Einstein-Hilbert action,

$$\mathcal{L}_\phi = -\partial_\mu \phi \partial^\mu \phi - V(\phi) , \quad \mathcal{L}_c = c_1 \alpha' e^{2\frac{\phi}{\phi_0}} \mathcal{L}_c^{(1)} + c_2 \alpha'^2 e^{4\frac{\phi}{\phi_0}} \mathcal{L}_c^{(2)} + c_3 \alpha'^3 e^{6\frac{\phi}{\phi_0}} \mathcal{L}_c^{(3)} , \quad (\text{II.168})$$

where $1/\alpha'$ is the string tension, $\mathcal{L}_c^{(1)}$, $\mathcal{L}_c^{(2)}$, and $\mathcal{L}_c^{(3)}$ express the leading-order (Gauss-Bonnet term $\mathcal{G}$ in (II.124)), the second-order, and the third-order curvature corrections, respectively. The terms $\mathcal{L}_c^{(1)}$, $\mathcal{L}_c^{(2)}$ and $\mathcal{L}_c^{(3)}$ in the Lagrangian have the following form

$$\mathcal{L}_c^{(1)} = \Omega_2 , \quad \mathcal{L}_c^{(2)} = 2\Omega_3 + R_{\alpha\beta}^{\mu\nu} R_{\lambda\rho}^{\alpha\beta} R_{\mu\nu}^{\lambda\rho} , \quad \mathcal{L}_c^{(3)} = \mathcal{L}_{31} - \delta_H \mathcal{L}_{32} - \frac{\delta_B}{2} \mathcal{L}_{33} . \quad (\text{II.169})$$

Here, δ_B and δ_H take the value of 0 or 1 and

$$\begin{aligned} \Omega_2 &= \mathcal{G} , \\ \Omega_3 &\propto \epsilon^{\mu\nu\rho\sigma\tau\eta} \epsilon_{\mu'\nu'\rho'\sigma'\tau'\eta'} R_{\mu\nu}^{\mu'\nu'} R_{\rho\sigma}^{\rho'\sigma'} R_{\tau\eta}^{\tau'\eta'} , \\ \mathcal{L}_{31} &= \zeta(3) R_{\mu\nu\rho\sigma} R^{\alpha\nu\rho\beta} \left(R_{\delta\beta}^{\mu\gamma} R_{\alpha\gamma}^{\delta\sigma} - 2 R_{\delta\alpha}^{\mu\gamma} R_{\beta\gamma}^{\delta\sigma} \right) , \\ \mathcal{L}_{32} &= \frac{1}{8} (R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta})^2 + \frac{1}{4} R_{\mu\nu}^{\gamma\delta} R_{\gamma\delta}^{\rho\sigma} R_{\rho\sigma}^{\alpha\beta} R_{\alpha\beta}^{\mu\nu} - \frac{1}{2} R_{\mu\nu}^{\alpha\beta} R_{\alpha\beta}^{\rho\sigma} R_{\sigma\gamma\delta}^{\mu} R_{\rho}^{\nu\gamma\delta} - R_{\mu\nu}^{\alpha\beta} R_{\alpha\beta}^{\rho\nu} R_{\rho\sigma}^{\gamma\delta} R_{\gamma\delta}^{\mu\sigma} , \\ \mathcal{L}_{33} &= (R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta})^2 - 10 R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\sigma} R_{\sigma\gamma\delta\rho} R^{\beta\gamma\delta\rho} - R_{\mu\nu\alpha\beta} R^{\mu\nu\rho}_{\sigma} R^{\beta\sigma\gamma\delta} R_{\delta\gamma\rho}^{\alpha} . \end{aligned} \quad (\text{II.170})$$

The correction terms are different depending on the type of string theory; the dependence is encoded in the curvature invariants and in the coefficients (c_1, c_2, c_3) and δ_H , δ_B , as follows,

- For the Type II superstring theory: $(c_1, c_2, c_3) = (0, 0, 1/8)$ and $\delta_H = \delta_B = 0$.
- For the heterotic superstring theory: $(c_1, c_2, c_3) = (1/8, 0, 1/8)$ and $\delta_H = 1, \delta_B = 0$.
- For the bosonic superstring theory: $(c_1, c_2, c_3) = (1/4, 1/48, 1/8)$ and $\delta_H = 0, \delta_B = 1$.

Motivated by the string considerations, we consider the scalar-Einstein-Gauss-Bonnet gravity² based on [55, 60]. It was first proposed in ref. [55] to consider such a theory as a gravitational alternative for dark energy, so-called Gauss-Bonnet dark energy.

The starting action is:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \xi(\phi) \mathcal{G} \right]. \quad (\text{II.171})$$

Here, we do not restrict the forms of $V(\phi)$ and $\xi(\phi)$ which should be given by the non-perturbative string theory (II.168). Note also that the action (II.171) is given by adding the kinetic term for the scalar field ϕ to the action of the $F(\mathcal{G})$ gravity in the scalar-tensor form that appeared in the previous section.

Making the variation of the action (II.171) over the metric $g_{\mu\nu}$, the field equations follow as:

$$\begin{aligned} 0 = & \frac{1}{\kappa^2} \left(-R^{\mu\nu} + \frac{1}{2} g^{\mu\nu} R \right) + \frac{1}{2} \partial^\mu \phi \partial^\nu \phi - \frac{1}{4} g^{\mu\nu} \partial_\rho \phi \partial^\rho \phi + \frac{1}{2} g^{\mu\nu} (-V(\phi) + \xi(\phi) \mathcal{G}) \\ & - 2\xi(\phi) R R^{\mu\nu} - 4\xi(\phi) R^\mu{}_\rho R^{\nu\rho} - 2\xi(\phi) R^{\mu\rho\sigma\tau} R^\nu{}_{\rho\sigma\tau} + 4\xi(\phi) R^{\mu\rho\nu\sigma} R_{\rho\sigma} \\ & + 2(\nabla^\mu \nabla^\nu \xi(\phi)) R - 2g^{\mu\nu} (\nabla^2 \xi(\phi)) R - 4(\nabla_\rho \nabla^\mu \xi(\phi)) R^{\nu\rho} - 4(\nabla_\rho \nabla^\nu \xi(\phi)) R^{\mu\rho} \\ & + 4(\nabla^2 \xi(\phi)) R^{\mu\nu} + 4g^{\mu\nu} (\nabla_\rho \nabla_\sigma \xi(\phi)) R^{\rho\sigma} + 4(\nabla_\rho \nabla_\sigma \xi(\phi)) R^{\mu\rho\nu\sigma}. \end{aligned} \quad (\text{II.172})$$

In Eq. (II.172), the derivatives of curvature such as ∇R , do not appear. Therefore, the derivatives higher than two do not appear, which can be contrasted with a general $\alpha R^2 + \beta R_{\mu\nu} R^{\mu\nu} + \gamma R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}$ gravity, where fourth derivatives of $g_{\mu\nu}$ appear. For the classical theory, if we specify the values of $g_{\mu\nu}$ and $\dot{g}_{\mu\nu}$ on a spatial surface as an initial condition, the time development is uniquely determined. This situation is similar to the case in classical mechanics, in which one only needs to specify the values of position and velocity of particle as initial conditions. In general $\alpha R^2 + \beta R_{\mu\nu} R^{\mu\nu} + \gamma R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}$ gravity, we need to specify the values of $\ddot{g}_{\mu\nu}$ and $\ddot{\ddot{g}}_{\mu\nu}$ in addition to $g_{\mu\nu}$, $\dot{g}_{\mu\nu}$ so that a unique time development will follow. In Einstein gravity, only the specific value of $g_{\mu\nu}$, $\dot{g}_{\mu\nu}$, should be given as an initial condition. Thus, the scalar-Gauss-Bonnet gravity is a natural extension of the Einstein gravity.

In the FRW universe (II.6), Eq. (II.172) becomes the following:

$$0 = -\frac{3}{\kappa^2} H^2 + \frac{1}{2} \dot{\phi}^2 + V(\phi) + 24H^3 \frac{d\xi(\phi(t))}{dt}, \quad (\text{II.173})$$

$$0 = \frac{1}{\kappa^2} \left(2\dot{H} + 3H^2 \right) + \frac{1}{2} \dot{\phi}^2 - V(\phi) - 8H^2 \frac{d^2 \xi(\phi(t))}{dt^2} - 16H\dot{H} \frac{d\xi(\phi(t))}{dt} - 16H^3 \frac{d\xi(\phi(t))}{dt}. \quad (\text{II.174})$$

On the other hand, by the variation of the action (II.171) with respect to the scalar field, the scalar equation of motion follows as

$$0 = \ddot{\phi} + 3H\dot{\phi} + V'(\phi) + \xi'(\phi) \mathcal{G}. \quad (\text{II.175})$$

In particular when we consider the following string-inspired model [55],

$$V = V_0 e^{-\frac{2\phi}{\phi_0}}, \quad \xi(\phi) = \xi_0 e^{\frac{2\phi}{\phi_0}}, \quad (\text{II.176})$$

the de Sitter space solution follows:

$$H^2 = H_0^2 \equiv -\frac{e^{-\frac{2\varphi_0}{\phi_0}}}{8\xi_0\kappa^2}, \quad \phi = \varphi_0. \quad (\text{II.177})$$

² For pioneering work on the scalar-Einstein-Gauss-Bonnet gravity, see [59].

Here, φ_0 is an arbitrary constant. If φ_0 is chosen to be larger, the Hubble rate $H = H_0$ becomes smaller. Then, if $\xi_0 \sim \mathcal{O}(1)$, by choosing $\varphi_0/\phi_0 \sim 140$, the value of the Hubble rate $H = H_0$ is consistent with the observations. The model (II.176) also has another solution:

$$\begin{aligned} H &= \frac{h_0}{t}, \quad \phi = \phi_0 \ln \frac{t}{t_1} \quad \text{when } h_0 > 0, \\ H &= -\frac{h_0}{t_s - t}, \quad \phi = \phi_0 \ln \frac{t_s - t}{t_1} \quad \text{when } h_0 < 0. \end{aligned} \quad (\text{II.178})$$

Here, h_0 is obtained by solving the following algebraic equations:

$$0 = -\frac{3h_0^2}{\kappa^2} + \frac{\phi_0^2}{2} + V_0 t_1^2 - \frac{48\xi_0 h_0^3}{t_1^2}, \quad 0 = (1 - 3h_0)\phi_0^2 + 2V_0 t_1^2 + \frac{48\xi_0 h_0^3}{t_1^2}(h_0 - 1). \quad (\text{II.179})$$

Eqs. (II.179) can be rewritten as

$$V_0 t_1^2 = -\frac{1}{\kappa^2(1+h_0)} \left\{ 3h_0^2(1-h_0) + \frac{\phi_0^2 \kappa^2 (1-5h_0)}{2} \right\}, \quad (\text{II.180})$$

$$\frac{48\xi_0 h_0^2}{t_1^2} = -\frac{6}{\kappa^2(1+h_0)} \left(h_0 - \frac{\phi_0^2 \kappa^2}{2} \right). \quad (\text{II.181})$$

The arbitrary value of h_0 can be realized by properly choosing V_0 and ξ_0 . With the appropriate choice of V_0 and ξ_0 , we can obtain a negative h_0 and, therefore, the effective EoS parameter (II.4) is less than -1 , $w_{\text{eff}} < -1$, which corresponds to the effective phantom. In usual (canonical) scalar-tensor theory without the Gauss-Bonnet term, the phantom cannot be realized by the canonical scalar.

For example, if $h_0 = -80/3 < -1$ and, therefore, $w = -1.025$, which is consistent with the observed value, we find

$$V_0 t_1^2 = \frac{1}{\kappa^2} \left(\frac{531200}{231} + \frac{403}{154} \gamma \phi_0^2 \kappa^2 \right) > 0, \quad \frac{f_0}{t_1^2} = -\frac{1}{\kappa^2} \left(\frac{9}{49280} + \frac{27}{7884800} \gamma \phi_0^2 \kappa^2 \right). \quad (\text{II.182})$$

For other choices of scalar potentials one can realize other types of dark energy universes, for instance, the effective quintessence. Moreover, one can propose the potentials in such a way that the unification of the inflation with dark energy naturally occurs. Different cosmological aspects of scalar-Gauss-Bonnet dark energy have been studied in refs. [61]. An extension of these studies taking into account the Euler invariant and other higher-derivative stringy terms has also been done [62].

D. Non-local gravity

1. Non-local $F(R)$ gravity

In this section, we review non-local gravity [63–65] and present its local (scalar-tensor) formulation. The explicit example of such a theory, which naturally leads to the unification of inflation with late-time cosmic acceleration, is worked out.

The starting action of the simplest non-local gravity is given by

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} \{ R(1 + f(\square^{-1}R)) - 2\Lambda \} + \mathcal{L}_{\text{matter}}(Q; g) \right\}. \quad (\text{II.183})$$

Here, f is some function, $\square$ is the d'Alembertian for the scalar field, Λ is a cosmological constant and Q represents the matter fields. The above action can be rewritten by introducing two scalar fields, η and ξ , in the following form:

$$\begin{aligned} S &= \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} \{ R(1 + f(\eta)) + \xi(\square\eta - R) - 2\Lambda \} + \mathcal{L}_{\text{matter}} \right] \\ &= \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} \{ R(1 + f(\eta)) - \partial_\mu \xi \partial^\mu \eta - \xi R - 2\Lambda \} + \mathcal{L}_{\text{matter}} \right]. \end{aligned} \quad (\text{II.184})$$

By the variation of the action (II.184) over ξ , we obtain $\square\eta = R$ or $\eta = \square^{-1}R$. Substituting the above equation into (II.184), one re-obtains (II.183).

The variation of (II.184) with respect to the metric tensor $g_{\mu\nu}$ gives

$$\begin{aligned} 0 = & \frac{1}{2}g_{\mu\nu} \{R(1+f(\eta)-\xi) - \partial_\rho \xi \partial^\rho \eta - 2\Lambda\} - R_{\mu\nu}(1+f(\eta)-\xi) \\ & + \frac{1}{2}(\partial_\mu \xi \partial_\nu \eta + \partial_\mu \eta \partial_\nu \xi) - (g_{\mu\nu}\square - \nabla_\mu \nabla_\nu)(f(\eta)-\xi) + \kappa^2 T_{\mu\nu}. \end{aligned} \quad (\text{II.185})$$

On the other hand, the variation with respect to η gives

$$0 = \square \xi + f'(\eta)R. \quad (\text{II.186})$$

Now, we assume that the spatially flat FRW metric and the scalar fields η and ξ only depend on time. Then, Eq. (II.185) has the following form:

$$0 = -3H^2(1+f(\eta)-\xi) + \frac{1}{2}\dot{\xi}\dot{\eta} - 3H(f'(\eta)\dot{\eta} - \dot{\xi}) + \Lambda + \kappa^2 \rho_{\text{matter}}, \quad (\text{II.187})$$

$$0 = (2\dot{H} + 3H^2)(1+f(\eta)-\xi) + \frac{1}{2}\dot{\xi}\dot{\eta} + \left(\frac{d^2}{dt^2} + 2H\frac{d}{dt}\right)(f(\eta)-\xi) - \Lambda + \kappa^2 p. \quad (\text{II.188})$$

On the other hand, the scalar equations are:

$$0 = \ddot{\eta} + 3H\dot{\eta} + 6\dot{H} + 12H^2, \quad (\text{II.189})$$

$$0 = \ddot{\xi} + 3H\dot{\xi} - (6\dot{H} + 12H^2)f'(\eta). \quad (\text{II.190})$$

For the de Sitter solution $H = H_0$, Eq. (II.189) can be solved as

$$\eta = -4H_0 t - \eta_0 e^{-3H_0 t} + \eta_1, \quad (\text{II.191})$$

with constants of integration η_0 and η_1 . For simplicity, let $\eta_0 = \eta_1 = 0$ and $f(\eta)$ be given by

$$f(\eta) = f_0 e^{\frac{\eta}{\beta}} = f_0 e^{-\frac{4H_0 t}{\beta}}. \quad (\text{II.192})$$

Here, β is a constant. Then, Eq. (II.190) can be solved as follows:

$$\xi = -\frac{3f_0\beta}{3\beta-4}e^{-\frac{4H_0 t}{\beta}} + \frac{\xi_0}{3H_0}e^{-3H_0 t} - \xi_1. \quad (\text{II.193})$$

Here, ξ_0 and ξ_1 are constants. For the de Sitter space a behaves as $a = a_0 e^{H_0 t}$. For matter with a constant equation of state w , we find

$$\rho_{\text{matter}} = \rho_0 e^{-3(w+1)H_0 t}. \quad (\text{II.194})$$

Then, after setting $\xi_0 = 0$, by substituting (II.191), (II.193), and (II.194) into (II.187), one obtains

$$0 = -3H_0^2(1+\xi_1) + 6H_0^2 f_0 \left(\frac{2}{\beta} - 1\right) e^{-\frac{4H_0 t}{\beta}} + \Lambda + \kappa^2 \rho_0 e^{-3(w+1)H_0 t}. \quad (\text{II.195})$$

When $\rho_0 = 0$, with the choice

$$\beta = 2, \quad \xi_1 = -1 + \frac{\Lambda}{3H_0^2}, \quad (\text{II.196})$$

the de Sitter space can be a solution. Even if $\rho_{\text{matter}} \neq 0$, if we choose

$$\beta = \frac{4}{3(1+w)}, \quad f_0 = \frac{\kappa^2 \rho_0}{3H_0^2(1+3w)}, \quad \xi_1 = -1 + \frac{\Lambda}{3H_0^2}, \quad (\text{II.197})$$

there is a de Sitter solution.

Eq. (II.196) or (II.197) shows that

$$H_0^2 = \frac{3\Lambda}{1+\xi_1}. \quad (\text{II.198})$$

This indicates that the cosmological constant Λ is effectively screened by ξ . We should also note that in the case without the cosmological constant $\Lambda = 0$, if we choose $\xi_1 = -1$, H_0 can be arbitrary. Then, H_0 can be determined by the initial condition. Because H_0 can be small or large, the theory with the function (II.192) with $\beta = 2$ in (II.196) can describe the early-time inflation or current cosmic acceleration.

In the presence of matter with $w \neq 0$, the de Sitter solution $H = H_0$ is realized even if $f(\eta)$ given by

$$f(\eta) = f_0 e^{\eta/2} + f_1 e^{3(w+1)\eta/4}. \quad (\text{II.199})$$

The following solution exists:

$$\eta = -4H_0 t, \quad \xi = 1 + 3f_0 e^{-2H_0 t} + \frac{f_1}{w} e^{-3(w+1)H_0 t}, \quad \rho_{\text{matter}} = -\frac{3(3w+1)H_0^2 f_1}{\kappa^2} e^{-3(1+w)H_0 t}. \quad (\text{II.200})$$

The theory with multiple de Sitter solutions may describe the unification of inflation with dark energy [64]. Other cosmological aspects of such non-local gravity were studied in refs. [66].

One can construct more general non-local gravity along the above line. Let us start from a general non-local action:

$$S = \int d^4x \sqrt{-g} [F(R, \square R, \square^2 R, \dots, \square^m R, \square^{-1} R, \square^{-2} R, \dots, \square^{-n} R) + \mathcal{L}_{\text{matter}}], \quad (\text{II.201})$$

with m and n being the positive integers. By introducing scalar fields A , B , χ_k , η_k , ($k = 1, 2, \dots, m$), and ξ_l , ϕ_l , ($l = 1, 2, \dots, n$), the action (II.201) can be rewritten in the following form:

$$S = \int d^4x \sqrt{-g} \left[BR - BA + F(A, \eta_1, \eta_2, \dots, \eta_m, \phi_1, \phi_2, \dots, \phi_n) + \partial_\mu \chi \partial^\mu A + \sum_{k=2}^m \partial_\mu \chi_k \partial^\mu \eta_{k-1} \right. \\ \left. + \sum_{l=1}^n \partial_\mu \xi_l \partial^\mu \phi_l + \sum_{k=1}^m \chi_1 \eta_1 + A \xi_1 + \sum_{l=2}^n \xi_l \phi_{l-1} + \mathcal{L}_{\text{matter}} \right]. \quad (\text{II.202})$$

The variations over A , B , χ_k , and ξ_l lead to the following equations

$$0 = R - A = \eta_1 - \square A = \eta_k - \square \eta_{k-1} = \square \phi_1 - A = \square \phi_l - \phi_{l-1} \quad (k = 2, \dots, m, \quad l = 2, \dots, n), \quad (\text{II.203})$$

which give $\eta_k = \square^k R$ and $\square^l \phi_l = R$. Thus, the actions (II.202) and (II.201) are equivalent.

Let us now consider a scale transformation given by

$$g_{\mu\nu} \rightarrow \frac{1}{2B} g_{\mu\nu}. \quad (\text{II.204})$$

Then, the action (II.202) is transformed into the action in the Einstein frame:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2} - \frac{3}{2B^2} \partial_\mu B \partial^\mu B + \frac{1}{2B} \left(\partial_\mu \chi \partial^\mu A + \sum_{k=2}^m \partial_\mu \chi_k \partial^\mu \eta_{k-1} + \sum_{l=1}^n \partial_\mu \xi_l \partial^\mu \phi_l \right) \right. \\ \left. + \frac{1}{4B^2} \left(-BA + F(A, \eta_1, \dots, \eta_m, \phi_1, \dots, \phi_n) + \sum_{k=1}^m \chi_1 \eta_1 + A \xi_1 + \sum_{l=2}^n \xi_l \phi_{l-1} \right) + \mathcal{L}_{\text{matter}}^A \right]. \quad (\text{II.205})$$

Here, $\mathcal{L}_{\text{matter}}^A$ can be obtained by scale transformation of the metric tensor in $\mathcal{L}_{\text{matter}}$ by (II.204). The above action can be regarded as a non-linear σ model coupled to gravity. The action (II.183) can be considered as a particular case of the general non-local action. In this case, however, the continuity equation (II.144) is no longer valid; additional terms on the right-hand side of this equation are generated and are responsible for inducing the interaction of matter within the background. This aspect is crucial for the discussion of local gravity constraints for such models based on the modified theories of gravity.

2. Non-local Gauss-Bonnet gravity

It may be interesting to discuss the extension of the model presented in the above section by including the Gauss-Bonnet invariant [67, 68].

The corresponding non-local action generalizing the modified Gauss-Bonnet gravity is

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{\kappa^2}{2\alpha} \mathcal{G} \square^{-1} \mathcal{G} + \mathcal{L}_{\text{matter}} \right). \quad (\text{II.206})$$

Here, $\mathcal{L}_{\text{matter}}$ is the matter Lagrangian, and $\mathcal{G}$ is the Gauss-Bonnet invariant. By introducing the scalar field ϕ , one may rewrite the action (II.206) in a local form:

$$S = \int d^4x \left(\frac{R}{2\kappa^2} - \frac{\alpha}{2\kappa^2} \partial_\mu \phi \partial^\mu \phi + \phi \mathcal{G} + \mathcal{L}_{\text{matter}} \right). \quad (\text{II.207})$$

In fact, from the ϕ equation, it follows that $\phi = -\frac{\kappa^2}{\alpha} \square^{-1} \mathcal{G}$. By substituting this expression into (II.207), we re-obtain (II.206). It is remarkable that the above action belongs to a specific class of string-inspired, scalar-Gauss-Bonnet gravity that is known to be consistent with the local tests and may serve as a realistic candidate for dark energy [55] (see section C of this chapter). Unlike the situation in standard stringy compactifications, the scalar-Gauss-Bonnet coupling is not an exponential scalar function. The scalar potential is zero, as it should be in the low-energy, perturbative approach to the string effective action.

Let us now study the de Sitter solutions that may describe late-time universe acceleration in such non-local gravity. Considering the FRW metric with a flat spatial part and a ϕ that only depends on the cosmological time, the equations of motion are

$$0 = -\frac{3H^2}{\kappa^2} + \frac{\alpha}{2\kappa^2} \dot{\phi}^2 - 24\dot{\phi}H^3 + \rho_{\text{matter}}, \quad 0 = -\frac{\alpha}{\kappa^2} (\ddot{\phi} + 3H\dot{\phi}) + 24(\dot{H}H^2 + H^4). \quad (\text{II.208})$$

When the matter contribution is neglected, assuming $\dot{\phi}$ and H are constant $\dot{\phi} = c_\phi$, $H = H_0$, one can solve Eqs. (II.208) as follows: $c_\phi = \frac{8\kappa^2}{\alpha} H_0^3$, $H_0^4 = -\frac{3\alpha}{160\kappa^4}$. Therefore, if α is allowed to be negative, there is a de Sitter universe solution, namely

$$H = H_0 = \left(-\frac{3\alpha}{160\kappa^4} \right)^{1/4}. \quad (\text{II.209})$$

On the other hand, a negative α means that the scalar field has a ‘wrong’ kinetic term, like a ghost or phantom. As we will see later in (II.218), the de Sitter space solution corresponds to the case $a \rightarrow \infty$ (late-time acceleration) when matter is included. Even in the early universe, before matter was created, this solution can be used for the description of the inflationary epoch.

Note that the second equation (II.208) yields

$$\left(-\frac{\alpha}{\kappa^2} \dot{\phi} + 8H^2 \right) a^3 = C. \quad (\text{II.210})$$

Here, C is a constant. The action (II.207) is invariant under a constant shift of the scalar field ϕ : $\phi \rightarrow \phi + c$ (c is a constant) because $\mathcal{G}$ is a total derivative. The quantity C is the Noether charge corresponding to the invariance. By using (II.210), we may delete $\dot{\phi}$ in the first equation in (II.208) and obtain

$$0 = -\frac{3H^2}{\kappa^2} - \frac{160\kappa^2 H^6}{\alpha} + \frac{16\kappa^2 C H^3}{\alpha a^3} + \frac{\kappa^2 C^2}{\alpha a^6} + \rho_{\text{matter}}. \quad (\text{II.211})$$

When ρ_{matter} is given by the sum of the contributions from all forms of matter with a constant EoS parameter w_i , one gets $\rho_{\text{matter}}(a) = \sum_i \rho_{0i} a^{-3(w_i+1)}$. Then, we can solve algebraically Eq. (II.211) with respect to H as a function of a : $H = \mathcal{H}(a)$. Using the definition of H , the equation has the form $\dot{a} = a\mathcal{H}(a)$. Thus, $t = \int \frac{da}{a\mathcal{H}(a)}$, which gives, at least formally, the solution of (II.211) and, therefore, the FRW universe.

As a simple case, one may consider $C = 0$. Then, Eq. (II.211) reduces to the following form

$$0 = \mathcal{F}(x) \equiv x^3 + \frac{3\alpha}{160\kappa^4} x - \frac{\alpha \rho_{\text{matter}}(a)}{160\kappa^2}. \quad (\text{II.212})$$

Here, $x \equiv H^2$ and, therefore, $x \geq 0$. Because $\mathcal{F}'(x) = 3\left(x^2 + \frac{\alpha}{160\kappa^4}\right)$, when $\alpha > 0$, $\mathcal{F}(x)$ is a monotonically increasing function of x . On the other hand, when $\alpha < 0$, we find

$$\mathcal{F}'(x_\pm) = 0, \quad \mathcal{F}(x_\pm) = \mp \left(-\frac{\alpha}{160\kappa^4} \right)^{3/2} - \frac{\alpha \rho_{\text{matter}}}{160\kappa^2}, \quad x_\pm \equiv \pm \left(-\frac{\alpha}{160\kappa^4} \right)^{1/2}. \quad (\text{II.213})$$

Because

$$\mathcal{F}(0) = -\frac{\alpha\rho_{\text{matter}}}{160\kappa^2}, \quad (\text{II.214})$$

when $\alpha > 0$, because $\mathcal{F}(0) < 0$ and $\mathcal{F}(x)$ is a monotonically increasing function, there is only one positive solution in (II.212):

$$H^2 = x = \alpha_+ + \alpha_-, \quad \alpha_{\pm}^3 = \frac{\alpha\rho_{\text{matter}}}{320\kappa^2} \left(1 \pm \sqrt{1 + \frac{\alpha}{40\kappa^8\rho_{\text{matter}}^2}} \right). \quad (\text{II.215})$$

We now consider the case that ρ_{matter} comes from a unique kind of matter with a constant EoS parameter w . Then, when $a \rightarrow 0$, one obtains $\alpha_+^3 \rightarrow \frac{\alpha\rho_0}{160\kappa^2} a^{-3(1+w)}$ and $\alpha_- \rightarrow 0$. Therefore, $H \propto a^{-(1+w)/2}$ and $a \propto t^{2/(1+w)}$, which could be compared with the usual FRW universe in Einstein gravity, where $a \propto t^{2/3(1+w)}$. On the other hand, when $a \rightarrow \infty$, Eq. (II.215) implies that $H \propto a^{-3(1+w)/2}$ and, therefore, $a \propto t^{2/3(1+w)}$, which reproduces the result in general relativity. This indicates that such a non-local term becomes less dominant as compared with the Einstein-Hilbert term in the late universe.

We may consider the case when $\alpha < 0$. Because $\mathcal{F}(0) > 0$ in this case, if $\mathcal{F}(x_+) > 0$, there is no positive solution for x . On the other hand, if $\mathcal{F}(x_+) < 0$, that is, when

$$-\alpha > 40\kappa^6\rho_{\text{matter}}^2, \quad (\text{II.216})$$

there are two positive solutions and one negative solution in (II.212), namely

$$x = \tilde{\alpha}_+ + \tilde{\alpha}_-, \quad \tilde{\alpha}_+\xi + \tilde{\alpha}_-\xi^2, \quad \tilde{\alpha}_+\xi^2 + \tilde{\alpha}_-\xi, \quad \tilde{\alpha}_{\pm}^3 = \frac{\alpha\rho_{\text{matter}}}{320\kappa^2} \left(1 \pm i\sqrt{-1 - \frac{\alpha}{40\kappa^8\rho_{\text{matter}}^2}} \right). \quad (\text{II.217})$$

Eq. (II.216) indicates that there is a lower bound in a as $a \geq a_{\min} \equiv \left(-\frac{\alpha}{40\kappa^4\rho_0^2} \right)^{1/6(1+w)}$. Here, it is assumed that $w > -1$. Note that even if $a = a_{\min}$, x and, therefore, H do not vanish, which means that the existence of $a_{\min}$ does not imply the bounce solution. It is likely that a universe with $a = a_{\min}$ could appear, for example, as a quantum correction.

When a is large, we find that $\mathcal{F}(0)$ in (II.214) vanishes and $\alpha_{\pm}$ tends to a constant: $\alpha_{\pm}^3 \rightarrow \pm i\frac{1}{640\kappa^6}\sqrt{-\frac{\alpha^3}{10}}$, which gives

$$H^2 = x = 0, \quad \pm\sqrt{-\frac{3\alpha}{160\kappa^4}}. \quad (\text{II.218})$$

The positive solution of x in (II.218) corresponds to the de Sitter solution (II.209).

When the Einstein-Hilbert term in (II.206) or (II.207) can be neglected, by assuming

$$\phi = \frac{a_{\phi}}{t^2}, \quad H = \frac{h_0}{t} \quad (a = a_0 t^{h_0}), \quad (\text{II.219})$$

Eqs. (II.208) reduce to the following algebraic equations

$$0 = a_{\phi} \left(\frac{\alpha}{\kappa^2} a_{\phi} + 24h_0^3 \right), \quad 0 = (h_0 - 1) \left(\frac{\alpha}{\kappa^2} a_{\phi} + 4h_0^3 \right). \quad (\text{II.220})$$

The equations in (II.220) have a non-trivial solution ($a_{\phi} \neq 0$ and $h_0 \neq 0$): $h_0 = 1$ and $a_{\phi} = -\frac{24\kappa^2}{\alpha}$.

Matter can also be accounted in the model; in that case, the first equation in (II.220) is modified as

$$0 = -\frac{3H^2}{\kappa^2} + \frac{\alpha}{2\kappa^2}\dot{\phi}^2 - 24\dot{\phi}H^3 + \rho_{\text{matter}}. \quad (\text{II.221})$$

When one may neglect the Einstein-Hilbert term in (II.206) or (II.207), and if we assume (II.219), there is a non-trivial solution when $h_0 = \frac{4}{3(w+1)}$. In fact, Eqs. (II.208) reduce to the following algebraic expressions

$$0 = \frac{2\alpha}{\kappa^2} a_{\phi}^2 + \rho_0 a_0^{-3(w+1)} + 48a_{\phi} \left(\frac{4}{3(1+w)} \right)^3, \quad 0 = \left(\frac{4}{3(1+w)} - 1 \right) \left(\frac{\alpha}{\kappa^2} a_{\phi} + 4 \left(\frac{4}{3(1+w)} \right)^3 \right). \quad (\text{II.222})$$

If the matter is radiation with $w = 1/3$, the second equation is satisfied and the first equation acquires the following form: $0 = \frac{2\alpha}{\kappa^2} a_\phi^2 + \rho_0 a_0^{-4} + 48a_\phi$. Then, if $\tilde{D} \equiv (24)^2 - \frac{2\alpha\rho_0}{\kappa^2 a_0^4} > 0$, there are non-trivial solutions: $a_\phi = \frac{\kappa^2}{2\alpha} \left(-24 \pm \sqrt{\tilde{D}} \right)$. On the other hand, when $w \neq 1/3$, we find $a_\phi = -\frac{4\kappa^2}{\alpha} \left(\frac{4}{3(1+w)} \right)^3$, $\rho_0 a_0^{-4} = \frac{160\kappa^2}{\alpha} \left(\frac{4}{3(1+w)} \right)^6$. The second equation requires α to be positive. Thus, a phantom like universe solution from purely non-local gravity, without the R term, has been found. In the same way, in the case where several de Sitter solutions exist one can expect the unified description of inflation with dark energy.

3. Other classical non-local Gauss-Bonnet models and their de Sitter solutions

Another model giving rise to an interesting non-local action is the following

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{\kappa^2}{2a} F(\mathcal{G}) \square^{-1} F(\mathcal{G}) \right], \quad (\text{II.223})$$

where a is a dimensional constant and $F(\mathcal{G})$ is an adequate function of $\mathcal{G}$. By introducing three scalar fields, ϕ , ξ , and η , one can rewrite the action (II.223) in the following form:

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} + \frac{a}{2\kappa^2} \partial_\mu \phi \partial^\mu \phi + \phi F(\eta) + \xi (\eta - \mathcal{G}) \right). \quad (\text{II.224})$$

We may further add a potential $V(\phi)$ to the action

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} + \frac{a}{2\kappa^2} \partial_\mu \phi \partial^\mu \phi + \phi F(\eta) + \xi (\eta - \mathcal{G}) - V(\phi) \right). \quad (\text{II.225})$$

In the FRW universe, this action leads to the following equations

$$\begin{aligned} 0 &= -\frac{3}{\kappa^2} H^2 - \frac{a}{2\kappa^2} \dot{\phi}^2 + 24\xi \dot{H}^3 + V(\phi), \quad 0 = \frac{a}{\kappa^2} (\ddot{\phi} + 3H\dot{\phi}) + F(\eta) - V'(\phi), \\ 0 &= \phi F(\eta) + \xi, \quad 0 = \eta - 24(H^4 + \dot{H}H^2). \end{aligned} \quad (\text{II.226})$$

However, after adding the potential $V(\phi)$, it is difficult to obtain the corresponding non-local action explicitly.

When $V(\phi) = 0$, assuming that $\phi = c_\phi t$, $\eta = \eta_0$, $\xi = c_\xi t$, and $H = H_0$, with constant c_ϕ , η_0 , c_ξ , and H_0 , the equations in (II.226) reduce to the algebraic equations

$$0 = -\frac{3}{\kappa^2} H_0^2 - \frac{a}{2\kappa^2} c_\phi^2 + 24c_\xi H_0^3, \quad 0 = \frac{3a}{\kappa^2} H_0 c_\phi + F(\eta_0), \quad 0 = c_\phi F'(\eta_0) + c_\xi, \quad 0 = \eta_0 - 24H_0^4. \quad (\text{II.227})$$

One can solve Eqs. (II.227) with respect to c_ϕ , c_ξ , and η_0 as follows: $c_\phi = -\frac{\kappa^2 F(24H_0^4)}{3aH_0}$, $c_\xi = -\frac{\kappa^2}{3aH_0} F(24H_0^4) F'(24H_0^4)$, $\eta_0 = 24H_0^4$, and we find

$$0 = -\frac{G_0}{8\kappa^2} - \frac{\kappa^2}{18a} F(G_0)^2 + \frac{\kappa^2}{9a} G_0 F(G_0) F'(G_0). \quad (\text{II.228})$$

Here, $G_0 = 24H_0^4$.

For example, if

$$F(\mathcal{G}) = f_0 \mathcal{G}^2, \quad (\text{II.229})$$

Eq. (II.228) gives

$$0 = G_0 \left(-\frac{1}{8\kappa^2} + \frac{11\kappa^2 f_0^2}{18a} G_0^3 \right), \quad (\text{II.230})$$

which has a trivial solution $G_0 = 0$ corresponding to the flat background and

$$G_0^3 = \frac{9a}{44\kappa^4 f_0^2}, \quad (\text{II.231})$$

which corresponds to the de Sitter universe.

As another example, one can choose

$$F(\mathcal{G})^2 = -g_0(\mathcal{G} + g_1)(\mathcal{G} - g_2). \quad (\text{II.232})$$

Here, g_0 , g_1 , and g_2 are positive constants. For $F(\mathcal{G})$ to be real, the value of $\mathcal{G}$ is restricted as $-g_1 < \mathcal{G} < g_2$. Then, Eq. (II.228) yields

$$0 = G_0^2 + \frac{9a}{4g_0\kappa^2}G_0 + g_1g_2, \quad (\text{II.233})$$

which can be solved as

$$G_0 = G_{0\pm} \equiv \frac{1}{2} \left(-\frac{9a}{4g_0\kappa^2} \pm \sqrt{\left(\frac{9a}{4g_0\kappa^2}\right)^2 + 4g_1g_2} \right). \quad (\text{II.234})$$

The above solution is positive, provided that $a < 0$. For $-g_1 < G_{0\pm} < g_2$, we require

$$g_1 > -\frac{9a}{4\kappa^4g_0}, \quad g_2 > -\frac{9a}{8\kappa^4g_0}. \quad (\text{II.235})$$

Then, as long as Eq. (II.235) is satisfied, there are two solutions describing the de Sitter universe. By choosing a more general $F(\mathcal{G})$, we may find multiple de Sitter universe solutions, among which the solution with the largest value of H_0 could perfectly correspond to the inflation epoch and the one with the smallest value of H_0 to the dark energy era. Note that, in the model (II.232), as long as Eq. (II.235) is satisfied, there is no singularity, for example at $\mathcal{G} = 0$. Thus, a smooth transition could occur and there exists a true possibility to realize matter dominance before late-time acceleration. The important lesson from the above investigation is again the fact that the inflation and dark energy epochs may be natural solutions of the field equations.

E. Gravity non-minimally coupled with the matter Lagrangian

1. Non-minimally coupled scalar theory

It could be interesting to investigate the role of the non-standard non-minimal coupling of modified gravity with the matter Lagrangian in the cosmic acceleration epoch. The corresponding, quite general theory has been suggested in refs. [69, 70] (see also [71]). As an example of such a theory, the following action is considered:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} + f(R) L_d \right\}. \quad (\text{II.236})$$

Here, L_d is a standard matter like action that produces dark energy due to non-trivial coupling with the curvature. In principle, the first term may be more general, for instance, of the $f(R)$ form. However, for illustrative purposes it is enough to consider this simpler variant of the non-minimal theory. The second term in the above action describes the non-linear coupling of matter with gravity. Very often, such terms may be induced by quantum effects such as the non-linear effective action. Thus, it is natural to consider that the second term belongs to matter sector.

By the variation over $g_{\mu\nu}$, the equation of motion follows as:

$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta g_{\mu\nu}} = \frac{1}{2\kappa^2} \left\{ \frac{1}{2} g^{\mu\nu} R - R^{\mu\nu} \right\} + \tilde{T}^{\mu\nu}. \quad (\text{II.237})$$

Here, the effective energy-momentum tensor $\tilde{T}_{\mu\nu}$ is defined by

$$\begin{aligned} \tilde{T}^{\mu\nu} &\equiv -f'(R)R^{\mu\nu}L_d + (\nabla^\mu \nabla^\nu - g^{\mu\nu} \nabla^2)(f'(R)L_d) + f(R)T^{\mu\nu}, \\ T^{\mu\nu} &\equiv \frac{1}{\sqrt{-g}} \frac{\delta}{\delta g_{\mu\nu}} \left(\int d^4x \sqrt{-g} L_d \right). \end{aligned} \quad (\text{II.238})$$

As a matter we consider the massless scalar

$$L_d = -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi. \quad (\text{II.239})$$

Then, the equation given by the variation over ϕ has the following form:

$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \phi} = \frac{1}{\sqrt{-g}} \partial_\mu (f(R) \sqrt{-g} g^{\mu\nu} \partial_\nu \phi) . \quad (\text{II.240})$$

It is enough to consider the case that

$$f(R) = \left(\frac{R}{\mu^2} \right)^\alpha . \quad (\text{II.241})$$

We again assume the spatially flat FRW universe (II.6) and that the scalar ϕ only depends on t ($\phi = \phi(t)$). The solution of the scalar field equation (II.240) is given by

$$\dot{\phi} = qa^{-3} R^{-\alpha} . \quad (\text{II.242})$$

Here, q is a constant of the integration. Thus, $R^\alpha L_d = \frac{q^2}{2a^6 R^\alpha}$, which becomes dominant when R is small (large) compared with the Einstein term $\frac{R}{2\kappa^2}$ if $\alpha > -1$ ($\alpha < -1$). Thus, one arrives at the remarkable possibility [69, 70] that dark energy grows to asymptotic dominance over the usual matter with a decrease of the curvature. In the current universe, this solves the coincidence problem (the equality of the energy density for dark energy and for matter) simply by the fact of universe expansion.

Substituting (II.242) into (II.237), the $(\mu, \nu) = (t, t)$ component of the equation of motion has the following form:

$$\begin{aligned} 0 = & -\frac{3}{2\kappa^2} H^2 + \frac{36q^2}{\mu^{2\alpha} a^6 (6\dot{H} + 12H^2)^{\alpha+2}} \left\{ \frac{\alpha(\alpha+1)}{4} \ddot{H} H + \frac{\alpha+1}{4} \dot{H}^2 \right. \\ & \left. + \left(1 + \frac{13}{4}\alpha + \alpha^2 \right) \dot{H} H^2 + \left(1 + \frac{7}{2}\alpha \right) H^4 \right\} . \end{aligned} \quad (\text{II.243})$$

The accelerated FRW solution of (II.243) exists as [69, 70]

$$a = a_0 t^{\frac{\alpha+1}{3}} \left(H = \frac{\alpha+1}{3t} \right) , \quad a_0^6 \equiv \frac{2\kappa^2 q^2 (2\alpha-1)(\alpha-1)}{\mu^{2\alpha} 3(\alpha+1)^{\alpha+1} \left(\frac{2}{3}(2\alpha-1) \right)^{\alpha+2}} . \quad (\text{II.244})$$

Eq. (II.244) shows that the universe accelerates, that is, $\ddot{a} > 0$ if $\alpha > 2$. If $\alpha < -1$, the solution (II.244) describes a shrinking universe if $t > 0$. If the time is shifted as $t \rightarrow t - t_s$ (t_s is a constant), the accelerating and expanding universe occurs when $t < t_s$. In the solution with $\alpha < -1$ a Big Rip singularity appears at $t = t_s$. For $a \propto t^{h_0}$, the effective EoS parameter w_{eff} in (II.21) is given by $w_{\text{eff}} = -1 + \frac{2}{3h_0}$, and the accelerating universe expansion ($h_0 > 1$) occurs if $-1 < w_{\text{eff}} < -\frac{1}{3}$. For the case of (II.244), one finds

$$w_{\text{eff}} = \frac{1-\alpha}{1+\alpha} . \quad (\text{II.245})$$

Then, if $\alpha < -1$, we have $w_{\text{eff}} < -1$, which is an effective phantom regime. For general matter with the constant EoS parameter w , the energy E and the energy density ρ_{matter} behave as $E \sim a^{-3w}$ and $\rho_{\text{matter}} \sim a^{-3(w+1)}$. Thus, for the standard phantom with $w < -1$, the density becomes large with time and might generate the Big Rip. Thus, it is demonstrated that the non-linear coupling of gravity with matter may produce an accelerated cosmological expansion with the effective EoS parameter close (above, equal or below) to -1 . More complicated couplings [69, 70, 72] may be considered in the same fashion. Note that such theories show some very interesting properties, like the induction of extra gravitational force [73]. Other cosmological aspects of such non-minimally coupled non-linear theories are discussed in refs. [74].

2. Non-minimally coupled vector model

The non-minimal coupling of gravity with the matter Lagrangian of the previous section may be considered for more complicated models. Let us investigate the abelian vector field Lagrangian non-minimally coupled with gravity [75]:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \frac{I(R)}{4} F_{\mu\nu} F^{\mu\nu} \right\} . \quad (\text{II.246})$$

Here, $I(R)$ is an appropriate function of the scalar curvature R , and $F_{\mu\nu}$ is the curvature of the $U(1)$ gauge field. Making the variation of the action Eq. (II.246) with respect to the metric $g_{\mu\nu}$ and the $U(1)$ gauge field A_μ , we obtain the following equations:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa^2 T_{\mu\nu}^{(\text{EM})}, \quad (\text{II.247})$$

$$T_{\mu\nu}^{(\text{EM})} = I(R) \left(g^{\alpha\beta} F_{\mu\beta} F_{\nu\alpha} - \frac{1}{4} g_{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} \right) + \frac{1}{2} \left\{ f'(R) F_{\alpha\beta} F^{\alpha\beta} R_{\mu\nu} + g_{\mu\nu} \square [f'(R) F_{\alpha\beta} F^{\alpha\beta}] - \nabla_\mu \nabla_\nu [f'(R) F_{\alpha\beta} F^{\alpha\beta}] \right\}, \quad (\text{II.248})$$

and

$$-\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} I(R) F^{\mu\nu}) = 0. \quad (\text{II.249})$$

In the FRW universe (II.6), the $(\mu, \nu) = (0, 0)$ component and $(\mu, \nu) = (i, j)$ component of Eq. (II.247) gives

$$H^2 = \frac{\kappa^2}{3} \left\{ I(R) \left(g^{\alpha\beta} F_{0\beta} F_{0\alpha} + \frac{1}{4} F_{\alpha\beta} F^{\alpha\beta} \right) + \frac{3}{2} \left[-f'(R) (\dot{H} + H^2) + 6f''(R) H (\ddot{H} + 4H\dot{H}) \right] F_{\alpha\beta} F^{\alpha\beta} + \frac{3}{2} f'(R) H \frac{d}{dt} (F_{\alpha\beta} F^{\alpha\beta}) - \frac{1}{2} f'(R) \frac{1}{a^2} \Delta^{(3)} (F_{\alpha\beta} F^{\alpha\beta}) \right\}, \quad (\text{II.250})$$

$$2\dot{H} + 3H^2 = \frac{\kappa^2}{2} \left\{ \frac{1}{6} I(R) F_{\alpha\beta} F^{\alpha\beta} + \left[-f'(R) (\dot{H} + 3H^2) + 6f''(R) (\ddot{H} + 7H\dot{H} + 4\dot{H}^2 + 12H^2\dot{H}) + 36f'''(R) (\ddot{H} + 4H\dot{H})^2 \right] F_{\alpha\beta} F^{\alpha\beta} + 3 \left[f'(R) H + 4f''(R) (\ddot{H} + 4H\dot{H}) \right] \frac{d}{dt} (F_{\alpha\beta} F^{\alpha\beta}) + f'(R) \frac{d^2}{dt^2} (F_{\alpha\beta} F^{\alpha\beta}) - \frac{2}{3} f'(R) \frac{1}{a^2} \Delta^{(3)} (F_{\alpha\beta} F^{\alpha\beta}) \right\}. \quad (\text{II.251})$$

By properly choosing $I(R)$, a model unifying the late acceleration and the inflation emerges. However, the properties of the model may be investigated only qualitatively [75] or numerically because the equations of motion are very complicated. Alternatively, as will be shown in the next chapter the cosmological reconstruction of the theory under investigation may be conducted so that unified cosmic history is realized. Different properties of the non-minimally coupled vector theory have been studied in refs. [76].

3. Non-minimal Yang-Mills theory

Here, we consider the cosmology in the non-abelian non-minimal vector- $F(R)$ gravity based on ref. [77]. Different cosmological proposals regarding the non-minimal Yang-Mills theory may be found in refs. [78].

The proposed action follows:

$$\bar{S}_{\text{MG}} = \int d^4x \sqrt{-g} [\mathcal{L}_{\text{MG}} + \mathcal{L}_{\text{V}}], \quad \mathcal{L}_{\text{MG}} = \frac{1}{2\kappa^2} [R + F(R)], \quad \mathcal{L}_{\text{V}} = I(R) \left\{ -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} - V[A^2] \right\}, \quad A^2 \equiv A^{a2} \quad (\text{II.252})$$

where $F_{\mu\nu}^a$ is given by

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c, \quad (\text{II.253})$$

and $A^2 = g^{\mu\nu} A_\mu^a A_\nu^a$. We should note that the last term $V[A^2]$ in the action (II.252) is not gauge invariant, but it can be rewritten in a gauge-invariant way. For example, if the gauge group is a unitary group, one may introduce a σ model like field U , which satisfies $U^\dagger U = 1$. Then, the last term could be rewritten in the gauge-invariant form:

$$V[A^2] \rightarrow V[\text{ctr}(U^\dagger A_\mu^a U)(U^\dagger A^{a\mu} U)]. \quad (\text{II.254})$$

Here, $\bar{c}$ is the normalization constant. Choosing the unitary gauge $U = 1$, the term in (II.254) reduces to the original one: $V[A^2]$. This may indicate that the action (II.252) describes the theory in which the gauge group is spontaneously broken.

The field equations can be derived by taking variations of the action (II.252) with respect to the metric $g_{\mu\nu}$ and the vector field A_μ^a as follows:

$$[1 + F'(R)] R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} [R + F(R)] + g_{\mu\nu} \square F'(R) - \nabla_\mu \nabla_\nu F'(R) = \kappa^2 T_{\mu\nu}^{(V)}, \quad (\text{II.255})$$

with

$$T_{\mu\nu}^{(V)} = I(R) \left\{ g^{\alpha\beta} F_{\mu\beta}^a F_{\nu\alpha}^a + 2A_\mu^a A_\nu^a \frac{dV[A^2]}{dA^2} - \frac{1}{4} g_{\mu\nu} \bar{\mathcal{F}} \right\} + \frac{1}{2} \{ f'(R) \bar{\mathcal{F}} R_{\mu\nu} + g_{\mu\nu} \square [f'(R) \bar{\mathcal{F}}] - \nabla_\mu \nabla_\nu [f'(R) \bar{\mathcal{F}}] \}, \quad (\text{II.256})$$

$$\bar{\mathcal{F}} = F_{\mu\nu}^a F^{a\mu\nu} + 4V[A^2], \quad (\text{II.257})$$

and

$$\frac{1}{\sqrt{-g}} \partial_\mu [\sqrt{-g} I(R) F^{a\mu\nu}] - I(R) \left\{ f^{abc} A_\mu^b F^{c\mu\nu} + 2 \frac{dV[A^2]}{dA^2} A^{a\nu} \right\} = 0, \quad (\text{II.258})$$

where $T_{\mu\nu}^{(V)}$ is the contribution to the energy-momentum tensor from A_μ^a . As will be shown in the next chapter, the cosmological reconstruction of the above theory may result in the accelerating expanding universe. The direct solution of the equations of motion is extremely complicated.

F. Modified $F(R)$ Hořava-Lifshitz gravity

1. General formulation

This section is devoted to a review of the $F(R)$ Hořava-Lifshitz gravity [79]. The FRW equations for this theory are also formulated. The Hořava-Lifshitz version of general relativity was originally introduced as a candidate for renormalizable quantum gravity at the price of the violation of Lorentz invariance. The below model represents natural generalization of the standard $F(R)$ gravity (II.1) discussed at the beginning of this chapter.

Let us present the general formulation of the theory. By using the ADM decomposition [80, 81] (for reviews, see [82, 83]), one can write the metric in the following form:

$$ds^2 = -N^2 dt^2 + g_{ij}^{(3)} (dx^i + N^i dt) (dx^j + N^j dt), \quad i = 1, 2, 3. \quad (\text{II.259})$$

Here, N is called the lapse variable and N^i 's are the shift variables. The scalar curvature R becomes:

$$R = K^{ij} K_{ij} - K^2 + R^{(3)} + 2\nabla_\mu (n^\mu \nabla_\nu n^\nu - n^\nu \nabla_\nu n^\mu), \quad (\text{II.260})$$

and $\sqrt{-g} = \sqrt{g^{(3)}} N$. Here, $R^{(3)}$ is the three-dimensional scalar curvature defined by the metric $g_{ij}^{(3)}$, and K_{ij} is the extrinsic curvature defined by

$$K_{ij} = \frac{1}{2N} \left(\dot{g}_{ij}^{(3)} - \nabla_i^{(3)} N_j - \nabla_j^{(3)} N_i \right), \quad K = K^i_i. \quad (\text{II.261})$$

n^μ is a unit vector perpendicular to the three-dimensional hypersurface Σ_t defined by $t = \text{constant}$ and $\nabla_i^{(3)}$ expresses the covariant derivative on the hypersurface Σ_t .

The first attempt of the extension of $F(R)$ gravity to a Hořava-Lifshitz-type theory [84] which is known to be a power-counting renormalizable model was proposed in ref. [85]. The starting action is

$$S_{F_{\text{HL}}(R)} = \int d^4x \sqrt{g^{(3)}} N F(R_{\text{HL}}), \quad R_{\text{HL}} \equiv K^{ij} K_{ij} - \lambda K^2 - E^{ij} \mathcal{G}_{ijkl} E^{kl}. \quad (\text{II.262})$$

Here, λ is a real constant in the “generalized De Witt metric” or “super-metric”,

$$\mathcal{G}^{ijkl} = \frac{1}{2} \left(g^{(3)ik} g^{(3)jl} + g^{(3)il} g^{(3)jk} \right) - \lambda g^{(3)ij} g^{(3)kl}, \quad (\text{II.263})$$

defined on the three-dimensional hypersurface Σ_t , and E^{ij} can be defined by the so-called “detailed balance condition” by using an action $W[g_{kl}^{(3)}]$ on the hypersurface Σ_t

$$\sqrt{g^{(3)}} E^{ij} = \frac{\delta W[g_{kl}^{(3)}]}{\delta g_{ij}}, \quad (\text{II.264})$$

and the inverse of $\mathcal{G}^{ijkl}$ is written as

$$\mathcal{G}_{ijkl} = \frac{1}{2} \left(g_{ik}^{(3)} g_{jl}^{(3)} + g_{il}^{(3)} g_{jk}^{(3)} \right) - \tilde{\lambda} g_{ij}^{(3)} g_{kl}^{(3)}, \quad \tilde{\lambda} = \frac{\lambda}{3\lambda - 1}. \quad (\text{II.265})$$

The action $W[g_{kl}^{(3)}]$ is assumed to be defined by the metric and the covariant derivatives on the hypersurface Σ_t . The original motivation for the detailed balance condition is its ability to simplify the quantum behavior and renormalization properties of theories that satisfy the detailed balance condition. Otherwise, there is no a priori physical reason to restrict E^{ij} to be defined by (II.264).

There exists an anisotropy between space and time in the Hořava-Lifshitz approach. In the ultraviolet (high-energy) region, the time coordinate and the spatial coordinates are assumed to behave as

$$\mathbf{x} \rightarrow b\mathbf{x}, \quad t \rightarrow b^z t, \quad z = 2, 3, \dots, \quad (\text{II.266})$$

under the scale transformation. In ref. [84], $W[g_{kl}^{(3)}]$ is explicitly given for the case $z = 2$,

$$W = \frac{1}{\kappa_W^2} \int d^3\mathbf{x} \sqrt{g^{(3)}} (R - 2\Lambda_W), \quad (\text{II.267})$$

and for the case $z = 3$,

$$W = \frac{1}{w^2} \int_{\Sigma_t} \omega_3(\Gamma). \quad (\text{II.268})$$

Here, κ_W is a coupling constant of dimension $-1/2$ and w^2 is the dimensionless coupling constant. $\omega_3(\Gamma)$ is given by

$$\omega_3(\Gamma) = \text{Tr} \left(\Gamma \wedge d\Gamma + \frac{2}{3} \Gamma \wedge \Gamma \wedge \Gamma \right) \equiv \varepsilon^{ijk} \left(\Gamma_{il}^m \partial_j \Gamma_{km}^l + \frac{2}{3} \Gamma_{il}^n \Gamma_{jm}^l \Gamma_{kn}^m \right) d^3\mathbf{x}. \quad (\text{II.269})$$

A general E^{ij} consists of all contributions to W up to the chosen value z .

In the Hořava-Lifshitz like $F(R)$ gravity, it is assumed that N can only depend on the time coordinate t , which is called the “projectability condition”. The reason for this is that Hořava-Lifshitz gravity does not have the full diffeomorphism invariance, but rather is invariant only under “foliation-preserving” diffeomorphisms, i.e., under the transformations

$$\delta x^i = \zeta^i(t, \mathbf{x}), \quad \delta t = f(t). \quad (\text{II.270})$$

If N depends on the spatial coordinates, it could not be fixed to be unity ($N = 1$) by using the foliation-preserving diffeomorphisms. There exists a version of Hořava-Lifshitz gravity without the projectability condition, but it is suspected to possess a few additional consistency problems [86, 87]. Therefore, we prefer to assume that N depends only on the time coordinate t .

Let us consider the FRW universe with a flat spatial part and the lapse function N :

$$ds^2 = -N^2 dt^2 + a(t)^2 \sum_{i=1,2,3} (dx^i)^2. \quad (\text{II.271})$$

It is clear from the explicit expressions (II.267) and (II.268) that $W[g_{kl}^{(3)}]$ vanishes identically if $\Lambda_W = 0$, which is assumed because a non-vanishing Λ_W gives a cosmological constant. Then, one can obtain

$$R = \frac{12H^2}{N^2} + \frac{6}{N} \frac{d}{dt} \left(\frac{H}{N} \right) = -\frac{6H^2}{N} + \frac{6}{a^3 N} \frac{d}{dt} \left(\frac{Ha^3}{N} \right), \quad R_{\text{HL}} = \frac{(3 - 9\lambda) H^2}{N^2}. \quad (\text{II.272})$$

In the case of general relativity, the second term in the last expression for R becomes a total derivative:

$$\int d^4x \sqrt{-g} R = \int d^4x a^3 N \left\{ -\frac{6H^2}{N} + \frac{6}{a^3 N} \frac{d}{dt} \left(\frac{Ha^3}{N} \right) \right\} = \int d^4x \left\{ -6H^2 a^3 + 6 \frac{d}{dt} \left(\frac{Ha^3}{N} \right) \right\}. \quad (\text{II.273})$$

Therefore, this term can be dropped in Einstein gravity. The total derivative term comes from the last term $2\nabla_\mu (n^\mu \nabla_\nu n^\nu - n^\nu \nabla_\nu n^\mu)$ in (II.260), which is dropped in the usual Hořava-Lifshitz gravity. In $F(R)$ gravity, however, this term cannot be dropped due to its non-linearity. Then, if we consider FRW cosmology with the flat spatial part, there is almost no qualitative difference between Einstein gravity and Hořava-Lifshitz gravity, except that an effective dark matter could appear as a kind of integration constant in Hořava-Lifshitz gravity [88]. The effective dark matter appears because the constraint given by the variation over N becomes global in the projectable Hořava-Lifshitz gravity.

Now, we introduce very general Hořava-Lifshitz like $F(R)$ gravity [79] by

$$S_{F(\tilde{R})} = \int d^4x \sqrt{g^{(3)}} N F(\tilde{R}), \quad \tilde{R} \equiv K^{ij} K_{ij} - \lambda K^2 + 2\mu \nabla_\mu (n^\mu \nabla_\nu n^\nu - n^\nu \nabla_\nu n^\mu) - E^{ij} \mathcal{G}_{ijkl} E^{kl}. \quad (\text{II.274})$$

In the FRW universe with a flat spatial part, $\tilde{R}$ has the following form:

$$\tilde{R} = \frac{(3 - 9\lambda) H^2}{N^2} + \frac{6\mu}{a^3 N} \frac{d}{dt} \left(\frac{H a^3}{N} \right) = \frac{(3 - 9\lambda + 18\mu) H^2}{N^2} + \frac{6\mu}{N} \frac{d}{dt} \left(\frac{H}{N} \right). \quad (\text{II.275})$$

The case one obtains with the choice of parameters $\lambda = \mu = 1$ corresponds to the usual $F(R)$ gravity as long as we consider spatially flat FRW cosmology, because $\tilde{R}$ reduces to R in (II.272). On the other hand, in the case of $\mu = 0$, $\tilde{R}$ reduces to R_{HL} (II.272) and, therefore, the action (II.274) becomes identical with the action (II.262) of the Hořava-Lifshitz like $F(R)$ gravity in ref. [85]. Thus, the $\mu = 0$ version corresponds to some degenerate limit of the above general $F(R)$ Hořava-Lifshitz gravity. We call this limit degenerate because it is very difficult to obtain the FRW equations when $\mu = 0$ is set from the very beginning. In our theory the FRW equations can be obtained quite easily, and then $\mu = 0$ is a simple limit.

For the action (II.274), the FRW equation given by the variation over $g_{ij}^{(3)}$ has the following form after assuming the FRW space-time (II.271) and setting $N = 1$:

$$0 = F(\tilde{R}) - 2(1 - 3\lambda + 3\mu) (\dot{H} + 3H^2) F'(\tilde{R}) - 2(1 - 3\lambda) H \frac{dF'(\tilde{R})}{dt} + 2\mu \frac{d^2 F'(\tilde{R})}{dt^2} + p_{\text{matter}}. \quad (\text{II.276})$$

On the other hand, the variation over N gives the global constraint:

$$0 = \int d^3x \left[F(\tilde{R}) - 6 \left\{ (1 - 3\lambda + 3\mu) H^2 + \mu \dot{H} \right\} F'(\tilde{R}) + 6\mu H \frac{dF'(\tilde{R})}{dt} - \rho_{\text{matter}} \right], \quad (\text{II.277})$$

after setting $N = 1$. Because N only depends on t , and thus does not depend on the spatial coordinates, we only obtain the global constraint given by the integration. If the standard conservation law (II.144) is used, Eq. (II.276) can be integrated to give

$$0 = F(\tilde{R}) - 6 \left\{ (1 - 3\lambda + 3\mu) H^2 + \mu \dot{H} \right\} F'(\tilde{R}) + 6\mu H \frac{dF'(\tilde{R})}{dt} - \rho_{\text{matter}} - \frac{C}{a^3}. \quad (\text{II.278})$$

Here, C is the integration constant. Using (II.277), one finds $C = 0$. In [88], however, it was claimed that C need not always vanish in a local region, because (II.277) needs to be satisfied throughout the entire universe. In the region $C > 0$, the $C a^{-3}$ term in (II.278) may be regarded as dark matter.

Note that Eq. (II.278) corresponds to the first FRW equation and (II.276) to the second one. Specifically, if we choose $\lambda = \mu = 1$ and $C = 0$, Eq. (II.278) reduces to

$$\begin{aligned} 0 &= F(\tilde{R}) - 6 (H^2 + \dot{H}) F'(\tilde{R}) + 6H \frac{dF'(\tilde{R})}{dt} - \rho_{\text{matter}} \\ &= F(\tilde{R}) - 6 (H^2 + \dot{H}) F'(\tilde{R}) + 36 (4H^2 \dot{H} + \ddot{H}) F''(\tilde{R}) - \rho_{\text{matter}}. \end{aligned} \quad (\text{II.279})$$

This coincides with the corresponding FRW equation for traditional $F(R)$ gravity introduced in the first section of this chapter. Note that in the degenerate $\mu = 0$ case [85], the action (II.274) or (II.262) does not contain any term with second derivatives with respect to the coordinates. Such a term appears in the standard $F(R)$ gravity. The existence of the second derivatives in the usual $F(R)$ gravity induces the third and fourth derivatives in the FRW equation as in

(II.276). Due to such higher derivatives, an extra scalar mode appears, which is often called the scalaron in the usual $F(R)$ gravity. This scalar mode often affects the correction to Newton's law as well as other solar tests. Therefore, such a scalar mode does not appear in the $F(R)$ Hořava-Lifshitz gravity with $\mu = 0$. Thus, we have formulated a general Hořava-Lifshitz $F(R)$ gravity that describes the standard $F(R)$ gravity or its non-degenerate Hořava-Lifshitz extension in a consistent way.

2. FRW cosmology

This section is devoted to the study of the FRW Eqs. (II.276) and (II.277), which admit a de Sitter universe solution. We follow ref. [79]. It is convenient to neglect the matter contribution: $p_{\text{matter}} = \rho_{\text{matter}} = 0$. Assuming $H = H_0$, both of Eqs. (II.276) and (II.277) lead to the same equation

$$0 = F(3(1 - 3\lambda + 6\mu)H_0^2) - 6(1 - 3\lambda + 3\mu)H_0^2 F'(3(1 - 3\lambda + 6\mu)H_0^2), \quad (\text{II.280})$$

as long as the integration constant vanishes ($C = 0$) in Eq. (II.278).

First, we consider the popular case that

$$F(\tilde{R}) \propto \tilde{R} + \beta \tilde{R}^2. \quad (\text{II.281})$$

Then, Eq. (II.280) gives

$$0 = H_0^2 \{1 - 3\lambda + 9\beta(1 - 3\lambda + 6\mu)(1 - 3\lambda + 2\mu)H_0^2\}. \quad (\text{II.282})$$

In the case of usual $F(R)$ gravity, where $\lambda = \mu = 1$ and, therefore, $1 - 3\lambda + 2\mu = 0$, there is formally only a trivial solution $H_0^2 = 0$. For our general case, however, there exists the non-trivial solution

$$H_0^2 = -\frac{1 - 3\lambda}{\beta(1 - 3\lambda + 6\mu)(1 - 3\lambda + 2\mu)}, \quad (\text{II.283})$$

as long as the right-hand side of (II.283) is positive. If the magnitude of this non-trivial solution is small enough, this solution might correspond to the late-time accelerating expansion. Thus, the R^2 term may generate the late-time acceleration. On the other hand, the above solution may serve as an inflationary solution for the early universe (with the corresponding choice of parameters).

Instead of (II.281) one may consider the following model:

$$F(\tilde{R}) \propto \tilde{R} + \beta \tilde{R}^2 + \gamma \tilde{R}^3. \quad (\text{II.284})$$

Eq. (II.280) becomes

$$0 = H_0^2 \left\{ 1 - 3\lambda + 9\beta(1 - 3\lambda + 6\mu)(1 - 3\lambda + 2\mu)H_0^2 + 9\gamma(1 - 3\lambda + 6\mu)^2(5 - 15\lambda + 12\mu)H_0^4 \right\}, \quad (\text{II.285})$$

which has the following two non-trivial solutions,

$$H_0^2 = -\frac{(1 - 3\lambda + 2\mu)\beta}{2(1 - 3\lambda + 6\mu)(5 - 15\lambda + 12\mu)\gamma} \left(1 \pm \sqrt{1 - \frac{4(1 - 3\lambda)(5 - 15\lambda + 12\mu)\gamma}{9(1 - 3\lambda + 2\mu)^2\beta^2}} \right), \quad (\text{II.286})$$

as long as the r.h.s. is real and positive. If

$$\left| \frac{4(1 - 3\lambda)(5 - 15\lambda + 12\mu)\gamma}{9(1 - 3\lambda + 2\mu)^2\beta^2} \right| \ll 1, \quad (\text{II.287})$$

one of the two solutions is much smaller than the other solution. Then, one may regard that the larger solution corresponds to the inflation and the smaller one to the late-time acceleration, similarly to the modified gravity model [2], where such unification had been first proposed. Note that some of the above models may possess the future singularity in the same way as the usual $F(R)$ gravity. However, it would be possible to demonstrate that adding terms with even higher derivatives might cure this singularity, as the addition of the R^2 term did in the standard $F(R)$ gravity. Thus, we demonstrated the qualitative possibility to unify the early-time inflation with the late-time acceleration in the modified Hořava-Lifshitz $F(R)$ gravity.

3. Unified inflation and dark energy in modified Hořava-Lifshitz gravity

Let us consider here some viable $F(\tilde{R})$ gravities that admit the unification [2] of inflation with late-time acceleration. In the traditional $F(R)$ theory, a number of viable models (see [30–32]), which pass all local tests and are able to unify the inflationary and the current cosmic accelerated epochs, have been proposed. Here, following ref. [89], it is shown that this class of models may be extended to the Hořava-Lifshitz gravity. The starting action is

$$F(\tilde{R}) = \tilde{R} + f(\tilde{R}), \quad (\text{II.288})$$

where it is assumed that the term $f(\tilde{R})$ becomes important at cosmological scales, whereas for scales comparable to that of the solar system, the theory becomes linear on $\tilde{R}$. As an example, the following function [31] is considered

$$f(\tilde{R}) = \frac{\tilde{R}^n(\alpha\tilde{R}^n - \beta)}{1 + \gamma\tilde{R}^n}, \quad (\text{II.289})$$

where α , β , and γ are constants and $n > 1$. This theory reproduces the inflationary and cosmic acceleration epochs in convenient $F(R)$ gravity (see ref. [31]). During inflation, it is assumed that the curvature scalar tends to infinity. In this case, the model (II.288), with (II.289), behaves as

$$\lim_{\tilde{R} \rightarrow \infty} F(\tilde{R}) = \alpha\tilde{R}^n. \quad (\text{II.290})$$

Then, by solving the FRW equation (II.278) with $C = 0$, this kind of function yields a power-law solution of the type

$$H(t) = \frac{h_1}{t}, \quad \text{where} \quad h_1 = \frac{2\mu(n-1)(2n-1)}{1-3\lambda+6\mu-2n(1-3\lambda+3\mu)}. \quad (\text{II.291})$$

This solution produces acceleration during the inflationary epoch if the parameters of the theory are properly defined. The acceleration parameter is given by $\frac{\ddot{a}}{a} = h_1(h_1 - 1)/t^2$; thus, for $h_1 > 1$, the inflationary epoch is well reproduced by the model (II.289). Alternatively, the function (II.289) has a minimum at $\tilde{R}_0$, given by

$$\tilde{R}_0 \sim \left(\frac{\beta}{\alpha\gamma}\right)^{1/4}, \quad f'(\tilde{R}) = 0, \quad f(\tilde{R}) = -2\Lambda \sim -\frac{\beta}{\gamma}, \quad (\text{II.292})$$

where the condition $\beta\gamma/\alpha \gg 1$ is imposed. Then, at the current epoch, the scalar curvature acquires a small value that can be fixed to coincide with the minimum (II.292), such that the FRW equations (II.276) and (II.278) with $C = 0$ yield

$$H^2 = \frac{\kappa^2}{3(3\lambda-1)}\rho_{\text{matter}} + \frac{2\Lambda}{3(3\lambda-1)}, \quad \dot{H} = -\kappa^2 \frac{\rho_{\text{matter}} + p_{\text{matter}}}{3\lambda-1}, \quad (\text{II.293})$$

which look very similar to the standard FRW equations in general relativity, except for the parameter λ . (For a first consideration of the FRW dynamics in Hořava-Lifshitz gravity based on general relativity, see refs. [90–111].) As has been pointed out, at the current epoch, the scalar $\tilde{R}$ is small, so the theory is in the IR limit where the parameter $\lambda \sim 1$, and the equations approach the usual equations for $F(R)$ gravity. Thus, the FRW equations (II.293) reproduce the behavior of the well-known Λ CDM model with no need to introduce a dark energy fluid to explain the current universe acceleration.

As another example of the models described by (II.288), one can consider the function [15, 30, 32],

$$f(\tilde{R}) = -\frac{(\tilde{R} - \tilde{R}_0)^{2n+1} + \tilde{R}_0^{2n+1}}{f_0 + f_1[(\tilde{R} - \tilde{R}_0)^{2n+1} + \tilde{R}_0^{2n+1}]} = -\frac{1}{f_1} + \frac{f_0/f_1}{f_0 + f_1[(\tilde{R} - \tilde{R}_0)^{2n+1} + \tilde{R}_0^{2n+1}]}. \quad (\text{II.294})$$

This function could also serve as the unification of inflation and cosmic acceleration; however, in this case, when one takes the limit $\tilde{R} \rightarrow \infty$, one gets

$$\lim_{\tilde{R} \rightarrow \infty} F(\tilde{R}) = \tilde{R} - 2\Lambda_i, \quad \text{where} \quad \Lambda_i = 1/2f_1, \quad (\text{II.295})$$

where the subscript i denotes the inflationary. By inserting this into Eqs. (II.276) and (II.278) with $C = 0$, the FRW equations take the same form as in (II.293). Then, for the function (II.294), the inflationary epoch is produced by an

effective cosmological constant, which implies that the parameter $\lambda > 1/3$, or the equations themselves will present inconsistencies, as was discussed in the above section. For the current epoch, it is easy to see that the function (II.294) exhibits a minimum for $\tilde{R} = \tilde{R}_0$, which implies, as in the model above, an effective cosmological constant for late time that can produce the cosmic acceleration. The emergence of matter dominance before the dark energy epoch can be exhibited, in analogy with the case of the convenient theory. Thus, we have shown that the model (II.294) also unifies the cosmic expansion history, although with different properties during the inflationary epoch as compared with the model (II.289).

It is also interesting to explore the de Sitter solutions allowed by the theory (II.262). By taking $H(t) = H_0$, the FRW equation (II.278), in the absence of any kind of matter and with $C = 0$, reduces to

$$0 = F(\tilde{R}_0) - 6H_0^2(1 - 3\lambda + 3\mu)F'(\tilde{R}_0), \quad (\text{II.296})$$

which represents an algebraic equation that can be solved to yield the possible de Sitter points allowed by the theory. As an example, let us consider the model (II.289), where Eq. (II.296) takes the form

$$\tilde{R}_0 + \frac{\tilde{R}_0^n(\alpha\tilde{R}_0^n - \beta)}{1 + \gamma\tilde{R}_0^n} + \frac{6H_0^2(-1 + 3\lambda - 3\mu) \left[1 + n\alpha\gamma\tilde{R}_0^{3n-1} + \tilde{R}_0^{n-1}(2\gamma\tilde{R}_0 - n\beta) + \tilde{R}_0^{2n-1}(\gamma^2\tilde{R}_0 + 2n\alpha) \right]}{(1 + \gamma\tilde{R}_0^n)^2} = 0. \quad (\text{II.297})$$

Here, $\tilde{R}_0 = 3(1 - 3\lambda + 6\mu)H_0^2$. By specifying the free parameters of the theory, one can solve Eq. (II.297), which yields several de Sitter points, as the one studied above. The de Sitter points can be used to explain the coincidence problem, with the argument that the present will not be the only late-time accelerated epoch experienced by our universe. In standard $F(R)$ gravity, it was found that this model contains at least two de Sitter points along the cosmic history (see [113]). In the same way, the second model studied here (II.294) provides several de Sitter points in the course of cosmic history. Note that when $\mu = \lambda - \frac{1}{3}$, Eq. (II.296) turns out to be much more simple; it reduces to $F(\tilde{R}_0) = 0$, where the de Sitter points are the roots. For example, for (II.297) we have $\tilde{R}_0(1 + \gamma\tilde{R}_0^n) + \tilde{R}_0^n(\alpha\tilde{R}_0^n - \beta) = 0$, where the number of positive roots (de Sitter points) depends on the free parameters of the theory.

In summary, it has been demonstrated here that, as in $F(\tilde{R})$ Hořava-Lifshitz gravity, the so-called viable models, as in (II.289) or (II.294), can in fact reproduce the whole cosmological history of the universe, in the same way as convenient modified gravity.

G. Covariant power-counting renormalizable gravity

Motivated by the idea of ref. [84], a model of covariant, power-counting renormalizable gravity was proposed [114, 115]. Its prototype using the perfect fluid is given in ref. [100]. Such a theory seems to be more natural than models [79, 84] because Lorentz invariance is not broken explicitly. Moreover, the theory looks very similar to R^2 gravity with a scalar. The present section is devoted to review of such covariant model in the FRW universe.

1. Perfect fluid formulation

Let us briefly review the covariant, power-counting renormalizable gravity of ref. [100]. The starting action is

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \alpha (T^{\mu\nu} R_{\mu\nu} + \beta TR)^2 \right\}. \quad (\text{II.298})$$

Here, $T_{\mu\nu}$ is the energy-momentum tensor of some string-inspired fluid. The action (II.298) is fully diffeomorphism invariant. We consider the perturbation from the flat background $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$. The following gauge conditions are chosen: $h_{tt} = h_{ti} = h_{it} = 0$. The fluid energy-momentum tensor in the flat background has the following form:

$$T_{tt} = \rho, \quad T_{ij} = p\delta_{ij} = w\rho\delta_{ij}. \quad (\text{II.299})$$

Here, w is the EoS parameter. Then, one finds

$$\begin{aligned} & T^{\mu\nu} R_{\mu\nu} + \beta TR \\ &= \rho \left[\left\{ -\frac{1}{2} + \frac{w}{2} + (-1 + 3w)\beta \right\} \partial_t^2 (\delta^{ij} h_{ij}) + (w - \beta + 3w\beta) \partial^i \partial^j h_{ij} \right. \\ & \quad \left. + (-w + \beta - 3w\beta) \partial_k \partial^k (\delta^{ij} h_{ij}) \right]. \end{aligned} \quad (\text{II.300})$$

If

$$\beta = -\frac{w-1}{2(3w-1)}, \quad (\text{II.301})$$

the second term in the action (II.298) becomes

$$\alpha (T^{\mu\nu} R_{\mu\nu} + \beta TR)^2 = \alpha \rho^2 \left(\frac{w}{2} + \frac{1}{2} \right)^2 \left\{ \partial^i \partial^j h_{ij} - \partial_k \partial^k (\delta^{ij} h_{ij}) \right\}^2, \quad (\text{II.302})$$

which does not contain the derivative with respect to t and breaks the Lorentz invariance. Imagine ρ is almost constant. Then, in the ultraviolet region, in which $\mathbf{k}$ is large, the second term in the action (II.298) gives the propagator behaving as $1/|\mathbf{k}|^4$. It renders the ultraviolet behavior (compared with Eq. (1.4) in [84]). Note that the form (II.301) indicates that the longitudinal mode does not propagate; only the transverse mode propagates.

The action (II.298) gives $z = 2$ theory. For the theory to be ultraviolet, power-counting renormalizable in $3 + 1$ dimensions, $z = 3$ theory is necessary. To obtain such a theory, we note that, for any scalar quantity Φ with the choice

$$\gamma = \frac{1}{3w-1}, \quad (\text{II.303})$$

one obtains

$$T^{\mu\nu} \nabla_\mu \nabla_\nu \Phi + \gamma T \nabla^\rho \nabla_\rho \Phi = \rho (w+1) \partial_k \partial^k \Phi, \quad (\text{II.304})$$

which does not contain the derivative with respect to the time coordinate t . This is true even if the coordinate frame is not the local Lorentz frame. The derivative with respect to the time coordinate t is not contained in any coordinate frame where the perfect fluid does not flow. Thus, the proposed action looks as

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \alpha (T^{\mu\nu} R_{\mu\nu} + \beta TR) (T^{\mu\nu} \nabla_\mu \nabla_\nu + \gamma T \nabla^\rho \nabla_\rho) (T^{\mu\nu} R_{\mu\nu} + \beta TR) \right\}, \quad (\text{II.305})$$

with $\beta = -\frac{w-1}{2(3w-1)}$ and $\gamma = \frac{1}{3w-1}$. It corresponds to $z = 3$ theory, which seems to be (power-counting) renormalizable. In general, for the case

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \alpha \{ (T^{\mu\nu} \nabla_\mu \nabla_\nu + \gamma T \nabla^\rho \nabla_\rho)^n (T^{\mu\nu} R_{\mu\nu} + \beta TR) \}^2 \right], \quad (\text{II.306})$$

with a constant n , $z = 2n + 2$ theory, which is super-renormalizable for $n \geq 1$ emerges.

The second terms in the actions (II.298), (II.305), and (II.306), which effectively break the Lorentz symmetry, are relevant only in the high-energy/UV region because they contain higher-derivative terms. In the IR region, these terms do not dominate, and the usual Einstein gravity follows as a limit.

2. Covariant field theory of gravity with a Lagrange multiplier

We now show that the perfect fluid, which appeared in (II.298), (II.305), and (II.306), can be realized by a scalar field and the Lagrange multiplier field. As a result, we obtain a covariant, power-counting renormalizable field theory of gravity.

The following constrained action for the scalar field ϕ is proposed:

$$S_\phi = \int d^4x \sqrt{-g} \left\{ -\lambda \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + U(\phi) \right) \right\}. \quad (\text{II.307})$$

Here, λ is the Lagrange multiplier field, which gives a constraint of

$$\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + U(\phi) = 0, \quad (\text{II.308})$$

that is, the vector $(\partial_\mu \phi)$ is time like. At least locally, one can choose the direction of time to be parallel to $(\partial_\mu \phi)$. Then, Eq. (II.308) has the following form:

$$\frac{1}{2} \left(\frac{d\phi}{dt} \right)^2 = U(\phi). \quad (\text{II.309})$$

The equation given by the variation over ϕ will be discussed later.

The scalar field energy-momentum tensor $T_{\mu\nu}^\phi$ with an arbitrary scalar potential is defined as follows:

$$T_{\mu\nu}^\phi = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} \partial_\rho \phi \partial^\rho \phi + V(\phi) \right). \quad (\text{II.310})$$

The “energy density” ρ_ϕ and “pressure” p_ϕ become:

$$p_\phi = \frac{1}{2} \left(\frac{d\phi}{dt} \right)^2 - V(\phi) = U(\phi) - V(\phi), \quad \rho_\phi = \frac{1}{2} \left(\frac{d\phi}{dt} \right)^2 + V(\phi) = U(\phi) + V(\phi). \quad (\text{II.311})$$

Here, the constraint (II.309) is used. Note that (unknown, for the moment) $V(\phi)$ is not identical with $U(\phi)$: $V(\phi) \neq U(\phi)$. In the case that $V(\phi) = U(\phi)$, Eq. (II.311) shows that $p_\phi = 0$, which corresponds to dust with $w_\phi \equiv p_\phi/\rho_\phi = 0$. Of course, the classical and quantum dynamics of such constrained theories are quite non-trivial [117].

For simplicity, we choose $V(\phi)$ and $U(\phi)$ to be constants:

$$U(\phi) = U_0, \quad V(\phi) = V_0. \quad (\text{II.312})$$

Then, if $U_0 = V_0$, the EoS parameter w_ϕ vanishes. In the general case, one has $w_\phi = \frac{U_0 - V_0}{U_0 + V_0}$. Let us now use $T_{\mu\nu}^\phi$ as an energy-momentum tensor in the previous section. Eq.(II.301) shows $\beta = -\frac{w-1}{2(3w-1)} = \frac{V_0}{2U_0 - 4V_0}$. One can simplify

$$T^{\phi\mu\nu} R_{\mu\nu} + \beta T^\phi R = \partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R. \quad (\text{II.313})$$

Here, Eqs. (II.308), (II.310), and (II.312) are used. Similarly, γ in (II.303) has the following form: $\gamma = \frac{U_0 - V_0}{2U_0 - 4V_0}$, which gives, by using (II.304),

$$T^{\phi\mu\nu} \nabla_\mu \nabla_\nu \Phi + \gamma T^\phi \nabla^\rho \nabla_\rho \Phi = \partial^\mu \phi \partial^\nu \phi \nabla_\mu \nabla_\nu \Phi + 2U_0 \nabla^\rho \nabla_\rho \Phi. \quad (\text{II.314})$$

Eq. (II.313) enables us to write down the $z = 2$ total action corresponding to (II.298) as

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \alpha (\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R)^2 - \lambda \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + U_0 \right) \right\}. \quad (\text{II.315})$$

On the other hand, the $z = 3$ total action corresponding to (II.305) is:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \alpha (\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R) (\partial^\mu \phi \partial^\nu \phi \nabla_\mu \nabla_\nu + 2U_0 \nabla^\rho \nabla_\rho) (\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R) - \lambda \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + U_0 \right) \right\}, \quad (\text{II.316})$$

and

$$S_{2n+2} = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \alpha \{ (\partial^\mu \phi \partial^\nu \phi \nabla_\mu \nabla_\nu + 2U_0 \nabla^\rho \nabla_\rho)^n (\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R) \}^2 - \lambda \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + U_0 \right) \right\}. \quad (\text{II.317})$$

for the $z = 2n + 2$ model ($n = 0, 1, 2, \dots$), and

$$S_{2n+3} = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - \alpha \{ (\partial^\mu \phi \partial^\nu \phi \nabla_\mu \nabla_\nu + 2U_0 \nabla^\rho \nabla_\rho)^n (\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R) \} \times \left\{ (\partial^\mu \phi \partial^\nu \phi \nabla_\mu \nabla_\nu + 2U_0 \nabla^\rho \nabla_\rho)^{n+1} (\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R) \right\} - \lambda \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + U_0 \right) \right\}, \quad (\text{II.318})$$

for the $z = 2n + 3$ model ($n = 0, 1, 2, \dots$) (compare with [79])³. Note that the actions (II.315), (II.316), (II.317), and (II.318) are totally diffeomorphism invariant and are only given in terms of the local fields. The actions (II.315), (II.316), (II.317), and (II.318) do not depend on V_0 .

³ Note that the model (II.317) or (II.318) has better renormalization properties and no scalar graviton when the action is modified a little bit and the scalar projector is inserted in second term [116].

By the variation over ϕ , for example for the $z = 2$ ($n = 0$) case in (II.317), one finds

$$0 = 4\alpha\partial^\mu \{ \partial^\nu \phi R_{\mu\nu} (\partial^\rho \phi \partial^\sigma \phi R_{\rho\sigma} + U_0 R) \} + \partial^\mu (\lambda \partial_\mu \phi) . \quad (\text{II.319})$$

For the $z \geq 3$ ($n \geq 1$ case in (II.317) or $n \geq 0$ case in (II.318)), one obtains a rather complicated equation.

The dispersion relation of the graviton is now given by

$$\omega = \alpha c_0 k^z , \quad (\text{II.320})$$

in the high-energy region. Here, c_0 is a constant, ω is the angular frequency corresponding to the energy and k is the wave number corresponding to momentum. If $\alpha < 0$, the dispersion relation becomes inconsistent and, therefore, α should be positive.

3. FRW cosmology

The gravitational terms different from general relativity in (II.317) and (II.318) are relevant in the high-energy region. Such terms might affect the inflationary era. In this section, we briefly study FRW cosmology within the theory under discussion. To obtain the FRW equations, the following form of the metric is assumed:

$$ds^2 = -e^{2b(t)} dt^2 + a(t)^2 \sum_{i=1,2,3} (dx^i)^2 , \quad (\text{II.321})$$

and it is assumed that the scalar field ϕ only depends on time. Then, Eq. (II.308) looks as:

$$\frac{1}{2} \left(\frac{d\phi}{dt} \right)^2 = e^{2b(t)} U_0 . \quad (\text{II.322})$$

Thus, one gets

$$\partial^\mu \phi \partial^\nu \phi R_{\mu\nu} + U_0 R = 6U_0 e^{-2b} H^2 , \quad \partial^\mu \phi \partial^\nu \phi \nabla_\mu \nabla_\nu + 2U_0 \nabla^\rho \nabla_\rho = -6U_0 e^{-2b} H \partial_t , \quad (\text{II.323})$$

and the actions (II.317) and (II.318) have the following form:

$$S_{2n+2} = \int d^4 x a^3 \left[\frac{e^{-b}}{2\kappa^2} (6\dot{H} + 12H^2 - 6\dot{b}H) - (6U_0)^{2n+2} e^b \left\{ (e^{-2b} H \partial_t)^n (H^2 e^{-2b}) \right\}^2 - \lambda \left(-\frac{e^{-b}}{2} \left(\frac{d\phi}{dt} \right)^2 + e^b U_0 \right) \right] , \quad (\text{II.324})$$

$$S_{2n+3} = \int d^4 x a^3 \left[\frac{e^{-b}}{2\kappa^2} (6\dot{H} + 12H^2 - 6\dot{b}H) - 2^{2n+3} \cdot 3^2 \alpha U_0^{2n+2} e^b \left\{ (e^{-2b} H \partial_t)^n (H^2 e^{-2b}) \right\} \left\{ (e^{-2b} H \partial_t)^{n+1} (H^2 e^{-2b}) \right\} - \lambda \left(-\frac{e^{-b}}{2} \left(\frac{d\phi}{dt} \right)^2 + e^b U_0 \right) \right] . \quad (\text{II.325})$$

The first FRW equation looks as follows:

$$0 = \frac{3}{\kappa^2} H^2 - (6U_0)^{2n+2} \alpha a^{-3} \left[a^3 (D^n (H^2))^2 - 4(-1)^n H^2 \bar{D}^n (a^3 D^n (H^2)) - 4 \sum_{k=0}^{n-1} (-1)^k (\bar{D}^{n-k} (H^2)) (\bar{D}^k (a^3 D^n (H^2))) \right] - 2\lambda U_0 - \rho_{\text{matter}} , \quad (\text{II.326})$$

for (II.324) and as follows:

$$0 = \frac{3}{\kappa^2} H^2 - (6U_0)^{2n+3} \alpha a^{-3} \left[a^3 (D^n (H^2)) (D^{n+1} (H^2)) - 2(-1)^n H^2 \bar{D}^n (a^3 D^{n+1} (H^2)) - 2(-1)^{n+1} H^2 \bar{D}^{n+1} (a^3 D^n (H^2)) - 2 \sum_{k=0}^{n-1} (-1)^k (\bar{D}^{n-k} (H^2)) (\bar{D}^k (a^3 D^{n+1} (H^2))) - 2 \sum_{k=0}^n (-1)^k (\bar{D}^{n+1-k} (H^2)) (\bar{D}^k (a^3 D^n (H^2))) \right] - 2\lambda U_0 - \rho_{\text{matter}} , \quad (\text{II.327})$$

for (II.325). We have also put $b = 0$ after the variation over b , where the metric (II.321) reduces to the standard FRW metric and the operations of D and $\bar{D}$ for a scalar φ are defined by

$$D\varphi \equiv H \frac{d\varphi}{dt}, \quad \bar{D}\varphi \equiv \frac{d(H\varphi)}{dt}. \quad (\text{II.328})$$

On the other hand, by the variation over a , we get

$$\begin{aligned} 0 = & \frac{1}{\kappa^2} \left(2\dot{H} + 3H^2 \right) + 2^{2n+2} 3^{2n+1} \alpha U_0^{2n+2} \left\{ -3 \left(D^n (H^2) \right)^2 + 4 (-1)^n a^{-3} \frac{d}{dt} \left(H \bar{D}^n (a^3 \bar{D}^n (H^2)) \right) \right. \\ & \left. + 2 \sum_{k=1}^n a^{-3} \frac{d}{dt} \left(\left(\frac{d}{dt} (D^{n-k} (H^2)) \right) (\bar{D}^{k-1} (a^3 D^n (H^2))) \right) \right\} + p_{\text{matter}}, \end{aligned} \quad (\text{II.329})$$

for (II.324) and

$$\begin{aligned} 0 = & \frac{1}{\kappa^2} \left(2\dot{H} + 3H^2 \right) + 2^{2n+3} 3^{2n+2} \alpha U_0^{2n+2} \left\{ -3 \left(D^n (H^2) \right) \left(D^{n+1} (H^2) \right) \right. \\ & + 2 (-1)^n a^{-3} \frac{d}{dt} \left(H \bar{D}^n (a^3 \bar{D}^{n+1} (H^2)) \right) + 2 (-1)^{n+1} a^{-3} \frac{d}{dt} \left(H \bar{D}^{n+1} (a^3 \bar{D}^n (H^2)) \right) \\ & + \sum_{k=1}^n a^{-3} \frac{d}{dt} \left(\left(\frac{d}{dt} (D^{n-k} (H^2)) \right) (\bar{D}^{k-1} (a^3 D^{n+1} (H^2))) \right) \\ & \left. + \sum_{k=1}^{n+1} a^{-3} \frac{d}{dt} \left(\left(\frac{d}{dt} (D^{n-k+1} (H^2)) \right) (\bar{D}^{k-1} (a^3 D^n (H^2))) \right) \right\} + p_{\text{matter}}, \end{aligned} \quad (\text{II.330})$$

for (II.325). The simplest case is $n = 0$ in (II.324) when the FRW equations are

$$\frac{3}{\kappa^2} H^2 = -108 \alpha U_0^2 H^4 + 2 \lambda U_0 + \rho_{\text{matter}}, \quad (\text{II.331})$$

$$-\frac{1}{\kappa^2} (2\dot{H} + 3H^2) = 36 \alpha U_0^2 (3H^4 + 4H^2 \dot{H}) + p_{\text{matter}}. \quad (\text{II.332})$$

In the early universe, in which the curvature was large, the contribution from the Einstein term, which corresponds to the right-hand side in (II.331) and (II.332), and the contributions from the matter ρ_{matter} and p_{matter} , can be neglected. Then, a solution of (II.332) is given by

$$H = \frac{4}{3t}, \quad (\text{II.333})$$

which expresses the (power-law) accelerated universe expansion corresponding to that with a perfect fluid with $w = -1/2$. Eq. (II.331) gives

$$\lambda = \frac{32 \alpha U_0}{3t^4}. \quad (\text{II.334})$$

For more complicated versions of the above theory, one can expect to find a more rich class of background cosmological solutions. The cosmological reconstruction may also be used for such models.

H. Dark fluid with an inhomogeneous equation of state

Usually, a perfect fluid with a constant EoS parameter w is discussed in the cosmological-related literature. One may, however, consider a general equation of state fluid:

$$p = w \left(\rho, a, H, \dot{H}, \ddot{H}, \dots \right) \rho + \left(\rho, a, H, \dot{H}, \ddot{H}, \dots \right). \quad (\text{II.335})$$

It is clear that any modified gravity may be presented as such an effective gravitational fluid coupled with general relativity as was already mentioned in section A of the present chapter. Such an approach to modified gravity can make things simpler when background evolution is investigated. However, it may lead to a number of problems in the

study of cosmological perturbation theory or structure formation. Indeed, when developing such an approach, one often forgets that the initial modified gravity is a higher-derivative theory and that perturbation theory should also be a higher-derivative theory. Non-linearities are also effectively dropped. That is why we prefer to work with the explicit modified gravity formulation but not with general relativity coupled to gravitational fluid.

The above fluid formulation is very general. Moreover, even if we restrict it to the case that the pressure p depends only on the energy density ρ , the EoS fluid, which arbitrarily reproduced in FRW cosmology [118], may be constructed. Needless to say, the unification of inflation with the dark energy epoch may already be achieved for such a fluid. The FRW equations (II.3) show that, for the perfect fluid with the generalized equation of state

$$p = -\rho - \frac{2}{\kappa^2} f' \left(f^{-1} \left(\kappa \sqrt{\frac{\rho}{3}} \right) \right), \quad (\text{II.336})$$

the solution of the FRW equations is given by $H = f(t)$. Here, $f(t)$ is an arbitrary function of the cosmological time t . Therefore, the arbitrary universe evolution given by $H = f(t)$ can be realized via the perfect fluid whose equation of state is given by (II.336).

It is often convenient to use e-folding $N = \ln \frac{a}{a_0}$ instead of the cosmological time t because e-folding is related with the redshift z , which is directly given by observations (for more details, see the next chapter). One may consider the reconstruction using N instead of the cosmological time t . Then, the second equation (II.3) can be rewritten as

$$p_{\text{total}} = -\frac{1}{\kappa^2} (2HH' + 3H^2). \quad (\text{II.337})$$

Here, $'$ expresses the derivative with respect to N : $' \equiv d/dN$. Thus, the fluid with the generalized equation of state

$$p = -\rho - \frac{2}{\kappa} \tilde{f}' \left(\tilde{f}^{-1} \left(\kappa \sqrt{\frac{\rho}{3}} \right) \right) \sqrt{\frac{\rho}{3}}, \quad (\text{II.338})$$

gives the solution of the FRW equations by $H = \tilde{f}(N)$.

It was already mentioned in section A that a phantom fluid whose EoS parameter w is less than -1 leads to the future Big Rip singularity [1]. Several types of softer future singularities may occur for an effective quintessence fluid. The classification of four future finite-time singularities is given in section A following ref. [34].

The Type I singularity corresponds to the Big Rip singularity [1, 33], which emerges when $w < -1$. The Type II singularity corresponds to the sudden future singularity [35] at which a and ρ are finite but p diverges. The Type III singularity appears, for instance, for the model with $p = -\rho - A\rho^\alpha$ [119], which is different from the sudden future singularity in the sense that ρ diverges. This type of singularity was discovered in the model of ref. [119], where the corresponding Lagrangian model of a scalar field with a potential was constructed.

Let us show that all above four types of future singularities may naturally occur for fluid dark energy. As an example, one may start from the following fluid dark energy

$$p = -\rho - f(\rho), \quad (\text{II.339})$$

where $f(\rho)$ is generally an arbitrary function. The case of $f(\rho) \propto \rho^\alpha$ with a constant α was proposed and investigated in detail in ref. [119]. Using the conservation law (II.144) for such a choice, the scale factor a is given by

$$a = a_0 \exp \left(\frac{1}{3} \int \frac{d\rho}{f(\rho)} \right). \quad (\text{II.340})$$

Using the first equation (II.3), the cosmological time t is

$$t = \int \frac{d\rho}{\kappa \sqrt{3\rho f(\rho)}}. \quad (\text{II.341})$$

In the case

$$f(\rho) = A\rho^\alpha, \quad (\text{II.342})$$

by using Eq. (II.340), it follows that

$$a = a_0 \exp \left[\frac{\rho^{1-\alpha}}{3(1-\alpha)A} \right]. \quad (\text{II.343})$$

When $\alpha > 1$, the scale factor remains finite even if ρ goes to infinity. When $\alpha < 1$, $a \rightarrow \infty$ ($a \rightarrow 0$) as $\rho \rightarrow \infty$ for $A > 0$ ($A < 0$). Because the pressure is now given by

$$p = -\rho - A\rho^\alpha, \quad (\text{II.344})$$

p always diverges when ρ becomes infinite. If $\alpha > 1$, the EoS parameter $w = p/\rho$ also goes to infinity, that is, $w \rightarrow +\infty$ ($-\infty$) for $A < 0$ ($A > 0$). When $\alpha < 1$, we have $w \rightarrow -1 + 0$ ($-1 - 0$) for $A < 0$ ($A > 0$) as $\rho \rightarrow \infty$.

Using Eq. (II.341) for (II.342), one finds [34]

$$t = t_s + \frac{2}{\sqrt{3\kappa A}} \frac{\rho^{-\alpha+1/2}}{1-2\alpha}, \quad \text{for } \alpha \neq \frac{1}{2}, \quad (\text{II.345})$$

and

$$t = t_s + \frac{\ln\left(\frac{\rho}{\rho_0}\right)}{\sqrt{3\kappa A}}, \quad \text{for } \alpha = \frac{1}{2}. \quad (\text{II.346})$$

Therefore if $\alpha \leq 1/2$, ρ diverges into an infinite future or past. On the other hand, if $\alpha > 1/2$, the divergence of ρ corresponds to a finite future or past. In the case of a finite future, the singularity could be regarded as a Big Rip or a Type I singularity (compare with the case in section A).

For the equation of state (II.342), the following cases were discussed [34]:

- In the case $\alpha = 1/2$ or $\alpha = 0$, there does not appear any singularity.
- In the case $\alpha > 1$, Eq. (II.345) indicates that when $t \rightarrow t_s$, the energy density behaves as $\rho \rightarrow \infty$ and, therefore, $|p| \rightarrow \infty$ due to (II.344). Eq. (II.343) shows that the scale factor a is finite even if $\rho \rightarrow \infty$. Therefore, the $\alpha > 1$ case corresponds to a Type III singularity.
- The $\alpha = 1$ case corresponds to $p = w\rho$ with a constant w if we replace $-1 - A$ with w . Therefore, if $A > 0$, either the Big Rip or a Type I singularity occurs, but if $A \leq 0$, no future singularity appears.
- In the case that $1/2 < \alpha < 1$, when $t \rightarrow t_s$, all of ρ , $|p|$, and a diverge if $A > 0$; this corresponds to a Type I singularity.
- In the case that $0 < \alpha < 1/2$, when $t \rightarrow t_s$, we find ρ , $|p| \rightarrow 0$ and $a \rightarrow a_0$, but by combining (II.343) and (II.345), we find

$$\ln a \sim |t - t_s|^{\frac{\alpha-1}{\alpha-1/2}}. \quad (\text{II.347})$$

Because the exponent $(\alpha - 1)/(\alpha - 1/2)$ is not always an integer, even if a is finite, the higher derivatives of H diverge in general. Therefore, this case corresponds to a Type IV singularity.

- In the case that $\alpha < 0$, when $t \rightarrow t_s$, we find $\rho \rightarrow 0$, $a \rightarrow a_0$, but $|p| \rightarrow \infty$. Therefore, this case corresponds to a Type II singularity.

Thus, it is explicitly demonstrated that simple fluid dark energy may bring the universe evolution to one of four possible future singularities as well as to an asymptotically non-singular universe. The particular future behavior is defined by the choice of the equation of state and of the parameters of this equation. Precisely the same phenomenon occurs for any field model of dark energy or modified gravity, so there is no qualitative difference between different dark energy models in this respect. The emergence of a finite-time future singularity in different modified gravities will be described in the fourth chapter.

In summary, this chapter provided a review of several popular models of modified gravity is given. Of course, the explicit choice of the models is related to our scientific interests. In particular, the following models were discussed: $F(R)$ gravity, modified Gauss-Bonnet theory, scalar-Einstein-Gauss-Bonnet gravity, non-local gravity and non-minimally coupled gravity. More exotic theories with explicit Lorentz symmetry breaking like Hořava-Lifshitz $F(R)$ gravity or with apparent Lorentz symmetry breaking like covariant power-counting renormalizable field gravity were also presented. In all cases, the spatially flat FRW cosmology was investigated. The qualitative possibility of a unified description of early-time inflation with late-time acceleration was also demonstrated. It turns out that the easiest and very realistic unification of these two eras is possible in $F(R)$ gravity.

III. COSMOLOGICAL RECONSTRUCTION OF MODIFIED GRAVITY

To explain the unified cosmic history of the universe, especially the inflation and/or the late-time accelerating expansion of the present universe, several models of extended or modified gravity were proposed in the previous chapter. Usually, we start from a theory, which is defined by the action, and solve equations of motion to define the background dynamics. In this chapter, however, we consider the inverse problem, i.e., the cosmological reconstruction of gravitational theories. In other words, using the fact that modified gravity is defined in terms of some arbitrary function (or with the help of some potential), we show how the complicated background cosmology, which complies with observational data, may be reconstructed. Eventually, this leads to a particular choice of the above arbitrary function (or potential). This provides the qualitative difference between general relativity and modified gravity; the latter theory (more exactly, some class of it) may have as a solution any given gravitational space-time. The problem is, of course, how to define the specific version of the theory that has this particular space-time as an explicit (exact or approximate) solution. The reconstruction scheme is explicitly constructed here for several modified gravities introduced in the previous chapter. The general approach to reconstruction in modified gravity and dark energy models was developed in refs. [18, 120, 121]. This kind of reconstruction may also be used for a spherically symmetric solution like black holes (see, for example, [122]).

A. Scalar-tensor gravity

As the first example, let us consider the reconstruction of scalar-Einstein gravity (or scalar-tensor theory), whose action can be written as

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} R - \frac{1}{2} \omega(\phi) \partial_\mu \phi \partial^\mu \phi - V(\phi) + L_{\text{matter}} \right\}. \quad (\text{III.1})$$

Here, $\omega(\phi)$ and $V(\phi)$ are functions of the scalar ϕ . The function $\omega(\phi)$ is not relevant and can be absorbed into the redefinition of the scalar field ϕ . In fact, if one redefines the scalar field ϕ by

$$\varphi \equiv \int^\phi d\phi \sqrt{|\omega(\phi)|}, \quad (\text{III.2})$$

the kinetic term of the scalar field in the action (III.1) has the following form:

$$-\omega(\phi) \partial_\mu \phi \partial^\mu \phi = \begin{cases} -\partial_\mu \varphi \partial^\mu \varphi & \text{when } \omega(\phi) > 0 \\ \partial_\mu \varphi \partial^\mu \varphi & \text{when } \omega(\phi) < 0 \end{cases}. \quad (\text{III.3})$$

The case of $\omega(\phi) > 0$ corresponds to the quintessence or non-phantom scalar field, but the case of $\omega(\phi) < 0$ corresponds to the phantom scalar. Although $\omega(\phi)$ can be absorbed into the redefinition of the scalar field, we keep $\omega(\phi)$ for a later convenience.

1. The reconstruction using the cosmological time

This section is based on the studies [123–125].

For the action (III.1), in the FRW equations (II.3), the energy density and the pressure are as follows:

$$\rho = \frac{1}{2} \omega(\phi) \dot{\phi}^2 + V(\phi), \quad p = \frac{1}{2} \omega(\phi) \dot{\phi}^2 - V(\phi). \quad (\text{III.4})$$

Then

$$\omega(\phi) \dot{\phi}^2 = -\frac{2}{\kappa^2} \dot{H}, \quad V(\phi) = \frac{1}{\kappa^2} (3H^2 + \dot{H}). \quad (\text{III.5})$$

Assuming $\omega(\phi)$ and $V(\phi)$ are given by a single function $f(\phi)$, as follows,

$$\omega(\phi) = -\frac{2}{\kappa^2} f'(\phi), \quad V(\phi) = \frac{1}{\kappa^2} (3f(\phi)^2 + f'(\phi)), \quad (\text{III.6})$$

the exact solution of the FRW equations (II.3) with (III.4) (when we neglect the contribution from the matter) has the following form:

$$\phi = t, \quad H = f(t). \quad (\text{III.7})$$

It can be confirmed that the equation given by the variation over ϕ

$$0 = \omega(\phi)\ddot{\phi} + \frac{1}{2}\omega'(\phi)\dot{\phi}^2 + 3H\omega(\phi)\dot{\phi} + V'(\phi), \quad (\text{III.8})$$

is also satisfied by the solution (III.7). Then, the arbitrary universe evolution expressed by $H = f(t)$ can be realized by an appropriate choice of $\omega(\phi)$ and $V(\phi)$. In other words, by defining the particular type of universe evolution, the corresponding scalar-Einstein gravity may be found.

As mentioned in (III.2) and (III.3), $\omega(\phi)$ can be absorbed into the redefinition of the scalar field ϕ . By keeping $\omega(\phi)$, however, we can construct a model that exhibits a smooth transition from the non-phantom phase to the phantom phase (and vice versa).

Let the matter energy density ρ_{matter} and the matter pressure p_{matter} be given by the sum of the contributions from the matter whose EoS parameters are $w_{\text{matter}}^i = p_{\text{matter}}^i/\rho_{\text{matter}}^i$ is a constants. Here, the suffix “ i ” specifies the type of matter. Using the conservation law of the energy density (II.144), we find $\rho_{\text{matter}} = \sum \rho_0^i a^{-3(1+w_{\text{matter}}^i)}$. Here, ρ_0^i is a constant. If $\omega(\phi)$ and $V(\phi)$ are given in terms of a single function $g(\phi)$ as

$$\begin{aligned} \omega(\phi) &= -\frac{2}{\kappa^2}g''(\phi) - \sum_i \frac{w_{\text{matter}}^i + 1}{2} \rho_0^i a_0^{-3(1+w_{\text{matter}}^i)} e^{-3(1+w_{\text{matter}}^i)g(\phi)}, \\ V(\phi) &= \frac{1}{\kappa^2} (3g'(\phi)^2 + g''(\phi)) + \sum_i \frac{w_{\text{matter}}^i - 1}{2} \rho_0^i a_0^{-3(1+w_{\text{matter}}^i)} e^{-3(1+w_{\text{matter}}^i)g(\phi)}, \end{aligned} \quad (\text{III.9})$$

the solution of the FRW equations (II.3) is

$$\phi = t, \quad H = g'(t), \quad \left(a = a_0 e^{g(t)}\right). \quad (\text{III.10})$$

Thus, even in the presence of matter, any given cosmology defined by $H = g'(t)$ can be realized by potentials (III.9).

As an example, the model with only one kind of matter with the EoS parameter w_{matter} (III.9) can be considered. It is also assumed that $w_{\text{matter}} > -1$. Choosing $g(\phi)$ as

$$g(\phi) = \frac{2}{3(w_{\text{matter}} + 1)} \ln \left(\frac{\phi}{t_s - \phi} \right), \quad (\text{III.11})$$

one obtains

$$\begin{aligned} \omega(\phi) &= -\frac{4}{3(w_{\text{matter}} + 1)\kappa^2} \left(-\frac{1}{\phi^2} + \frac{1}{(t_s - \phi)^2} \right) - \frac{w_{\text{matter}} + 1}{2} \rho_0 a_0^{-3(1+w_{\text{matter}})} \frac{(t_s - \phi)^2}{\phi^2}, \\ V(\phi) &= \frac{1}{\kappa^2} \left\{ \frac{4}{3(w_{\text{matter}} + 1)^2} \left(\frac{1}{\phi} + \frac{1}{t_s - \phi} \right)^2 + \frac{2}{3(w_{\text{matter}} + 1)} \left(-\frac{1}{\phi^2} + \frac{1}{(t_s - \phi)^2} \right) \right\} \\ &\quad + \frac{w_{\text{matter}} - 1}{2} \rho_0 a_0^{-3(1+w_{\text{matter}})} \frac{(t_s - \phi)^2}{\phi^2}. \end{aligned} \quad (\text{III.12})$$

The Hubble rate H follows as

$$H = \frac{2}{3(w_{\text{matter}} + 1)} \left(\frac{1}{t} + \frac{1}{t_s - t} \right). \quad (\text{III.13})$$

Because

$$\dot{H} = \frac{2}{3(w_{\text{matter}} + 1)} \left(-\frac{1}{t^2} + \frac{1}{(t_s - t)^2} \right), \quad (\text{III.14})$$

the EoS parameter w_{eff} (II.4) goes to $w_{\text{matter}} > -1$ when $t \rightarrow 0$ and goes to $-2 - w_{\text{matter}} < -1$ at late times. The crossing $w_{\text{eff}} = -1$ occurs when $\dot{H} = 0$, that is, $t = t_s/2$. Note that

$$\frac{\ddot{a}}{a} = \frac{16t_s}{27(w_{\text{matter}} + 1)^3(t_s - t)^2 t^2} \left\{ t - \frac{(3w_{\text{matter}} + 1)t_s}{4} \right\}. \quad (\text{III.15})$$

Thus, if $w_{\text{matter}} > -1/3$, the deceleration of the universe turns to the acceleration at $t = t_a \equiv (3w_{\text{matter}} + 1)t_s/4$.

The second example in (III.9) is given by

$$g(\phi) = -\alpha \ln \left(1 - \beta \ln \frac{\phi}{\kappa} \right). \quad (\text{III.16})$$

Here, α and β are dimensionless positive constants. Let us choose $\beta \sim \mathcal{O}(10^{-2})$ for later convenience. Note that the parameter order of 10^{-2} is not so unnatural. When only one kind of matter with the EoS parameter w_{matter} is included, for simplicity, Eq. (III.16) gives the following expressions for $\omega(\phi)$ and $V(\phi)$:

$$\begin{aligned} \omega(\phi) &= -\frac{2\alpha\beta \left(\beta - 1 + \beta \ln \frac{\phi}{\kappa} \right)}{\kappa^2 \left(1 - \beta \ln \frac{\phi}{\kappa} \right)^2 \phi^2} - \frac{w_{\text{matter}} + 1}{2} \rho_0 a_0^{-3(1+w_{\text{matter}})} \left(1 - \beta \ln \frac{\phi}{\kappa} \right)^{3(w_{\text{matter}}+1)\alpha}, \\ V(\phi) &= \frac{\alpha\beta \left(3\alpha\beta + \beta - 1 + \beta \ln \frac{\phi}{\kappa} \right)}{\kappa^2 \left(1 - \beta \ln \frac{\phi}{\kappa} \right)^2 \phi^2} + \frac{w_{\text{matter}} - 1}{2} \rho_0 a_0^{-3(1+w_{\text{matter}})} \left(1 - \beta \ln \frac{\phi}{\kappa} \right)^{3(w_{\text{matter}}+1)\alpha}. \end{aligned} \quad (\text{III.17})$$

Supposing $\rho_0 a_0^{-3(1+w_{\text{matter}})} \sim \mathcal{O}(\kappa^{-2})$, any small parameter, compared with the Planck scale does not appear. Now, the Hubble rate is given by $H = \alpha\beta / (1 - \beta \ln(t/\kappa)) t$, which is positive if

$$0 < t < t_s \equiv \kappa e^{\frac{1}{\beta}}, \quad (\text{III.18})$$

and has a Big Rip-type singularity at $t = t_s$. Because $10^{61} \sim e^{140}$, with the choice $\beta \sim 1/140$, we obtain $t_s \sim \kappa \times 10^{61} \sim (10^{-33} \text{ eV})^{-1}$, whose order is that of the age of the present universe. Thus, due to the property of the exponential function (or logarithmic function), the small scale like t_s appears rather naturally. We should also note that if $\alpha\beta \sim \mathcal{O}(10^{0-2})$ and t is a present age of the universe $t \sim (10^{-33} \text{ eV})^{-1}$, the observed value of the Hubble rate $H \sim 10^{-33} \text{ eV}$ can also be reproduced. Because

$$\frac{\ddot{a}}{a} = \frac{\alpha\beta^2 \left(\ln \frac{t}{\kappa} + \alpha + 1 - \frac{1}{\beta} \right)}{\left(1 - \beta \ln \frac{t}{\kappa} \right)^2 t^2}, \quad (\text{III.19})$$

the universe changes to an accelerating from a decelerating universe when

$$t = t_a \equiv \kappa e^{\frac{1}{\beta} - \alpha - 1} < t_s. \quad (\text{III.20})$$

Because the energy density ρ of the scalar field ϕ and that of the matter ρ_{matter} are given by

$$\begin{aligned} \rho &= \frac{3\alpha^2\beta^2}{\kappa^2 \left(1 - \beta \ln \frac{t}{\kappa} \right)^2 t^2} - \rho_0 a_0^{-3(1+w_{\text{matter}})} \left(1 - \beta \ln \frac{t}{\kappa} \right)^{3(w_{\text{matter}}+1)\alpha}, \\ \rho_{\text{matter}} &= \rho_0 a_0^{-3(1+w_{\text{matter}})} \left(1 - \beta \ln \frac{t}{\kappa} \right)^{3(w_{\text{matter}}+1)\alpha}, \end{aligned} \quad (\text{III.21})$$

the coincidence time t_c could be given by solving the following equation:

$$\left(1 - \beta \ln \frac{t_c}{\kappa} \right)^{3(w_{\text{matter}}+1)\alpha+2} t_c^2 = \frac{3\alpha^2\beta^2}{\kappa^2 \rho_0 a_0^{-3(1+w_{\text{matter}})}}. \quad (\text{III.22})$$

One may regard ρ_{matter} as the sum of the energy density of usual matter, like baryons, and that of (cold) dark matter. If ρ corresponds to the energy density of the dark energy, the current data indicate that $\rho : \rho_{\text{matter}} \sim 7 : 3$. Then, in the present universe, it follows that

$$\left(1 - \beta \ln \frac{t}{\kappa} \right)^{3(w_{\text{matter}}+1)\alpha+2} t^2 \sim \frac{9\alpha^2\beta^2}{10\kappa^2 g_0}. \quad (\text{III.23})$$

Thus, in the model (III.16), the acceleration of the present universe and the coincidence problem seem to be explained rather naturally. This is achieved via explicit cosmological reconstruction of the scalar potentials.

2. The reconstruction of the scalar-tensor models using e-folding

In this subsection, the formulation using the e-folding N instead of cosmological time is considered as in Eqs. (II.337) and (II.338). The FRW equations (II.3) with (III.4) (when the matter contribution is neglected) are given in terms of N , as follows:

$$\frac{3}{\kappa^2}H^2 = \frac{H^2\omega(\phi)\phi'^2}{2} + V(\phi), \quad -\frac{1}{\kappa^2}(2HH' + 3H^2) = \frac{H^2\omega(\phi)\phi'^2}{2} - V(\phi). \quad (\text{III.24})$$

We now identify $\phi = N$; then,

$$\omega(\phi) = -\frac{2H'}{\kappa^2 H}, \quad V(\phi) = \frac{1}{\kappa^2}(HH' + 3H^2). \quad (\text{III.25})$$

The above equations (III.25) show that if one studies the model

$$\omega(\phi) = -\frac{2\tilde{f}'(\phi)}{\kappa^2 \tilde{f}(\phi)}, \quad V(\phi) = \frac{1}{\kappa^2}(\tilde{f}(\phi)\tilde{f}'(\phi) + 3\tilde{f}(\phi)^2), \quad (\text{III.26})$$

which is given in terms of a single function $\tilde{f}$, then the exact solution is given by $H(N) = \tilde{f}(N)$.

Let us consider an example by using the following function $\tilde{f}$:

$$\tilde{f}(\phi) = H_1\phi^{-\gamma}. \quad (\text{III.27})$$

Here, H_1 and γ are positive constants. In this case

$$\omega(\phi) = \frac{2\gamma}{\kappa^2\phi}, \quad V(\phi) = \frac{H_1^2}{\kappa^2}(-\gamma\phi^{-2\gamma-1} + 3\phi^{-2\gamma}), \quad (\text{III.28})$$

and the exact solution is given by $H = H_1N^{-\gamma}$. With a redefinition of ϕ by $\phi \rightarrow \varphi = \frac{2\sqrt{2\gamma}\phi}{\kappa}$, the kinetic term becomes canonical, and the potential is given by

$$V(\varphi) = \frac{H_1^2}{\kappa^2} \left\{ -\gamma \left(\frac{\kappa^2}{8\gamma} \right)^{-2\gamma-1} \varphi^{-4\gamma-2} + 3 \left(\frac{\kappa^2}{8\gamma} \right)^{-2\gamma} \varphi^{-4\gamma} \right\}. \quad (\text{III.29})$$

At late times (large N), only the last term is relevant. Conversely, at late times, if the potential is given by $V \propto \varphi^{-4\gamma}$, the solution has the form of $H \propto N^{-\gamma}$. The effective EoS parameter (II.4) is equal to

$$w_{\text{eff}} = -1 + \frac{2\gamma}{3N}, \quad (\text{III.30})$$

which goes to -1 when N goes to infinity.

Because we naively find $C \equiv \frac{\text{Plank scale}}{H_0} \sim 10^{61} \sim e^{140}$, if $\frac{H_1}{H_0} \sim C$, it follows that $N \sim \ln C$ and, therefore, $\gamma \sim 28$. Thus, the fine-tuning problem might be relaxed, but in this case one obtains $w_{\text{eff}} \sim -0.87$, which is a little bigger than the observational value $-0.14 < 1 + w < 0.12$ [126].

In the case that $\tilde{C} \equiv \frac{\text{Weak scale}}{H_0} \sim 10^{44} \sim e^{101}$, if $\frac{H_1}{H_0} \sim \tilde{C}$, it follows that $N \sim \ln \tilde{C}$ and $\gamma \sim 21$. This finishes the cosmological reconstruction of scalar-tensor theory using e-folding.

3. Two-scalar model

Note that when studying the transition from the non-phantom phase to the phantom phase, the instability of the single scalar-tensor gravity becomes infinite at the transition point [123, 124]. Thus, we only investigate two scalar-tensor theory, which has an instability that can always finite. Moreover, with two scalars it is easier to realize unified universe history expansion.

The action of two scalar-tensor gravity is

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} R - \frac{1}{2} \omega(\phi) \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} \eta(\chi) \partial_\mu \chi \partial^\mu \chi - V(\phi, \chi) \right\}. \quad (\text{III.31})$$

Here, $\omega(\phi)$ is a function of the scalar field ϕ , and $\eta(\chi)$ is a function of another scalar field χ . Note that the above theory parameterizes a large number of popular dark energy models. Indeed, if both scalars could be canonical scalars, then a kind of the quintessence model emerges. One of scalars may be phantom (wrong sign of the kinetic term) while another scalar remains the canonical one. This is dark energy sometimes called quintom theory. Finally, both scalars could be phantom. Moreover, two scalars may be organized so that a charged scalar with real and imaginary parts appears.

It is also assumed that the FRW metric is spatially flat (II.6) and that ϕ and χ only depend on the time coordinate t . The FRW equations lead to

$$\omega(\phi)\dot{\phi}^2 + \eta(\chi)\dot{\chi}^2 = -\frac{2}{\kappa^2}\dot{H}, \quad V(\phi, \chi) = \frac{1}{\kappa^2} (3H^2 + \dot{H}). \quad (\text{III.32})$$

If

$$\omega(t) + \eta(t) = -\frac{2}{\kappa^2}f'(t), \quad V(t, t) = \frac{1}{\kappa^2} (3f(t)^2 + f'(t)), \quad (\text{III.33})$$

the explicit solution follows as:

$$\phi = \chi = t, \quad H = f(t). \quad (\text{III.34})$$

One may choose ω to always be positive and η to always be negative, for example

$$\omega(\phi) = -\frac{2}{\kappa^2} \left\{ f'(\phi) - \sqrt{\alpha^2 + f'(\phi)^2} \right\} > 0, \quad \eta(\chi) = -\frac{2}{\kappa^2} \sqrt{\alpha^2 + f'(\chi)^2} < 0. \quad (\text{III.35})$$

Here, α is a constant. Define a new function $\tilde{f}(\phi, \chi)$ by

$$\tilde{f}(\phi, \chi) \equiv -\frac{\kappa^2}{2} \left(\int d\phi \omega(\phi) + \int d\chi \eta(\chi) \right). \quad (\text{III.36})$$

The constant of the integration could be fixed to require

$$\tilde{f}(t, t) = f(t). \quad (\text{III.37})$$

If $V(\phi, \chi)$ is given using $\tilde{f}(\phi, \chi)$ as

$$V(\phi, \chi) = \frac{1}{\kappa^2} \left(3\tilde{f}(\phi, \chi)^2 + \frac{\partial \tilde{f}(\phi, \chi)}{\partial \phi} + \frac{\partial \tilde{f}(\phi, \chi)}{\partial \chi} \right), \quad (\text{III.38})$$

the FRW and the scalar field equations are also satisfied:

$$0 = \omega(\phi)\ddot{\phi} + \frac{1}{2}\omega'(\phi)\dot{\phi}^2 + 3H\omega(\phi)\dot{\phi} + \frac{\partial \tilde{V}(\phi, \chi)}{\partial \phi}, \quad 0 = \eta(\chi)\ddot{\chi} + \frac{1}{2}\eta'(\chi)\dot{\chi}^2 + 3H\eta(\chi)\dot{\chi} + \frac{\partial \tilde{V}(\phi, \chi)}{\partial \chi}. \quad (\text{III.39})$$

Thus, for given cosmology the cosmological reconstruction (the calculation of scalar potentials) is made.

It is now interesting to investigate the (in)stability of the model under discussion. By introducing the new quantities, X_ϕ , X_χ , and Y as

$$X_\phi \equiv \dot{\phi}, \quad X_\chi \equiv \dot{\chi}, \quad Y \equiv \frac{\tilde{f}(\phi, \chi)}{H}, \quad (\text{III.40})$$

the FRW equations and the scalar field equations (III.39) are:

$$\begin{aligned} \frac{dX_\phi}{dN} &= -\frac{\omega'(\phi)}{2H\omega(\phi)} (X_\phi^2 - 1) - 3(X_\phi - Y), & \frac{dX_\chi}{dN} &= -\frac{\eta'(\chi)}{2H\eta(\chi)} (X_\chi^2 - 1) - 3(X_\chi - Y), \\ \frac{dY}{dN} &= \frac{1}{2\kappa^2 H^2} \{X_\phi (X_\phi Y - 1) + X_\chi (X_\chi Y - 1)\}. \end{aligned} \quad (\text{III.41})$$

Here, $d/dN \equiv H^{-1}d/dt$. In the solution (III.34), $X_\phi = X_\chi = Y = 1$. The following perturbation may be considered

$$X_\phi = 1 + \delta X_\phi, \quad X_\chi = 1 + \delta X_\chi, \quad Y = 1 + \delta Y. \quad (\text{III.42})$$

Thus,

$$\frac{d}{dN} \begin{pmatrix} \delta X_\phi \\ \delta X_\chi \\ \delta Y \end{pmatrix} = M \begin{pmatrix} \delta X_\phi \\ \delta X_\chi \\ \delta Y \end{pmatrix}, \quad M \equiv \begin{pmatrix} -\frac{\omega'(\phi)}{H\omega(\phi)} - 3 & 0 & 3 \\ 0 & -\frac{\eta'(\chi)}{H\eta(\chi)} - 3 & 3 \\ \frac{1}{2\kappa^2 H^2} & \frac{1}{2\kappa^2 H^2} & \frac{1}{\kappa^2 H^2} \end{pmatrix}. \quad (\text{III.43})$$

The eigenvalues of the matrix M are given by solving the following eigenvalue equation

$$0 = \left(\lambda + \frac{\omega'(\phi)}{H\omega(\phi)} + 3 \right) \left(\lambda + \frac{\eta'(\chi)}{H\eta(\chi)} + 3 \right) \left(\lambda - \frac{1}{\kappa^2 H^2} \right) + \frac{3}{2\kappa^2 H^2} \left(\lambda + \frac{\omega'(\phi)}{H\omega(\phi)} + 3 \right) + \frac{3}{2\kappa^2 H^2} \left(\lambda + \frac{\eta'(\chi)}{H\eta(\chi)} + 3 \right). \quad (\text{III.44})$$

The eigenvalues (III.44) for the two-scalar model are clearly finite. Thus, the instability could be finite. In fact, right at the transition point where $\dot{H} = f'(t) = 0$ and, therefore, $f'(\phi) = f'(\chi) = 0$, for the choice in (III.35), one gets

$$\omega(\phi) = -\eta(\chi) = \frac{2\alpha}{\kappa^2}, \quad \omega'(\phi) = -\frac{2\ddot{H}}{\kappa^2}, \quad \eta'(\chi) = 0. \quad (\text{III.45})$$

Then, the eigenvalue equation (III.44) reduces to

$$0 = \lambda^3 + (-A - B + 6)\lambda^2 + (AB - 3A - 3B + 9)\lambda - \frac{3}{2}AB + 9B, \quad A \equiv \frac{\ddot{H}}{\alpha}, \quad B \equiv \frac{1}{\kappa^2 H^2}. \quad (\text{III.46})$$

Here, $\alpha > 0$ is chosen. The eigenvalues are surely finite, which shows that even if the solution (III.34) is not stable, the solution has a non-vanishing measure and, therefore, the transition from the non-phantom phase to the phantom phase can occur. We should also note that the solution (III.34) can be stable. For example, for the case $A, B \rightarrow 0$. Eq. (III.46) further reduces to

$$0 = \lambda(\lambda + 3)^2. \quad (\text{III.47})$$

Then, the eigenvalues are given by 0 and -3 . Because there is no positive eigenvalue, the solution (III.34) is stable in this case.

It is not difficult to extend the above formulation to the multi-scalar model, whose action is given by

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} R - \frac{1}{2} \sum_i \omega_i(\phi_i) \partial_\mu \phi_i \partial^\mu \phi_i - V(\phi_i) \right\}. \quad (\text{III.48})$$

Here, $\omega_i(\phi_i)$ is a function of the scalar field ϕ_i . We now choose ω_i to satisfy

$$\sum_i \omega_i(t) = -\frac{2}{\kappa^2} f'(t), \quad (\text{III.49})$$

by a proper function. The potential $V(\phi_i)$ is chosen as

$$V(\phi_i) = \frac{1}{\kappa^2} \left(3\tilde{f}(\phi_i) + \sum_i \frac{\partial \tilde{f}}{\partial \phi_i} \right). \quad (\text{III.50})$$

Here

$$\tilde{f}(\phi_i) \equiv -\frac{\kappa^2}{2} \sum_i \int d\phi_i \omega_i(\phi_i). \quad (\text{III.51})$$

The constant of the integration in (III.51) is determined to satisfy

$$\tilde{f}(\phi_i) \Big|_{\text{all } \phi_i = t} = f(t). \quad (\text{III.52})$$

Then, a solution of the FRW equations and the scalar field equations is given by

$$\phi_i = t, \quad H = f(t). \quad (\text{III.53})$$

The remaining arguments coincide with those given for the two-scalar model.

It is interesting to demonstrate that the transition from the matter-dominant period to the acceleration period can be realized in the present formulation. In the following, the matter contribution can be neglected because the ratio of the matter with the (effective) dark matter may be small.

The first example is as follows:

$$H = f(t) = g'(t) = g_0 + \frac{g_1}{t}. \quad (\text{III.54})$$

When t is large, the first term in (III.54) dominates and the Hubble rate H becomes a constant. Therefore, the universe is asymptotically the de Sitter space, which corresponds to an accelerating universe. On the other hand, when t is small, the second term in (III.54) dominates and the scale factor behaves as $a \sim t^{g_1}$. Therefore, if $g_1 = 2/3$, the matter-dominated period emerges. Here, one of the scalars may be considered as usual matter.

Because

$$f'(t) = -\frac{g_1}{t} < 0, \quad (\text{III.55})$$

instead of (III.35), by using a positive constant α , one may choose

$$\omega(\phi) = -\frac{2(1+\alpha)}{\kappa^2} f'(\phi) > 0, \quad \eta(\chi) = \frac{2\alpha}{\kappa^2} f'(\phi) < 0, \quad (\text{III.56})$$

that is,

$$\omega(\phi) = \frac{2(1+\alpha)}{\kappa^2} \frac{g_1}{\phi^2}, \quad \eta(\chi) = \frac{2\alpha}{\kappa^2} \frac{g_1}{\phi^2}. \quad (\text{III.57})$$

One should note that in the limit $\alpha \rightarrow 0$, the scalar field χ decouples and the single-scalar model follows. Thus, one gets

$$\tilde{f}(\phi, \chi) = g_0 + \frac{(1+\alpha)g_1}{\phi} - \frac{\alpha g_1}{\chi}, \quad V(\phi, \chi) = \frac{1}{\kappa^2} \left\{ 3 \left(g_0 + \frac{(1+\alpha)g_1}{\phi} - \frac{\alpha g_1}{\chi} \right)^2 - \frac{(1+\alpha)g_1}{\phi^2} + \frac{\alpha g_1}{\chi^2} \right\}. \quad (\text{III.58})$$

Thus, for the scalar-tensor theory with such potentials the matter-dominant era occurs before the acceleration epoch. This fact is easily established within the reconstruction scheme.

Before going to the second example, we consider the Einstein gravity with cosmological constant and with matter characterized by the EOS parameter w . The FRW equation is as follows:

$$\frac{3}{\kappa^2} H^2 = \rho_0 a^{-3(1+w)} + \frac{3}{\kappa^2 l^2}. \quad (\text{III.59})$$

Here, l^2 is the inverse cosmological constant. The solution of (III.59) is given by

$$a = a_0 e^{g(t)}, \quad g(t) = \frac{2}{3(1+w)} \ln \left(\alpha \sinh \left(\frac{3(1+w)}{2l} (t - t_s) \right) \right). \quad (\text{III.60})$$

Here, t_s is a constant of the integration and

$$\alpha^2 \equiv \frac{1}{3} \kappa^2 l^2 \rho_0 a_0^{-3(1+w)}. \quad (\text{III.61})$$

Thus, this is the Λ CDM era as it follows in the commonly accepted approach.

As the second example, the reconstruction of the Λ CDM metric (III.60) for scalar-tensor theory is done below. Because

$$f(t) \equiv g'(t) = \frac{1}{l} \coth \left(\frac{3(1+w)}{2l} (t - t_s) \right), \quad f'(t) = g''(t) = -\frac{3(1+w)}{2l^2} \sinh^{-2} \left(\frac{3(1+w)}{2l} (t - t_s) \right) < 0, \quad (\text{III.62})$$

it is convenient to use (III.56) instead of (III.35). Then, one arrives at

$$\begin{aligned}
\omega(\phi) &= \frac{3(1+w)(1+\alpha)}{\kappa^2 l^2} \sinh^{-2} \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) > 0, \\
\eta(\chi) &= -\frac{3(1+w)\alpha}{\kappa^2 l^2} \sinh^{-2} \left(\frac{3(1+w)}{2l} (\chi - t_s) \right) < 0, \\
\tilde{f}(\phi, \chi) &= \frac{1+\alpha}{l} \coth \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) - \frac{\alpha}{l} \coth \left(\frac{3(1+w)}{2l} (\chi - t_s) \right), \\
V(\phi, \chi) &= \frac{1}{\kappa^2} \left[\frac{3}{l^2} \left\{ (1+\alpha) \coth \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) - \alpha \coth \left(\frac{3(1+w)}{2l} (\chi - t_s) \right) \right\}^2 \right. \\
&\quad \left. - \frac{3(1+w)(1+\alpha)}{l^2} \sinh^{-2} \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) + \frac{3(1+w)\alpha}{l^2} \sinh^{-2} \left(\frac{3(1+w)}{2l} (\chi - t_s) \right) \right]. \tag{III.63}
\end{aligned}$$

This completes the reconstruction of scalar potentials to reproduce the Λ CDM epoch.

Thus, in both examples, (III.54) and (III.60), the matter dominant stage, the transition from the matter dominant phase to the acceleration phase and acceleration epoch occur. In the acceleration phase of the above examples, the universe asymptotically approaches the de Sitter space. This does not conflict with the Wilkinson Microwave Anisotropy Probe (WMAP) data. Indeed, three years WMAP data have been analyzed in ref. [127–129]. The combined analysis of WMAP data with the Supernova Legacy Survey (SNLS) constrains the dark energy equation of state w_{DE} , pushing it toward the cosmological constant. The marginalized best fit values of the equation of state parameter at the 68% confidence level are given by $-1.14 \leq w_{\text{DE}} \leq -0.93$. In the prior case of a flat universe, the combined data give $-1.06 \leq w_{\text{DE}} \leq -0.90$. In the examples (III.54) and (III.60), the universe goes asymptotically to the de Sitter space, which gives $w_{\text{DE}} \rightarrow -1$, which does not conflict with the above constraints. Note, however, that one needs to fine-tune g_0 in (III.54) and $1/l$ in (III.60) to be $g_0 \sim 1/l \sim 10^{-33}$ eV, to reproduce the observed Hubble rate $H_0 \sim 70 \text{ km s}^{-1} \text{ Mpc}^{-1} \sim 10^{-33}$ eV.

As a final remark, there is no problem regarding the inclusion of the usual matter (for example, ideal fluid and dust) into the above reconstruction scenario and/or finding the scalar potentials corresponding to realistic cosmology in the multi-scalar case. Moreover, in the same way, one may include the radiation-dominated epoch where quantum effects may still be neglected in the above scenario. In the next section it is shown how similar reconstruction scheme may be developed for modified gravity.

4. Brans-Dicke gravity

The action of the original Brans-Dicke model [130] has the following form:

$$S_{\text{BD}} = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \left(\phi R - \omega \frac{\partial_\mu \phi \partial^\mu \phi}{\phi} \right), \tag{III.64}$$

which can be generalized as

$$S = \int d^4x \sqrt{-g} \left[\frac{e^{\varphi(\phi)} R}{2\kappa^2} - \frac{1}{2} \omega(\phi) \partial_\mu \phi \partial^\mu \phi - V(\phi) + \mathcal{L}_m \right]. \tag{III.65}$$

This kind of model was applied to the dark energy problem in [131], and it was found that even if $\omega(\phi) > 0$, that is, the scalar field is canonical, the phantom universe can be realized.

The FRW equations are as follows:

$$H^2 = \frac{e^{-\varphi} \kappa^2}{3} \left[\frac{1}{2} \omega(\phi) \dot{\phi}^2 + V(\phi) \right], \quad \dot{H} = -\frac{e^{-\varphi} \kappa^2}{2} \omega(\phi) \dot{\phi}^2 - \frac{1}{2} (\ddot{\varphi} + \dot{\varphi}^2), \tag{III.66}$$

This can be rewritten as

$$\omega(\phi) \dot{\phi}^2 = -\frac{2e^{\varphi}}{\kappa^2} \left[\dot{H} + \frac{1}{2} (\ddot{\varphi} + \dot{\varphi}^2) \right], \quad V(\phi) = \frac{2e^{\varphi}}{\kappa^2} \left[\dot{H} + 3H^2 + \frac{1}{2} (\ddot{\varphi} + \dot{\varphi}^2) \right]. \tag{III.67}$$

Then, if the following choice of scalar potentials using a function $f(\phi)$ [132] is taken:

$$\omega(\phi) = -\frac{2e^{\varphi(\phi)}}{\kappa^2} \left[f'(\phi) + \frac{1}{2} (\varphi''(\phi) + \varphi'(\phi)^2) \right], \quad V(\phi) = \frac{2e^{\varphi(\phi)}}{\kappa^2} \left[f'(\phi) + 3f(\phi)^2 + \frac{1}{2} (\varphi''(\phi) + \varphi'(\phi)^2) \right], \tag{III.68}$$

the explicit solution looks like

$$H = f(t), \quad \phi = t. \quad (\text{III.69})$$

Note that $\varphi(\phi)$ can be an arbitrary function of ϕ . By this reconstruction formalism, the phantom crossing ($w = -1$ crossing) can be realized for $\omega(\phi) > 0$, that is, when the scalar field is canonical (non-ghost). A slightly different version of the reconstruction formalism for above theory is given in [133].

B. The k -essence model

Here, based on [134], we consider the reconstruction of the k -essence model, which represents the evident generalization of quintessence theory. The k -essence model is a rather general model that includes only one scalar field and the action is given by

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - K(\phi, X) + L_{\text{matter}} \right), \quad X \equiv \partial^\mu \phi \partial_\mu \phi. \quad (\text{III.70})$$

Here, ϕ is a scalar field.

Now, the Einstein equation has the following form:

$$\frac{1}{\kappa^2} \left(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) = -K(\phi, X) g_{\mu\nu} + 2K_X(\phi, X) \partial_\mu \phi \partial_\nu \phi + T_{\mu\nu}. \quad (\text{III.71})$$

Here, $K_X(\phi, X) \equiv \partial K(\phi, X) / \partial X$. On the other hand, the variation over ϕ gives

$$0 = -K_\phi(\phi, X) + 2\nabla^\mu (K(\phi, X) \partial_\mu \phi). \quad (\text{III.72})$$

Here, $K_\phi(\phi, X) \equiv \partial K(\phi, X) / \partial \phi$ and it is assumed that the scalar field ϕ does not directly couple with the matter.

1. The reconstruction using the cosmological time

Let us start from the FRW universe (II.6) and assume that the scalar field ϕ only depends on time. The FRW equations are given by

$$\frac{3}{\kappa^2} H^2 = 2X \frac{\partial K(\phi, X)}{\partial X} - K(\phi, X) + \rho_{\text{matter}}(t), \quad -\frac{1}{\kappa^2} (2\dot{H} + 3H^2) = K(\phi, X) + p_{\text{matter}}(t). \quad (\text{III.73})$$

The FRW equations (III.73) show that for the following model

$$K(\phi, X) = \sum_{n=0}^{\infty} (X+1)^n K^{(n)}(\phi), \quad K^{(0)}(\phi) = -\frac{1}{\kappa^2} (2f'(\phi) + 3f(\phi)^2), \quad K^{(1)}(\phi) = \frac{1}{\kappa^2} f'(\phi), \quad (\text{III.74})$$

there exists a solution given by

$$H = f(t), \quad \phi = t. \quad (\text{III.75})$$

Note that in (III.74), $K^{(n)}(\phi)$ $n \geq 2$ can be an arbitrary function. To investigate the role of $K^{(n)}(\phi)$ ($n \geq 2$), the following perturbation may be discussed:

$$H = f(t) + \delta H, \quad \phi = t + \delta \phi, \quad (\text{III.76})$$

The stability of the solution (III.75) should now be investigated. In (III.76), it is assumed that δH and $\delta \phi$ do not depend on the spatial coordinates. Then, δH is solved as a function of $\delta \phi$: $\delta H = \delta H(\delta \phi)$, and one obtains

$$\frac{d\delta \dot{\phi}}{dN} = \left[-3 - \frac{g''}{g'^2} - \frac{d}{dN} \left\{ \frac{\kappa^2}{6g'^2} \left(8K^{(2)} - \frac{2}{\kappa^2} g'' \right) \right\} \right] \delta \dot{\phi}. \quad (\text{III.77})$$

If the quantity inside [] is negative, the solution (III.75) becomes stable. Because Eq. (III.77) contains $K^{(2)}$, if $K^{(2)}$ is chosen properly, the solution corresponding to an arbitrary evolution law becomes stable.

Let us consider the condition that there is a Schwarzschild or Schwarzschild-(A)dS solution. Let ϕ be a constant:

$$\phi = \phi_0. \quad (\text{III.78})$$

Eqs. (III.71) and (III.72) reduce to

$$\frac{1}{\kappa^2} \left(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) = -K(\phi_0, 0) g_{\mu\nu} + T_{\mu\nu}, \quad (\text{III.79})$$

$$0 = K_\phi(\phi_0, 0). \quad (\text{III.80})$$

When $T_{\mu\nu} = 0$, if $K(\phi_0, 0)$ does not vanish, a solution of (III.79) is given by Schwarzschild-(A)dS space-time. On the other hand, if $K(\phi, 0)$ vanishes, the Schwarzschild solution, which is asymptotically Minkowski space-time, is a solution. The equation (III.80) requires that, in general, $K_\phi(\phi, 0)$ has an extremum or $K(\phi, 0)$ is a constant independent of ϕ . Especially if $K(\phi, 0)$ vanishes identically, Eq. (III.79) gives the Schwarzschild solution.

$K(\phi, X)$ can be written as

$$K(\phi, X) = \sum_{n=0} (1+X)^n K_n(\phi). \quad (\text{III.81})$$

Then, if $K(\phi, 0) = 0$, it follows that

$$\sum_{n=0} K_n(\phi) = 0. \quad (\text{III.82})$$

In particular, one may choose

$$K_3(\phi) = -K_0(\phi) - K_1(\phi) - K_2(\phi). \quad (\text{III.83})$$

Thus, if the condition (III.82) or (III.83) is satisfied, the Schwarzschild space-time is always a solution independent of the value of ϕ as long as ϕ is a constant. Here, any point source of matter creates the Schwarzschild space-time which generates the Newton potential. The correction to Newton's law does not appear. Note that the value of the scalar field ϕ changes by the evolution of the universe, but as long as the condition (III.82) or (III.83) is satisfied, in a local region where ϕ is almost constant, the correction to Newton's law is negligible.

2. The reconstruction using e-folding

Because it is often convenient to use the redshift z or e-folding N , we rewrite the FRW equations (III.73) as

$$\frac{3}{\kappa^2} H^2 = 2X \frac{\partial K(\phi, X)}{\partial X} - K(\phi, X) + \rho_{\text{matter}}(N), \quad -\frac{1}{\kappa^2} (2HH' + 3H^2) = K(\phi, X) + p_{\text{matter}}(N). \quad (\text{III.84})$$

Here, $H' \equiv dH/dN$. If the matter energy density is given by a sum of the contributions from the matter with constant EoS parameters w_i , one finds

$$\begin{aligned} \rho_{\text{matter}}(N) &= \sum_i \rho_{0i} a^{-3(1+w_i)} = \sum_i \rho_{0i} e^{-3(1+w_i)(N-N_0)}, \\ p_{\text{matter}}(N) &= \sum_i w_i \rho_{0i} a^{-3(1+w_i)} = \sum_i w_i \rho_{0i} e^{-3(1+w_i)(N-N_0)}. \end{aligned} \quad (\text{III.85})$$

Here, ρ_{0i} 's are constants. For general energy density $\rho_{\text{matter}}(N)$, because the conservation law (II.144) can be rewritten in terms of N as

$$\rho'_{\text{matter}}(N) + 3(\rho_{\text{matter}}(N) + p_{\text{matter}}(N)) = 0, \quad (\text{III.86})$$

it follows that

$$p_{\text{matter}}(N) = -\rho_{\text{matter}}(N) - \frac{1}{3} \rho'_{\text{matter}}(N). \quad (\text{III.87})$$

With the help of the FRW equations (III.84), we find

$$\begin{aligned} K(\phi, X) &= -\frac{1}{\kappa^2} \left(2H \frac{dH}{dN} + 3H^2 \right) + \rho_{\text{matter}}(N) + \frac{1}{3} \rho'_{\text{matter}}(N), \\ H^2 \frac{\partial K(\phi, X)}{\partial X} &= \frac{1}{\kappa^2} H \frac{dH}{dN} - \frac{1}{6} \rho'_{\text{matter}}(N). \end{aligned} \quad (\text{III.88})$$

If a new variable $G(N) = H(N)^2$ is defined, the equations (III.88) can be rewritten as

$$\begin{aligned} K(\phi, X) &= -\frac{1}{\kappa^2} (G'(N) + 3G(N)) + \rho_{\text{matter}}(N) + \frac{1}{3} \rho'_{\text{matter}}(N), \\ H^2 \frac{\partial K(\phi, X)}{\partial X} &= \frac{1}{2\kappa^2} G'(N) - \frac{1}{6} \rho'_{\text{matter}}(N). \end{aligned} \quad (\text{III.89})$$

Using the appropriate function $g_\phi(\phi)$, if we choose

$$\begin{aligned} K(\phi, X) &= \sum_{n=0}^{\infty} \left(\frac{X}{g_\phi(\phi) + \frac{\kappa^2}{3} \rho'_{\text{matter}}(\phi)} + 1 \right)^n \tilde{K}^{(n)}(\phi), \\ \tilde{K}^{(0)}(\phi) &\equiv -\frac{1}{\kappa^2} (g'_\phi(\phi) + 3g_\phi(\phi)) \quad \tilde{K}^{(1)}(\phi) = \frac{1}{2\kappa^2} g'_\phi(\phi), \end{aligned} \quad (\text{III.90})$$

the explicit solution for the FRW equations (III.73) is found as

$$G(N) = H(N)^2 = g_\phi(N) + \frac{\kappa^2}{3} \rho_{\text{matter}}(N), \quad \phi = N \quad (X = -H^2). \quad (\text{III.91})$$

Here, $\tilde{K}^{(n)}(\phi)$ with $n \geq 2$ can be arbitrary. The reconstruction is fulfilled.

Let us now investigate the (in)stability of the solution (III.91). For this purpose, the perturbation from the solution (III.91) is taken as follows:

$$G(N) = G_0(N) + \delta G(N) \quad \left(G_0(N) \equiv g_\phi(N) + \frac{\kappa^2}{3} \rho'_{\text{matter}}(N) \right), \quad \phi = N + \delta\phi. \quad (\text{III.92})$$

Note that N -dependence in the energy density ρ_{matter} is usually given by a fixed function as in (III.85). Therefore, $\delta\rho_{\text{matter}} = 0$. Thus, the equations (III.89) give

$$\begin{aligned} -\frac{1}{\kappa^2} (g''_\phi(N) + 3g'_\phi(N)) \delta\phi(N) - \frac{g'_\phi(N)}{2\kappa^2} \left(\frac{\delta G(N)}{G_0(N)} + 2\delta\phi'(N) - \frac{G'_0(N)}{G_0(N)} \delta\phi(N) \right) &= -\frac{1}{\kappa^2} (\delta G'(N) + 3\delta G(N)), \\ \frac{g'_\phi(N)}{2\kappa^2} \frac{\delta G(N)}{G_0(N)} + \frac{g''_\phi(N)}{2\kappa^2} \delta\phi(N) - \frac{g'_\phi(N)}{2\kappa^2} \frac{G'_0(N)}{G_0(N)} \delta\phi(N) - 2\tilde{K}^{(2)}(N) \left(\frac{\delta G(N)}{G_0(N)} + 2\delta\phi'(N) - \frac{G'_0(N)}{G_0(N)} \delta\phi(N) \right) & \\ = \frac{1}{2\kappa^2} \delta G'(N). \end{aligned} \quad (\text{III.93})$$

Then, we find

$$\begin{aligned} \begin{pmatrix} \delta\phi'(N) \\ \delta G'(N) \end{pmatrix} &= \frac{1}{L(N)} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} \delta\phi(N) \\ \delta G(N) \end{pmatrix}, \\ A &\equiv -\left(3 + \frac{G'_0(N)}{G_0(N)} \right) g'_\phi(N) + \frac{G'_0(N)}{2G_0(N)} L(N), \quad B \equiv \left(3 + \frac{g'_\phi(N)}{G_0(N)} \right) - \frac{L(N)}{2G_0(N)}, \\ C &\equiv (g''_\phi(N) + 3g'_\phi(N)) L(N) - g'_\phi(N)^2 \left(3 + \frac{G'_0(N)}{G_0(N)} \right), \quad D \equiv -3L(N) + g'_\phi(N) \left(3 + \frac{g'_\phi(N)}{G_0(N)} \right). \end{aligned} \quad (\text{III.94})$$

Here,

$$L(N) \equiv g'_\phi(N) + 8\kappa^2 \tilde{K}^{(2)}(N). \quad (\text{III.95})$$

The stability of the solution requires $A + D < 0$ and $AD - BC > 0$, which gives

$$\frac{1}{L(N)} \frac{g'_\phi(N)}{G_0(N)} (g'_\phi(N) - G'_0(N)) < 3 - \frac{G'_0(N)}{2G_0(N)}, \quad (\text{III.96})$$

$$\frac{1}{L(N)} (-2g'_\phi(N)g''_\phi(N) - 9g'_\phi(N)^2 - 6g''_\phi(N)G_0(N) + 3G'_0(N)g''_\phi(N) + 6G'_0(N)g'_\phi(N)) > g''_\phi(N). \quad (\text{III.97})$$

It might be possible to make the system stable by choosing $L(N)$ and, therefore, $\tilde{K}^{(2)}(N)$ properly. Neglecting the matter, one finds $g_\phi(N) = G_0(N)$ and obtains

$$3 > \frac{G'_0(N)}{2G_0(N)}, \quad \frac{1}{L(N)} (G'_0(N)G''_0(N) - 3G'_0(N)^2 - 6G''_0(N)G_0(N)) > G''_0(N). \quad (\text{III.98})$$

Thus, as long as the first condition is satisfied, the second condition can be always satisfied by choosing $L(N)$, and therefore $\tilde{K}^{(2)}(N)$, as an appropriate function. Using the above-developed general reconstruction scheme, any specific accelerating universe may be formulated as a solution of such a theory.

C. Modified $F(R)$ gravity

1. The reconstruction using the cosmological time

It is easy now to generalize the cosmological reconstruction scheme for modified gravity with $F(R)$ action (II.2) [18, 120, 121]. The action (II.2) is equivalently rewritten as

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} (P(\phi)R + Q(\phi)) + \mathcal{L}_{\text{matter}} \right\}. \quad (\text{III.99})$$

Here, P and Q are proper functions of the auxiliary scalar ϕ . The presence of an auxiliary scalar may simplify the formulation below, making it more similar to the one for scalar-tensor gravity. In fact, by the variation over ϕ , it follows that $0 = P'(\phi)R + Q'(\phi)$, which may be solved with respect to ϕ as $\phi = \phi(R)$. By substituting the obtained expression of $\phi(R)$ into (III.99), one arrives again at the $F(R)$ -gravity:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{F(R)}{2\kappa^2} + \mathcal{L}_{\text{matter}} \right\}, \quad F(R) \equiv P(\phi(R))R + Q(\phi(R)). \quad (\text{III.100})$$

By the variation of the action (III.99) with respect to the metric $g_{\mu\nu}$, the standard spatially flat FRW equations are obtained as:

$$0 = -6H^2 P(\phi) - Q(\phi) - 6H \frac{dP(\phi(t))}{dt} + 2\kappa^2 \rho_{\text{matter}}, \quad (\text{III.101})$$

$$0 = (4\dot{H} + 6H^2) P(\phi) + Q(\phi) + 2 \frac{d^2 P(\phi(t))}{dt^2} + 4H \frac{dP(\phi(t))}{dt} + 2\kappa^2 p_{\text{matter}}. \quad (\text{III.102})$$

Simple algebra leads to the following equation

$$0 = 2 \frac{d^2 P(\phi(t))}{dt^2} - 2H \frac{dP(\phi(t))}{dt} + 4\dot{H} P(\phi) + 2\kappa^2 (p_{\text{matter}} + \rho_{\text{matter}}). \quad (\text{III.103})$$

As one can redefine the scalar field ϕ freely, a very natural choice is $\phi = t$. It is assumed that ρ_{matter} and p_{matter} are the sum from the contributions of the matters with constant equation of state parameters w_i . In particular, when a combination of the radiation and dust is assumed, one gets the standard expression

$$\rho_{\text{matter}} = \rho_{r0} a^{-4} + \rho_{d0} a^{-3}, \quad p_{\text{matter}} = \frac{\rho_{r0}}{3} a^{-4}, \quad (\text{III.104})$$

with constant ρ_{r0} and ρ_{d0} . If the scale factor a is given by a proper function $g(t)$ as $a = a_0 e^{g(t)}$ with a constant a_0 , Eq. (III.102) reduces to the second-rank differential equation:

$$0 = 2 \frac{d^2 P(\phi)}{d\phi^2} - 2g'(\phi) \frac{dP(\phi)}{d\phi} + 4g''(\phi) P(\phi) + 2\kappa^2 \sum_i (1 + w_i) \rho_{i0} a_0^{-3(1+w_i)} e^{-3(1+w_i)g(\phi)}. \quad (\text{III.105})$$

In principle, by solving (III.105) the form of $P(\phi)$ may be found. Using (III.101) (or equivalently (III.102)), the form of $Q(\phi)$ follows:

$$Q(\phi) = -6(g'(\phi))^2 P(\phi) - 6g'(\phi) \frac{dP(\phi)}{d\phi} + 2\kappa^2 \sum_i \rho_{i0} a_0^{-3(1+w_i)} e^{-3(1+w_i)g(\phi)}. \quad (\text{III.106})$$

Thus, in principle, any given cosmology expressed as $a = a_0 e^{g(t)}$ can be realized as the solution of some specific (reconstructed) $F(R)$ gravity. Some explicit examples of such a reconstruction are presented below. Of course, the reconstruction is applied to cosmological eras that are fundamentally important in modern cosmology.

Model reproducing Λ CDM-type cosmology

Let us investigate whether Λ CDM-type cosmology can be reconstructed exactly by $F(R)$ gravity in the present formulation when we include dust, which can be a sum of the baryon and dark matter, and radiation.

In Einstein gravity, when there is matter with the EOS parameter w and cosmological constant, the FRW equation has the form (III.59). The solution of (III.59) is given by (III.60). Let us show how it is possible to reconstruct $F(R)$ gravity by reproducing such an epoch (III.60). When matter is included, Eq. (III.105) has the following form:

$$0 = 2 \frac{d^2 P(\phi)}{d\phi^2} - \frac{2}{l} \coth \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) \frac{dP(\phi)}{d\phi} - \frac{6(1+w)}{l^2} \sinh^{-2} \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) P(\phi) \quad (\text{III.107})$$

$$+ \frac{4}{3} \rho_{r0} a_0^{-4} \left(\alpha \sinh \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) \right)^{-8/3(1+w)} + \rho_{d0} a_0^{-3} \left(\alpha \sinh \left(\frac{3(1+w)}{2l} (\phi - t_s) \right) \right)^{-2/(1+w)}.$$

Because this equation is a linear inhomogeneous equation, its general solution is given by the sum of the special solution and of the general solution that corresponds to the homogeneous equation. For the case without matter, by changing the variable from ϕ to z as follows,

$$z \equiv -\sinh^{-2} \left(\frac{3(1+w)}{2l} (t - t_s) \right), \quad (\text{III.108})$$

Eq. (III.107) without matter can be rewritten in the form of the Gauss hypergeometric differential equation:

$$0 = z(1-z) \frac{d^2 P}{dz^2} + \left[\tilde{\gamma} - (\tilde{\alpha} + \tilde{\beta} + 1) z \right] \frac{dP}{dz} - \tilde{\alpha} \tilde{\beta} P,$$

$$\tilde{\gamma} \equiv 4 + \frac{1}{3(1+w)}, \quad \tilde{\alpha} + \tilde{\beta} + 1 \equiv 6 + \frac{1}{3(1+w)}, \quad \tilde{\alpha} \tilde{\beta} \equiv -\frac{1}{3(1+w)}, \quad (\text{III.109})$$

whose solution is given by Gauss' hypergeometric function:

$$P = P_0 F(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}; z) \equiv P_0 \frac{\Gamma(\tilde{\gamma})}{\Gamma(\tilde{\alpha})\Gamma(\tilde{\beta})} \sum_{n=0}^{\infty} \frac{\Gamma(\tilde{\alpha}+n)\Gamma(\tilde{\beta}+n)}{\Gamma(\tilde{\gamma}+n)} \frac{z^n}{n!}. \quad (\text{III.110})$$

Here, Γ is the Γ function. There is one more linearly independent solution like $(1-z)^{\tilde{\gamma}-\tilde{\alpha}-\tilde{\beta}} F(\tilde{\gamma}-\tilde{\alpha}, \tilde{\gamma}-\tilde{\beta}, \tilde{\gamma}; z)$, but we drop it here, for simplicity. Using (III.106), one finds the form of $Q(\phi)$:

$$Q = -\frac{6(1-z)P_0}{l^2} F(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}; z) - \frac{3(1+w)z(1-z)P_0}{l^2(13+12w)} F(\tilde{\alpha}+1, \tilde{\beta}+1, \tilde{\gamma}+1; z). \quad (\text{III.111})$$

From (III.108), it follows that $z \rightarrow 0$ when $t = \phi \rightarrow +\infty$. Then in the limit of $t = \phi \rightarrow +\infty$, one arrives at $P(\phi)R + Q(\phi) \rightarrow P_0 R - 6P_0/l^2$. Identifying $P_0 = 1/2\kappa^2$ and $\Lambda = 6/l^2$, the Einstein theory with cosmological constant Λ can be reproduced. The action is not singular even in the limit of $t \rightarrow \infty$. Therefore, even without the cosmological constant nor cold dark matter, the cosmology of the Λ CDM model can be reproduced by $F(R)$ -gravity.

We now investigate the special solution of (III.107). By changing the variable as in (III.108), the inhomogeneous differential equation looks as follows:

$$0 = z(1-z) \frac{d^2 P}{dz^2} + \left[\tilde{\gamma} - (\tilde{\alpha} + \tilde{\beta} + 1) z \right] \frac{dP}{dz} - \tilde{\alpha} \tilde{\beta} P + \eta (-z)^{-2(1+3w)/3(1+w)} + \xi (-z)^{-\frac{1+2w}{1+w}},$$

$$\eta \equiv \frac{4l^2}{27(1+w)} \rho_{r0} a_0^{-4} \alpha^{-8/3(1+w)}, \quad \zeta \equiv \frac{l^2}{3(1+w)} \rho_{d0} a_0^{-3} \alpha^{-2/(1+w)}. \quad (\text{III.112})$$

It is not trivial to find the solution of (III.112). Let us consider the case that $w = 0$ and $z \rightarrow -\infty$, that is, $t \rightarrow t_s$. In the limit of $t \rightarrow t_s$, Eq. (III.112) reduces to

$$0 = -z^2 \frac{d^2 P}{dz^2} + \tilde{\gamma} z \frac{dP}{dz} - \tilde{\alpha} \tilde{\beta} P + \eta (-z)^{-2/3}, \quad (\text{III.113})$$

whose special solution is given by

$$P = P_0(-z)^{-2/3}, \quad P_0 = \frac{\eta}{\frac{10}{9} - \frac{2(\tilde{\alpha} + \tilde{\beta} + 1)}{3} + \tilde{\alpha}\tilde{\beta}} = -\frac{9\eta}{25}. \quad (\text{III.114})$$

In principle, other special solutions of Eq. (III.112) could be found. This proves that, in the presence (or in the absence) of traditional matter, the standard Λ CDM cosmology can be reproduced exactly by $F(R)$ gravity.

2. Reconstructed model realizing the phantom divide crossing

Using the above-developed method, we reconstruct an explicit model in which a crossing of the phantom divide can be realized [135]. In other words, the modified $F(R)$ theory that describes the transition from acceleration to superacceleration (phantom era). will be presented.

A solution of Eq. (III.105) without matter can be given by

$$P(\phi) = e^{\tilde{g}(\phi)/2} \tilde{p}(\phi), \quad (\text{III.115})$$

$$\tilde{g}(\phi) = -10 \ln \left[\left(\frac{\phi}{t_0} \right)^{-\gamma} - C \left(\frac{\phi}{t_0} \right)^{\gamma+1} \right], \quad (\text{III.116})$$

$$\tilde{p}(\phi) = \tilde{p}_+ \phi^{\beta_+} + \tilde{p}_- \phi^{\beta_-}, \quad (\text{III.117})$$

$$\beta_{\pm} = \frac{1 \pm \sqrt{1 + 100\gamma(\gamma+1)}}{2}, \quad (\text{III.118})$$

where γ and C are positive constants, t_0 is the present time, and $\tilde{p}_{\pm}$ are arbitrary constants.

From Eq. (III.116), it follows that $\tilde{g}(\phi)$ diverges at finite ϕ when

$$\phi = t_s \equiv t_0 C^{-1/(2\gamma+1)}, \quad (\text{III.119})$$

which shows that the Big Rip singularity could occur at $t = t_s$ [1]. One only needs to consider the period $0 < t < t_s$ because $\tilde{g}(\phi)$ should be a real number. Eq. (III.116) also gives the following Hubble rate $H(t)$:

$$H(t) = \frac{d\tilde{g}(\phi)}{d\phi} = \left(\frac{10}{t_0} \right) \left[\frac{\gamma \left(\frac{\phi}{t_0} \right)^{-\gamma-1} + (\gamma+1)C \left(\frac{\phi}{t_0} \right)^{\gamma}}{\left(\frac{\phi}{t_0} \right)^{-\gamma} - C \left(\frac{\phi}{t_0} \right)^{\gamma+1}} \right], \quad (\text{III.120})$$

where it is taken that $\phi = t$.

In the FRW background (II.6), as in (II.3), the effective energy density and pressure of the universe are given by $\rho_{\text{eff}} = 3H^2/\kappa^2$ and $p_{\text{eff}} = -(2\dot{H} + 3H^2)/\kappa^2$, respectively. Then, the effective EoS parameter w_{eff} is defined as in (II.4). For the case of $H(t)$ of Eq. (III.120), from Eq. (II.4), we find that w_{eff} is expressed as

$$w_{\text{eff}} = -1 + U(t), \quad (\text{III.121})$$

where

$$U(t) \equiv -\frac{2\dot{H}}{3H^2} = -\frac{-\gamma + 4\gamma(\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} + (\gamma+1) \left(\frac{t}{t_s} \right)^{2(2\gamma+1)}}{15 \left[\gamma + (\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]^2}. \quad (\text{III.122})$$

For the case of Eq. (III.120), the scalar curvature $R = 6(\dot{H} + 2H^2)$ is given by

$$R = \frac{60 \left[\gamma(20\gamma-1) + 44\gamma(\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} + (\gamma+1)(20\gamma+21) \left(\frac{t}{t_s} \right)^{2(2\gamma+1)} \right]}{t^2 \left[1 - \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]^2}. \quad (\text{III.123})$$

In deriving Eqs. (III.122) and (III.123), Eq. (III.119) was used.

When $t \rightarrow 0$, i.e., $t \ll t_s$, $H(t)$ behaves as

$$H(t) \sim \frac{10\gamma}{t}. \quad (\text{III.124})$$

In this limit, it follows from Eq. (II.4) that the effective EoS parameter is given by

$$w_{\text{eff}} = -1 + \frac{1}{15\gamma}. \quad (\text{III.125})$$

This behavior is identical to that in Einstein gravity with matter with an EoS parameter greater than -1 .

On the other hand, when $t \rightarrow t_s$, one finds that

$$H(t) \sim \frac{10}{t_s - t}. \quad (\text{III.126})$$

In this case, the scale factor is given by $a(t) \sim a_0 (t_s - t)^{-10}$. When $t \rightarrow t_s$, therefore, $a \rightarrow \infty$, namely, the Big Rip singularity appears. In this limit, the effective EoS parameter is given by

$$w_{\text{eff}} = -1 - \frac{1}{15} = -\frac{16}{15}. \quad (\text{III.127})$$

This behavior is identical with that of the case in which there is phantom matter that has an EoS parameter smaller than -1 . Thus, we have obtained an explicit model showing a crossing of the phantom divide.

It follows from Eq. (II.4) that the effective EoS parameter w_{eff} is equal to -1 when $\dot{H} = 0$. Solving $w_{\text{eff}} = -1$ with respect to t by using Eq. (III.121), namely, $U(t) = 0$, we find that the effective EoS parameter crosses the phantom divide at $t = t_c$ given by

$$t_c = t_s \left(-2\gamma + \sqrt{4\gamma^2 + \frac{\gamma}{\gamma+1}} \right)^{1/(2\gamma+1)}. \quad (\text{III.128})$$

From Eq. (III.122), one sees that when $t < t_c$, $U(t) > 0$ because $\gamma > 0$. Moreover, the time derivative of $U(t)$ is given by

$$\frac{dU(t)}{dt} = -\frac{2\gamma(\gamma+1)(2\gamma+1)^2}{15 \left[\gamma + (\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]^3} \left(\frac{1}{t_s} \right) \left(\frac{t}{t_s} \right)^{2\gamma} \left[1 - \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]. \quad (\text{III.129})$$

Eq. (III.129) shows that the relation $dU(t)/(dt) < 0$ is always satisfied because only the period $0 < t < t_s$ is considered as mentioned above. This means that $U(t)$ decreases monotonically. Thus, the value of $U(t)$ evolves from positive to negative. From Eq. (III.121), we see that the value of w_{eff} crosses -1 . Once the universe enters the phantom phase, it stays in this phase, namely, the value of w_{eff} remains less than -1 , and finally the Big Rip singularity occurs because $U(t)$ decreases monotonically. Note that other types of finite-time future singularities in modified gravity are possible as demonstrated after Eq. (II.96) in the second chapter (see also ref. [39]). It follows from Eqs. (III.115), (III.116), (III.117) and (III.119) that $P(t)$ is given by

$$P(t) = \left[\frac{\left(\frac{t}{t_0} \right)^\gamma}{1 - \left(\frac{t}{t_s} \right)^{2\gamma+1}} \right]^5 \sum_{j=\pm} \tilde{p}_j t^{\beta_j}. \quad (\text{III.130})$$

Using Eq. (III.130), one gets

$$Q(t) = -6H \left[\frac{\left(\frac{t}{t_0} \right)^\gamma}{1 - \left(\frac{t}{t_s} \right)^{2\gamma+1}} \right]^5 \sum_{j=\pm} \left(\frac{3}{2}H + \frac{\beta_j}{t} \right) \tilde{p}_j t^{\beta_j}. \quad (\text{III.131})$$

If one can solve Eq. (III.123) with respect to t as $t = t(R)$, in principle we can obtain the form of $F(R)$ by using this solution and Eqs. (III.100), (III.130) and (III.131). In fact, however, for the general case it is difficult to solve




FIG. 2: Behavior of $t_s^2 F(\tilde{R})$ as a function of $\tilde{R}$ for $\gamma = 1/2$, $\tilde{p}_+ = -1/t_s^{\beta_+}$, $\tilde{p}_- = 0$, $\beta_+ = (1 + 2\sqrt{19})/2$ and $t_s = 2t_0$.

Eq. (III.123) as $t = t(R)$. Thus, as a solvable example, we show the behavior of $t_s^2 F(\tilde{R})$ as a function of $\tilde{R} \equiv t_s^2 R$ in Fig. 1 for $\gamma = 1/2$, $\tilde{p}_+ = -1/t_s^{\beta_+}$, $\tilde{p}_- = 0$, $\beta_+ = (1 + 2\sqrt{19})/2$ and $t_s = 2t_0$. The quantities in Fig. III C 2 are shown as dimensionless value. The horizontal and vertical axes show $\tilde{R}$ and $t_s^2 F$, respectively. (Here, $\tilde{R} = t_s^2 R = 4R/R_0$, where R_0 is the current curvature. In deriving this relation, we have used $t_s = 2t_0$, $t_0 \approx H_0^{-1}$, where H_0 is the present Hubble parameter.) From Fig. 1, we see that the value of $F(R)$ increases as that of R becomes larger.

To examine the analytic form of $F(R)$ for the general case, we will now investigate the behavior of $F(R)$ in the limits $t \rightarrow 0$ and $t \rightarrow t_s$. When $t \rightarrow 0$, from Eq. (III.124) we find

$$t \sim \sqrt{\frac{60\gamma(20\gamma - 1)}{R}}. \quad (\text{III.132})$$

In this limit, it follows from Eqs. (III.100), (III.124), (III.130), (III.131) and (III.132) that the form of $F(R)$ is given by

$$F(R) \sim \left\{ \frac{\left[\frac{1}{t_0} \sqrt{60\gamma(20\gamma - 1)} R^{-1/2} \right]^\gamma}{1 - \left[\frac{1}{t_s} \sqrt{60\gamma(20\gamma - 1)} R^{-1/2} \right]^{2\gamma+1}} \right\}^5 R \times \sum_{j=\pm} \left\{ \left(\frac{5\gamma - 1 - \beta_j}{20\gamma - 1} \right) \tilde{p}_j [60\gamma(20\gamma - 1)]^{\beta_j/2} R^{-\beta_j/2} \right\}. \quad (\text{III.133})$$

On the other hand, when $t \rightarrow t_s$, from Eq. (III.126) we obtain

$$t \sim t_s - 3\sqrt{\frac{140}{R}}. \quad (\text{III.134})$$

In this limit, it follows from Eqs. (III.100), (III.126), (III.130), (III.131) and (III.134) that the form of $F(R)$ is given by

$$F(R) \sim \left(\frac{\left\{ \frac{1}{t_0} [t_s - 3\sqrt{140}R^{-1/2}] \right\}^\gamma}{1 - \left[1 - \frac{3\sqrt{140}}{t_s} R^{-1/2} \right]^{2\gamma+1}} \right)^5 R \sum_{j=\pm} \tilde{p}_j [t_s - 3\sqrt{140}R^{-1/2}]^{\beta_j} \times \left\{ 1 - \sqrt{\frac{20}{7}} \left[\sqrt{\frac{15}{84}} t_s + (\beta_j - 15) R^{-1/2} \right] \frac{1}{t_s - 3\sqrt{140}R^{-1/2}} \right\}. \quad (\text{III.135})$$

The above modified gravity may be considered as an approximated form of a more realistic and viable theory. For large R , namely, $t_s^2 R \gg 1$, the expression of $F(R)$ in (III.135) can be written approximately as

$$F(R) \approx \frac{2}{7} \left[\frac{1}{3\sqrt{140}(2\gamma+1)} \left(\frac{t_s}{t_0} \right)^\gamma \right]^5 \left(\sum_{j=\pm} \tilde{p}_j t_s^{\beta_j} \right) t_s^5 R^{7/2}. \quad (\text{III.136})$$

Note also that viable models that lead to a Big Rip singularity as is discussed in the second chapter necessarily predict the phantom divide crossing. This gives exact (non-asymptotic) and realistic modified gravities that show such phenomena. Moreover, some of the viable models with accelerating solutions that are asymptotically non-singular or tend to softer Type II, Type III and Type IV singularities may also show phantom divide crossing. However, in this case the phantom era turns out to be transient.

3. The reconstruction using e-folding

Let us demonstrate the reconstruction of $F(R)$ gravity in terms of e-folding. Moreover, unlike the previous subsection, the slightly different approach is developed without the introduction of an auxiliary scalar. We introduce the e-folding instead of the cosmological time t as: $N = \ln \frac{a}{a_0}$. The variable N is related to the redshift z by

$$e^{-N} = \frac{a_0}{a} = 1 + z. \quad (\text{III.137})$$

Because $\frac{d}{dt} = H \frac{d}{dN}$ and, therefore, $\frac{d^2}{dt^2} = H^2 \frac{d^2}{dN^2} + H \frac{dH}{dN} \frac{d}{dN}$, one can rewrite (II.7) by

$$0 = -\frac{F(R)}{2} + 3(H^2 + HH')F'(R) - 18(4H^3H' + H^2(H')^2 + H^3H'')F''(R) + \kappa^2 \rho_{\text{matter}}. \quad (\text{III.138})$$

Here, $H' \equiv dH/dN$ and $H'' \equiv d^2H/dN^2$. As usual, the matter energy density ρ_{matter} is given by the sum of the fluid densities with the constant EoS parameter w_i

$$\rho_{\text{matter}} = \sum_i \rho_{i0} a^{-3(1+w_i)} = \sum_i \rho_{i0} a_0^{-3(1+w_i)} e^{-3(1+w_i)N}. \quad (\text{III.139})$$

Let the Hubble rate be given in terms of N via the function $g(N)$ as

$$H = g(N) = g(-\ln(1+z)). \quad (\text{III.140})$$

Then, the scalar curvature takes the form: $R = 6g'(N)g(N) + 12g(N)^2$, which can be solved with respect to N as $N = N(R)$. Using (III.139) and (III.140), one can rewrite (III.138) as

$$\begin{aligned} 0 = & -18 \left(4g(N(R))^3 g'(N(R)) + g(N(R))^2 g'(N(R))^2 + g(N(R))^3 g''(N(R)) \right) \frac{d^2 F(R)}{dR^2} \\ & + 3 \left(g(N(R))^2 + g'(N(R))g(N(R)) \right) \frac{dF(R)}{dR} - \frac{F(R)}{2} + \kappa^2 \sum_i \rho_{i0} a_0^{-3(1+w_i)} e^{-3(1+w_i)N(R)}, \end{aligned} \quad (\text{III.141})$$

which constitutes a differential equation for $F(R)$, where the variable is the scalar curvature R . Instead of g , if $G(N) \equiv g(N)^2 = H^2$ is used, the expression (III.141) can be simplified as:

$$\begin{aligned} 0 = & -9G(N(R))(4G'(N(R)) + G''(N(R))) \frac{d^2 F(R)}{dR^2} + \left(3G(N(R)) + \frac{3}{2}G'(N(R)) \right) \frac{dF(R)}{dR} \\ & - \frac{F(R)}{2} + \kappa^2 \sum_i \rho_{i0} a_0^{-3(1+w_i)} e^{-3(1+w_i)N(R)}. \end{aligned} \quad (\text{III.142})$$

Note that $R = 3G'(N) + 12G(N)$. Thus, when we find an $F(R)$ that satisfies the differential equation (III.141) or (III.142), and such $F(R)$ theory yields a solution (III.140). Thus, such reconstructed $F(R)$ gravity realizes the above cosmological solution. It is interesting that the above scheme may be applied not only to general relativity but also to modified gravity (partial reconstruction). In other words, one can start from modified gravity which consistently describes the entire sequence of the universe evolution eras and satisfies local tests; however, some of its predictions seem to contradict the observational data. In this case, one can apply the partial reconstruction of such a model (for instance, slightly modifying it at the very early universe) so that the above contradiction is resolved but all good properties of the theory are preserved. This provides a way to avoid any future exclusion of a non-realistic modified gravity to arrive at more narrow and more realistic sub-class of such theories.

4. Stability of the cosmological solution

In this section, we investigate the condition of the stability of the given cosmological solution. The FRW equation (II.7) can be rewritten by using $G = H^2$ and the e-folding N as follows (III.142):

$$0 = -9G(N) (4G'(N) + G''(N)) \frac{d^2 F(R)}{dR^2} \Big|_{R=3G'(N)+12G(N)} + \left(3G(N) + \frac{3}{2}G'(N) \right) \frac{dF(R)}{dR} \Big|_{R=3G'(N)+12G(N)} - \frac{F(R)}{2} \Big|_{R=3G'(N)+12G(N)} + \kappa^2 \rho_{\text{matter}}(N). \quad (\text{III.143})$$

Assume that the N -dependence of ρ_{matter} is known as in (III.139). Let a solution of (III.143) be $G = G_0(N)$, and consider the perturbation from the following solution:

$$G(N) = G_0(N) + \delta G(N). \quad (\text{III.144})$$

Then, Eq. (III.143) gives

$$0 = G_0(N) \frac{d^2 F(R)}{dR^2} \Big|_{R=3G'_0(N)+12G_0(N)} \delta G''(N) + \left\{ 3G_0(N) (4G'_0(N) + G''_0(N)) \frac{d^3 F(R)}{dR^3} \Big|_{R=3G'_0(N)+12G_0(N)} + \left(3G_0(N) - \frac{1}{2}G'_0(N) \right) \frac{d^2 F(R)}{dR^2} \Big|_{R=3G'_0(N)+12G_0(N)} \right\} \delta G'(N) + \left\{ 12G_0(N) (4G'_0(N) + G''_0(N)) \frac{d^3 F(R)}{dR^3} \Big|_{R=3G'_0(N)+12G_0(N)} + (-4G_0(N) + 2G'_0(N) + G''_0(N)) \frac{d^2 F(R)}{dR^2} \Big|_{R=3G'_0(N)+12G_0(N)} + \frac{1}{3} \frac{dF(R)}{dR} \Big|_{R=3G'_0(N)+12G_0(N)} \right\} \delta G. \quad (\text{III.145})$$

Note that in this formulation, as it is assumed that the N -dependence of ρ_{matter} is known, we do not need to consider the fluctuation of ρ_{matter} . Using the cosmological time t instead of N , we need to include the fluctuation of ρ_{matter} . Then, the conditions of the stability are given by

$$6 (4G'_0(N) + G''_0(N)) \frac{d^3 F(R)}{dR^3} \Big|_{R=3G'_0(N)+12G_0(N)} \left(\frac{d^2 F(R)}{dR^2} \Big|_{R=3G'_0(N)+12G_0(N)} \right)^{-1} + 6 - \frac{G'_0(N)}{G_0(N)} > 0, \quad (\text{III.146})$$

$$36 (4G'_0(N) + G''_0(N)) \frac{d^3 F(R)}{dR^3} \Big|_{R=3G'_0(N)+12G_0(N)} \left(\frac{d^2 F(R)}{dR^2} \Big|_{R=3G'_0(N)+12G_0(N)} \right)^{-1} - 12 + \frac{6G'_0(N)}{G_0(N)} + \frac{3G''_0(N)}{G_0(N)} + \frac{1}{G_0(N)} \frac{dF(R)}{dR} \Big|_{R=3G'_0(N)+12G_0(N)} \left(\frac{d^2 F(R)}{dR^2} \Big|_{R=3G'_0(N)+12G_0(N)} \right)^{-1} > 0. \quad (\text{III.147})$$

In the case of de Sitter space, where H and, therefore, G_0 and $R = R_0 \equiv 12G_0$ are constants, Eq. (III.146) gives $6 > 0$ and is trivially satisfied. Eq. (III.147) becomes

$$-12G_0 + \frac{dF(R)}{dR} \Big|_{R=12G_0} \left(\frac{d^2 F(R)}{dR^2} \Big|_{R=12G_0} \right)^{-1} = -R_0 + \frac{dF(R)}{dR} \Big|_{R=R_0} \left(\frac{d^2 F(R)}{dR^2} \Big|_{R=R_0} \right)^{-1} > 0, \quad (\text{III.148})$$

which is the standard result.

One may consider the case that $F(R)$ gravity admits two de Sitter solutions. If one is stable but another is unstable, there could be a solution in which the unstable de Sitter solution could transit to the stable solution. If the Hubble rate H in the unstable solution is much larger than the Hubble rate in the stable one, the unstable solution may correspond to the inflation and the stable one to the late-time acceleration. Note that we may directly construct an $F(R)$ theory that admits the transition from an asymptotic de Sitter universe with a large Hubble rate to another

asymptotic de Sitter universe with a small Hubble rate. Then, if the solution and $F(R)$ satisfy the conditions (III.146) and (III.147), such a transition occurs.

Consider the stability of a de Sitter universe satisfying the condition (II.9), or equivalently (II.37). Eq. (II.9) can be rewritten as

$$0 = \frac{d}{dR} \left(\frac{F(R)}{R^2} \right). \quad (\text{III.149})$$

Let $R = R_0$ be a solution of (III.149). Then, $F(R)$ has the following form:

$$\frac{F(R)}{R^2} = f_0 + f(R) (R - R_0)^n. \quad (\text{III.150})$$

Here, f_0 is a constant, which should be positive if we require $F(R) > 0$ and n to be integers greater than or equal to two: $n \geq 2$. Assume that the function $f(R)$ does not vanish at $R = R_0$, $f(R_0) \neq 0$. When $n = 2$,

$$-R_0 + \frac{F'(R_0)}{F''(R_0)} = -\frac{f(R_0)A_0}{f_0 + f(A_0)}. \quad (\text{III.151})$$

Then, Eq. (III.148) shows that the de Sitter solution is stable if

$$-f_0 < f(A_0) < 0. \quad (\text{III.152})$$

When $n \geq 3$, one finds

$$-R_0 + \frac{F'(R_0)}{F''(R_0)} = 0. \quad (\text{III.153})$$

It is necessary to conduct a more detailed investigation to check the stability. Using the expression of m_σ^2 in (II.30), one can investigate the sign of m_σ^2 in the region $R \sim R_0$. Note that the expression (II.38) is not used because the expression is only valid at the point $R = R_0$. Thus, one gets

$$m_\sigma^2 \sim -\frac{3n(n-1)R_0^2 f(R_0)}{2f_0^2} (R - R_0)^{n-2}. \quad (\text{III.154})$$

Eq. (III.154) indicates that when n is an even integer, the de Sitter solution is stable if $f(R_0) < 0$ but unstable if $f(R_0) > 0$. On the other hand, when n is an odd integer, the de Sitter solution is always unstable. Note, however, that when $f(R_0) < 0$ ($f(R_0) > 0$), we find $m_\sigma^2 > 0$ ($m_\sigma^2 < 0$) if $R > R_0$ but $m_\sigma^2 < 0$ ($m_\sigma^2 > 0$) if $R < R_0$. Therefore, when $f(R_0) < 0$, R becomes smaller, but when $f(R_0) > 0$, R becomes larger. The stability condition may be used to obtain realistic (unstable) de Sitter inflation for specific $F(R)$ gravity.

5. FRW cosmology in $F(R)$ gravity with a Lagrange multiplier

Here, we consider $F(R)$ gravity with a Lagrange multiplier field, following ref. [136]. In usual $F(R)$ gravity, the scalar mode called scalaron appears, which often affects Newton's law. In this section, we try to suppress the propagation of the scalaron by imposing the constraint under the Lagrange multiplier field. As a result, however, a propagating mode in the Lagrange multiplier field seems to appear, which may break Newton's law but in an easier way. The solution of this question may require additional modifications of the constraints. Another purpose of this section is the reconstruction. In usual $F(R)$ gravity, one needs to solve the complicated differential equation developed in the previous sections to realize the reconstruction. It is shown that the reconstruction can be realized more easily in the model with the Lagrange multiplier field.

The starting action is given by

$$S = \int d^4x \sqrt{-g} \left\{ F_1(R) - \lambda \left(\frac{1}{2} \partial_\mu R \partial^\mu R + F_2(R) \right) \right\}. \quad (\text{III.155})$$

Here, λ is the Lagrange multiplier field, which gives a constraint of

$$\frac{1}{2} \partial_\mu R \partial^\mu R + F_2(R) = 0. \quad (\text{III.156})$$

On the other hand, by the variation over the metric $g_{\mu\nu}$, the equation of motion follows:

$$0 = \frac{1}{2}g_{\mu\nu}F_1(R) + \frac{\lambda}{2}\partial_\mu R\partial_\nu R + (-R_{\mu\nu} + \nabla_\mu\nabla_\nu - g_{\mu\nu}\nabla^2)(F'_1(R) - \lambda F'_2(R) - \nabla^\mu(\lambda\nabla_\mu R)) . \quad (\text{III.157})$$

If the Ricci tensor is covariantly constant and the scalar curvature is a constant,

$$R_{\mu\nu} = \frac{R_0}{4}g_{\mu\nu} , \quad R = R_0 , \quad (\text{III.158})$$

Eqs. (III.156) and (III.157) reduce to

$$0 = F_2(R_0) , \quad (\text{III.159})$$

$$0 = F_1(R_0) - \frac{1}{2}R_0(F'_1(R_0) - \lambda F'_2(R_0)) . \quad (\text{III.160})$$

If Eq. (III.159) has a solution, Eq. (III.160) can be solved with respect to the Lagrange multiplier field:

$$\lambda = \frac{-F_1(R_0) + R_0 F'_1(R_0)}{F'_2(R_0)} . \quad (\text{III.161})$$

Then, if R_0 is positive, the above solution describes de Sitter space-time, which may correspond to dark energy or the inflationary epoch. The spatially flat FRW metric Eqs. (III.156) and $(\mu, \nu) = (0, 0)$ -component of (III.157) have the following form:

$$0 = -\frac{1}{2}\dot{R}^2 + F_2(R) , \quad (\text{III.162})$$

$$0 = -\frac{1}{2}F_1(R) + 18\lambda(\ddot{H} + 4H\dot{H})^2 + \left\{ 3(\dot{H} + H^2) - 3H\frac{d}{dt} \right\} \left\{ F'_1(R) - \lambda F'_2(R) + \left(\frac{d}{dt} + 3H \right) \left(\lambda \frac{dR}{dt} \right) \right\} . \quad (\text{III.163})$$

When $F_2(R) > 0$, Eq. (III.162) results in

$$t = \int^R \frac{dR}{\sqrt{2F_2(R)}} , \quad (\text{III.164})$$

which can be solved with respect to R as a function of t $R = F_R(t)$. Because

$$R = 6\frac{dH}{dt} + 12H^2 , \quad (\text{III.165})$$

one can find the behavior of $H = \frac{\dot{a}}{a}$ by solving the differential equation

$$6\frac{dH}{dt} + 12H^2 = F_R(t) . \quad (\text{III.166})$$

Using the obtained solution for $H = H(t)$ (and $R = F_R(t)$), Eq. (III.163) becomes a differential equation of the multiplier field λ , and the behavior of λ : $\lambda = \lambda(t)$ follows. (Note that FRW cosmology in scalar theory with a Lagrange multiplier constraint was studied in refs. [137]).

Conversely, when the behavior of $H(t)$ is known from the observational data, one may reconstruct $F_2(R)$ to reproduce $H(t)$ by using (III.162). $H(t)$ gives the behavior of R as $R = R(t)$, which can be solved with respect to t as $t = t(R)$. Using (III.162), the explicit form of $F_2(R)$ is found to be

$$F_2(R) = \frac{1}{2} \left(\frac{dR}{dt} \right)^2 \Big|_{t=t(R)} . \quad (\text{III.167})$$

Note that $F_1(R)$ can be an arbitrary function. Thus, the cosmological reconstruction of the model can be performed more easily than that in the usual $F(R)$ gravity.

As an explicit example, one may consider

$$H(t) = \frac{h_0}{t}, \quad (\text{III.168})$$

where h_0 is a constant. Here,

$$R = \frac{-6h_0 + 12h_0^2}{t^2} \quad \text{or} \quad t = \sqrt{\frac{-6h_0 + 12h_0^2}{R}}. \quad (\text{III.169})$$

Therefore, we find

$$\frac{dR}{dt} = -\frac{12(-h_0 + 2h_0^2)}{t^3} = -\frac{2R^{\frac{3}{2}}}{\sqrt{6(-h_0 + 2h_0^2)}}, \quad (\text{III.170})$$

which gives

$$F_2(R) = \frac{R^3}{12(-h_0 + 2h_0^2)}. \quad (\text{III.171})$$

Another example is given by

$$R = \frac{R_-}{2}(1 - \tanh \omega t) + \frac{R_+}{2}(1 + \tanh \omega t). \quad (\text{III.172})$$

Here, $R_{\pm}$ and ω are constants. When $t \rightarrow \pm\infty$, $R \rightarrow \pm R_{\pm}$ and, therefore, the space-time becomes asymptotically de Sitter. One may identify the epoch of $t \rightarrow -\infty$ as inflation and $t \rightarrow +\infty$ as late acceleration. Because

$$\dot{R} = \frac{(R_- - R_+)\omega}{2 \cosh^2 \omega t} = \frac{(R_- - R_+)\omega}{2} \left(1 - \frac{(R_- + R_+ - 2R)^2}{(R_- - R_+)^2} \right), \quad (\text{III.173})$$

from Eq. (III.167), one gets

$$F_2(R) = \frac{(R_- - R_+)^2 \omega^2}{8} \left(1 - \frac{(R_- + R_+ - 2R)^2}{(R_- - R_+)^2} \right)^2, \quad (\text{III.174})$$

Thus, the unification of early-time inflation with dark energy epoch is possible also in Lagrange multiplier constrained modified gravity.

Thus, the universe evolution only depends on the constraint equation (III.162) and not on $F_1(R)$. $F_1(R)$ can affect the correction to Newton's law. In convenient $F_1(R)$ cosmology, all of the dynamics are defined by the form of this function. With the constraint (III.162), $F_1(R)$ becomes irrelevant. The cosmological dynamics are defined by the form of $F_2(R)$. In convenient $F(R)$ gravity, the scalar propagating mode appears, which often violates Newton's law. One can show that such a scalar does not propagate in the theory under investigation. Indeed, Eq. (III.157) contains the second derivative of the multiplier field λ . We now need to solve the second-order differential equation to find λ , which can indicate that λ could propagate and that a correction to Newton's law can be found. The magnitude of the correction depends on the choice of $F_1(R)$ and/or $F_2(R)$.

To investigate the regime of Newton's law, $F_1(R)$ is chosen as

$$F_1(R) = \frac{R}{2\kappa^2}. \quad (\text{III.175})$$

The matter presence is assumed. Eq. (III.157) has the following form:

$$0 = \frac{1}{2\kappa^2} \left(\frac{1}{2} g_{\mu\nu} R - R_{\mu\nu} \right) + \frac{1}{2} T_{\mu\nu} - (-R_{\mu\nu} + \nabla_\mu \nabla_\nu - g_{\mu\nu} \nabla^2) (\lambda F_2'(R) - \nabla^\mu (\lambda \nabla_\mu R)). \quad (\text{III.176})$$

For the solution with $\lambda = 0$, Eq. (III.157) reduces to the Einstein equation,

$$0 = \frac{1}{\kappa^2} \left(\frac{1}{2} g_{\mu\nu} R - R_{\mu\nu} \right) + T_{\mu\nu}. \quad (\text{III.177})$$

Here, $T_{\mu\nu}$ is the matter energy-momentum tensor. Without matter $T_{\mu\nu} = 0$, the Schwarzschild space-time with $R = R_{\mu\nu} = 0$ is a solution, which also satisfies the constraint equation (III.156) if $F_2(0) = 0$. In the case with matter $T_{\mu\nu} \neq 0$, however, the Einstein equation (III.177) gives

$$R = -\kappa^2 T. \quad (\text{III.178})$$

Here, T is the trace of the energy-momentum tensor. The constraint equation (III.156) is rewritten to

$$0 = \frac{\kappa^4}{2} \partial_\mu T \partial^\mu T + F_2(-\kappa^2 T), \quad (\text{III.179})$$

which is not always satisfied. Thus, in the presence of matter, the constraint equation (III.156) should be modified to be

$$0 = \frac{1}{2} \partial_\mu R \partial^\mu R + F_2(R) - \frac{\kappa^4}{2} \partial_\mu T \partial^\mu T - F_2(-\kappa^2 T). \quad (\text{III.180})$$

This indicates that the total constrained action with matter should be, instead of (III.155),

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \lambda \left\{ \frac{1}{2} \partial_\mu R \partial^\mu R + F_2(R) - \frac{\kappa^4}{2} \partial_\mu T \partial^\mu T - F_2(-\kappa^2 T) \right\} + \mathcal{L}_{\text{matter}} \right]. \quad (\text{III.181})$$

In this case, Newton's law can be easily reproduced. Of course, some other form of the constraint may also solve this problem.

We also note that the form of the constraint could be correct when $F_1(R)$ is given by (III.175). For general form of $F_1(R)$, the constraint should be changed.

Because T vanishes in the vacuum, the constraint (III.180) reduces to

$$0 = \frac{1}{2} \partial_\mu R \partial^\mu R + F_2(R) - F_2(0). \quad (\text{III.182})$$

If $F_2(0) = 0$, the constraint (III.180) gives (III.156) and (III.162) and the cosmological evolution can be generated. Note that $F_2(R)$ (III.171) satisfies the condition $F_2(0) = 0$ but $F_2(R)$ in (III.151) does not. When the condition $F_2(0) = 0$ is satisfied, there are two classes of solutions for the constraint (III.162). One is a trivial solution $R = 0$, and another corresponds to the non-trivial cosmological evolution given by (III.164). Near solar systems and galaxies, the solution could correspond to the trivial one with $R = 0$ to reproduce Newton's law, but at large scales, the solution should correspond to (III.164), so that the evolution of the universe can be generated. It is not so trivial to prove or to deny that the two solutions are connected in the intermediate region between the galaxy scales and large-scale universe. More careful (possibly numerical) analysis should be done in order to solve this problem.

D. $F(\mathcal{G})$ gravity

The cosmological reconstruction scheme developed above can be applied to an arbitrary modified gravity. As one more example of its application, we discuss the reconstruction in modified Gauss-Bonnet gravity. The action (II.123) can be rewritten by using the auxiliary scalar field ϕ :

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - V(\phi) - \xi(\phi) \mathcal{G} \right]. \quad (\text{III.183})$$

By the variation over the scalar field ϕ , an algebraic equation is obtained:

$$0 = V'(\phi) + \xi'(\phi) \mathcal{G}. \quad (\text{III.184})$$

It can be solved, at least locally, with respect to ϕ as $\phi = \phi(\mathcal{G})$. Inserting the obtained expression of $\phi(\mathcal{G})$ into the action (III.183), we find the action (II.123):

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} + f(\mathcal{G}) \right]. \quad (\text{III.185})$$

Here,

$$f(\mathcal{G}) \equiv -V(\phi(\mathcal{G})) + \xi(\phi(\mathcal{G})) \mathcal{G}. \quad (\text{III.186})$$

In the following, we use the results obtained in refs. [52–54] and consider the reconstruction of scalar- $F(\mathcal{G})$ gravity. By the variation over the metric in the FRW universe (II.6), one obtains the following equations:

$$\begin{aligned} 0 &= -\frac{3}{\kappa^2}H^2 + V(\phi) + 24H^3 \frac{d\xi(\phi(t))}{dt}, \\ 0 &= \frac{1}{\kappa^2} \left(2\dot{H} + 3H^2 \right) - V(\phi) - 8H^2 \frac{d^2\xi(\phi(t))}{dt^2} - 16H\dot{H} \frac{d\xi(\phi(t))}{dt} - 16H^3 \frac{d\xi(\phi(t))}{dt}. \end{aligned} \quad (\text{III.187})$$

Using (III.187), it follows that

$$\begin{aligned} \xi(\phi(t)) &= \frac{1}{8} \int^t dt_1 \frac{a(t_1)}{H(t_1)^2} W(t_1), \\ V(\phi(t)) &= \frac{3}{\kappa^2} H(t)^2 - 3a(t)H(t)W(t), \quad W(t) \equiv \frac{2}{\kappa^2} \int^t \frac{dt_1}{a(t_1)} \dot{H}(t_1). \end{aligned} \quad (\text{III.188})$$

Because there is no kinetic term of ϕ , one can redefine ϕ properly and even identify ϕ with the cosmological time: $\phi = t$. In the same way as in the previous section, if we consider the following $V(\phi)$ and ξ given in term of a single function g ,

$$\begin{aligned} V(\phi) &= \frac{3}{\kappa^2} g'(\phi)^2 - 3g'(\phi) e^{g(\phi)} U(\phi), \\ \xi(\phi) &= \frac{1}{8} \int^\phi d\phi_1 \frac{e^{g(\phi_1)}}{g'(\phi_1)^2} U(\phi_1), \quad U(\phi) \equiv \frac{2}{\kappa^2} \int^\phi d\phi_1 e^{-g(\phi_1)} g''(\phi_1), \end{aligned} \quad (\text{III.189})$$

the explicit solution follows:

$$a = a_0 e^{g(t)} \quad (H = g'(t)). \quad (\text{III.190})$$

In terms of e-folding N

$$\begin{aligned} \xi(\phi(N)) &= \frac{1}{8} \int^N dN_1 \frac{e^{N_1}}{H(N_1)^3} \tilde{W}(N_1), \\ V(\phi(N)) &= \frac{3}{\kappa^2} H(N)^2 - 3e^N H(N) \tilde{W}(N), \quad \tilde{W}(N) \equiv \frac{2}{\kappa^2} \int^N \frac{dN_1}{e^{N_1}} \dot{H}(N_1). \end{aligned} \quad (\text{III.191})$$

Now we identify ϕ with the e-folding N : $\phi = N$ instead of $\phi = t$. Then, if $V(\phi)$ and ξ are given by

$$\begin{aligned} V(\phi) &= \frac{3}{\kappa^2} h(\phi)^2 - 3h(\phi) e^\phi \tilde{U}(\phi), \\ \xi(\phi) &= \frac{1}{8} \int^\phi d\phi_1 \frac{e^{\phi_1}}{h(\phi_1)^3} \tilde{U}(\phi_1), \quad U(\phi) \equiv \frac{2}{\kappa^2} \int^\phi d\phi_1 e^{-\phi_1} h'(\phi_1), \end{aligned} \quad (\text{III.192})$$

we obtain the solution $H = h(N)$. This general scheme may now be applied to explicitly reconstruct the modified Gauss-Bonnet gravity by realizing any given FRW metric with a particular scale factor.

E. Scalar-Einstein-Gauss-Bonnet gravity

In the same way as above, let us reconstruct string-inspired scalar-Einstein-Gauss-Bonnet gravity in II C. The reconstructed theory realizes the solution with the arbitrary Hubble rate H . Combining (II.173) and (II.174) and deleting $V(\phi)$, we obtain

$$\begin{aligned} 0 &= \frac{2}{\kappa^2} \dot{H} + \dot{\phi}^2 - 8H^2 \frac{d^2\xi(\phi(t))}{dt^2} - 16H\dot{H} \frac{d\xi(\phi(t))}{dt} + 8H^3 \frac{d\xi(\phi(t))}{dt} \\ &= \frac{2}{\kappa^2} \dot{H} + \dot{\phi}^2 - 8a \frac{d}{dt} \left(\frac{H^2}{a} \frac{d\xi(\phi(t))}{dt} \right). \end{aligned} \quad (\text{III.193})$$

The following expressions of $\xi(\phi(t))$ and $V(\phi(t))$ may be obtained:

$$\begin{aligned}\xi(\phi(t)) &= \frac{1}{8} \int^t dt_1 \frac{a(t_1)}{H(t_1)^2} \int^{t_1} \frac{dt_2}{a(t_2)} \left(\frac{2}{\kappa^2} \dot{H}(t_2) + \dot{\phi}(t_2)^2 \right), \\ V(\phi(t)) &= \frac{3}{\kappa^2} H(t)^2 - \frac{1}{2} \dot{\phi}(t)^2 - 3a(t)H(t) \int^t \frac{dt_1}{a(t_1)} \left(\frac{2}{\kappa^2} \dot{H}(t_1) + \dot{\phi}(t_1)^2 \right).\end{aligned}\quad (\text{III.194})$$

Therefore, for the model where $V(\phi)$ and $\xi(\phi)$ are given by adequate functions $g(t)$ and $f(\phi)$ as follows,

$$\begin{aligned}V(\phi) &= \frac{3}{\kappa^2} g'(f(\phi))^2 - \frac{1}{2f'(\phi)^2} - 3g'(f(\phi)) e^{g(f(\phi))} \int^\phi d\phi_1 f'(\phi_1) e^{-g(f(\phi_1))} \left(\frac{2}{\kappa^2} g''(f(\phi_1)) + \frac{1}{f'(\phi_1)^2} \right), \\ \xi(\phi) &= \frac{1}{8} \int^\phi d\phi_1 \frac{f'(\phi_1) e^{g(f(\phi_1))}}{g'(\phi_1)^2} \int^{\phi_1} d\phi_2 f'(\phi_2) e^{-g(f(\phi_2))} \left(\frac{2}{\kappa^2} g''(f(\phi_2)) + \frac{1}{f'(\phi_2)^2} \right),\end{aligned}\quad (\text{III.195})$$

the solution of (II.173) and (II.174) is given by

$$\phi = f^{-1}(t) \quad (t = f(\phi)), \quad a = a_0 e^{g(t)} \quad (H = g'(t)). \quad (\text{III.196})$$

Similarly, the reconstruction of the model in terms of N , instead of the cosmological time t may be presented. Using the same steps that led to Eq.(III.194), we find

$$\begin{aligned}\xi(\phi(N)) &= \frac{1}{8} \int^N dN_1 \frac{e^{N_1}}{H(N_1)^3} \int^{N_1} \frac{dN_2}{e^{N_2}} \left(\frac{2}{\kappa^2} H'(N_2) + H(N_2) \phi'(N_2)^2 \right), \\ V(\phi(N)) &= \frac{3}{\kappa^2} H(N)^2 - \frac{1}{2} H(N)^2 \phi'(N)^2 - 3e^N H(N) \int^N \frac{dN_1}{e^{N_1}} \left(\frac{2}{\kappa^2} H'(N_1) + H(N_1) \phi'(N_1)^2 \right).\end{aligned}\quad (\text{III.197})$$

In terms of functions $h(\phi)$ and $\tilde{f}(\phi)$, if $V(\phi)$ and $\xi(\phi)$ are given by

$$\begin{aligned}V(\phi) &= \frac{3}{\kappa^2} h(\tilde{f}(\phi))^2 - \frac{h(\tilde{f}(\phi))^2}{2\tilde{f}'(\phi)^2} - 3h(\tilde{f}(\phi)) e^{\tilde{f}(\phi)} \int^\phi d\phi_1 \tilde{f}'(\phi_1) e^{-\tilde{f}(\phi_1)} \left(\frac{2}{\kappa^2} h'(\tilde{f}(\phi_1)) + \frac{h'(\tilde{f}(\phi_1))}{f'(\phi_1)^2} \right), \\ \xi(\phi) &= \frac{1}{8} \int^\phi d\phi_1 \frac{\tilde{f}'(\phi_1) e^{\tilde{f}(\phi_1)}}{h(\tilde{f}(\phi_1))^3} \int^{\phi_1} d\phi_2 f'(\phi_2) e^{-\tilde{f}(\phi_2)} \left(\frac{2}{\kappa^2} h'(\tilde{f}(\phi_2)) + \frac{h(\tilde{f}(\phi_2))}{\tilde{f}'(\phi_2)^2} \right),\end{aligned}\quad (\text{III.198})$$

one obtain the following solution:

$$\phi = \tilde{f}^{-1}(N) \quad (N = \tilde{f}(\phi)), \quad H = h(N). \quad (\text{III.199})$$

This may now be applied to obtain an explicit reconstruction of scalar-Gauss-Bonnet gravity, which admits any given FRW cosmology as a solution.

F. $F(R)$ Hořava-Lifshitz gravity

In this section, we consider the reconstruction of $F(R)$ Hořava-Lifshitz gravity. It is based on the results of ref. [79].

To start, let us analyze the simple model $F(\tilde{R}) = \tilde{R}$, of which the cosmology was studied in [90–112]. In such a case, the FRW equations are similar to those of general relativity,

$$H^2 = \frac{\kappa^2}{3(3\lambda - 1)} \rho_{\text{matter}}, \quad \dot{H} = -\frac{\kappa^2}{2(3\lambda - 1)} (\rho_{\text{matter}} + p_{\text{matter}}), \quad (\text{III.200})$$

where, for $\lambda \rightarrow 1$, the standard FRW equations are recovered. Note that the constant μ is now irrelevant because, as pointed out above, the term in front of μ in (II.274) becomes a total derivative. For such a theory, one has to introduce a dark energy source as well as an inflaton field, to reproduce the cosmic and inflationary accelerated epochs, respectively. It is also important to note that, for this case, the coupling constant is restricted to be $\lambda > 1/3$;

otherwise, Eqs. (III.200) become inconsistent. It seems reasonable to think that, for the current epoch, where $\tilde{R}$ has a small value, the IR limit of the theory is satisfied by $\lambda \sim 1$, but for the inflationary epoch, when the scalar curvature $\tilde{R}$ goes to infinity, λ will take a different value. It has been realized that for $\lambda = 1/3$, the theory develops an anisotropic Weyl invariance (see [84]), and thus it takes a special role, although for the present model this value is not allowed.

We now discuss some cosmological solutions of $F(\tilde{R})$ Hořava-Lifshitz gravity. The first FRW equation, given by (II.278) with $C = 0$, can be rewritten as a function of the number of e-foldings $N = \ln \frac{a}{a_0}$ instead of the usual time t . Here, we extend the reconstruction formalism to the Hořava-Lifshitz $F(R)$ gravity. Because $\frac{d}{dt} = H \frac{d}{dN}$ and $\frac{d^2}{dN^2} = H^2 \frac{d^2}{dN^2} + H \frac{dH}{dN} \frac{d}{dN}$, the first FRW equation (II.278) is rewritten as

$$0 = F(\tilde{R}) - 6 \left[(1 - 3\lambda + 3\mu)H^2 + \mu H H' \right] \frac{dF(\tilde{R})}{d\tilde{R}} + 36\mu H^2 \left[(1 - 3\lambda + 6\mu)H H' + \mu H'^2 + \mu H'' H \right] \frac{d^2 F(\tilde{R})}{d^2 \tilde{R}} - \kappa^2 \rho_{\text{matter}}, \quad (\text{III.201})$$

where the primes denote derivatives with respect to N . Thus, in this case, there is no restriction on the values of λ or μ . By using the energy conservation equation and assuming a perfect fluid with an EoS of $p_{\text{matter}} = w_{\text{matter}} \rho_{\text{matter}}$, the energy density yields

$$\rho_{\text{matter}} = \rho_0 a^{-3(1+w_{\text{matter}})} = \rho_0 a_0^{-3(1+w_{\text{matter}})} e^{-3(1+w_{\text{matter}})N}. \quad (\text{III.202})$$

As the Hubble parameter can be written as a function of the number of e-foldings, $H = H(N)$, the scalar curvature in (II.274) takes the form

$$\tilde{R} = 3(1 - 3\lambda + 6\mu)H^2 + 6\mu H H', \quad (\text{III.203})$$

which can be solved with respect to N as $N = N(\tilde{R})$, and one obtains an expression (III.201) that gives an equation of $F(\tilde{R})$ with the variable $\tilde{R}$. This can be simplified by writing $G(N) = H^2$ instead of the Hubble parameter. In such a case, the differential equation (III.201) yields

$$0 = F(\tilde{R}) - 6 \left[(1 - 3\lambda + 3\mu)G + \frac{\mu}{2}G' \right] \frac{dF(\tilde{R})}{d\tilde{R}} + 18\mu \left[(1 - 3\lambda + 6\mu)GG' + \mu GG'' \right] \frac{d^2 F(\tilde{R})}{d^2 \tilde{R}} - \kappa^2 \rho_0 a_0^{-3(1+w)} e^{-3(1+w)N}, \quad (\text{III.204})$$

and the scalar curvature is now written as $\tilde{R} = 3(1 - 3\lambda + 6\mu)G + 3\mu G'$. Thus, for a given cosmological solution $H^2 = G(N)$, one can resolve Eq. (III.204), and the $F(\tilde{R})$ that reproduces such solution is obtained.

As an example, we consider the Hubble parameter that reproduces the Λ CDM epoch. It is expressed as

$$H^2 = G(N) = H_0^2 + \frac{\kappa^2}{3} \rho_0 a^{-3} = H_0^2 + \frac{\kappa^2}{3} \rho_0 a_0^{-3} e^{-3N}, \quad (\text{III.205})$$

where H_0 and ρ_0 are constant. In general relativity, the terms on the r.h.s. of Eq. (III.205) correspond to an effective cosmological constant $\Lambda = 3H_0^2$ and to cold dark matter with an EoS parameter of $w = 0$. The corresponding $F(\tilde{R})$ can be reconstructed by following the same steps as described above. Using the expression for the scalar curvature $\tilde{R} = 3(1 - 3\lambda + 6\mu)G + 3\mu G'$, the relation between $\tilde{R}$ and N is obtained,

$$e^{-3N} = \frac{\tilde{R} - 3(1 - 3\lambda + 6\mu)H_0^2}{3k(1 + 3(\mu - \lambda))}, \quad (\text{III.206})$$

where $k = \frac{\kappa^2}{3} \rho_0 a_0^{-3}$. Then, substituting (III.205) and (III.206) into Eq. (III.204), one obtains the differential expression

$$\begin{aligned} 0 = & (1 - 3\lambda + 3\mu)F(\tilde{R}) - 2 \left(1 - 3\lambda + \frac{3}{2}\mu \right) \tilde{R} + 9\mu(1 - 3\lambda)H_0^2 \frac{dF(\tilde{R})}{d\tilde{R}} \\ & - 6\mu(\tilde{R} - 9\mu H_0^2)(\tilde{R} - 3H_0^2(1 - 3\lambda + 6\mu)) \frac{d^2 F(\tilde{R})}{d^2 \tilde{R}} - R - 3(1 - 3\lambda + 6\mu)H_0^2, \end{aligned} \quad (\text{III.207})$$

where, for simplicity, we have considered a pressureless fluid $w = 0$ in Eq. (III.204). Performing the change of variable $x = \frac{\tilde{R} - 9\mu H_0^2}{3H_0^2(1 + 3(\mu - \lambda))}$, the homogeneous part of Eq. (III.207) can be easily identified as a hypergeometric differential equation

$$0 = x(1 - x) \frac{d^2 F}{dx^2} + (\gamma - (\alpha + \beta + 1)x) \frac{dF}{dx} - \alpha\beta F, \quad (\text{III.208})$$

with the set of parameters (α, β, γ) being given by

$$\gamma = -\frac{1}{2}, \quad \alpha + \beta = \frac{1 - 3\lambda - \frac{3}{2}\mu}{3\mu}, \quad \alpha\beta = -\frac{1 + 3(\mu - \lambda)}{6\mu}. \quad (\text{III.209})$$

The complete solution of Eq. (III.208) is a Gauss hypergeometric function plus a linear term and a cosmological constant from the particular solution of Eq. (III.207), namely

$$F(\tilde{R}) = C_1 F(\alpha, \beta, \gamma; x) + C_2 x^{1-\gamma} F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; x) + \frac{1}{\kappa_1} \tilde{R} - 2\Lambda, \quad (\text{III.210})$$

where C_1 and C_2 are constants, $\kappa_1 = 3\lambda - 1$ and $\Lambda = -\frac{3H_0^2(1-3\lambda+9\mu)}{2(1-3\lambda+3\mu)}$. Note that for the metric (III.205), the convenient $F(R)$ gravity was reconstructed and studied in refs. [18, 120, 121, 138, 139]. In this case, the solution (III.210) behaves similarly to the convenient $F(R)$ theory, except that now the parameters of the theory depend on (λ, μ) , which are allowed to vary, as was noted above. Thus, the cosmic evolution described by the Hubble parameter (III.205) is reproduced by this class of theories.

One can also explore the solution (III.205) for a particular choice of the parameters $\mu = \lambda - \frac{1}{3}$, which plays a special role in the cosmological expansion as is shown below. In this case, the scalar $\tilde{R}$ turns out to be a constant, and Eq. (III.207), in the presence of a pressureless fluid, has the solution

$$F(\tilde{R}) = \frac{1}{\kappa_1} \tilde{R} - 2\Lambda, \quad \text{with} \quad \Lambda = \frac{3}{2}(3\lambda - 1)H_0^2. \quad (\text{III.211})$$

Thus, for this constraint on the parameters, the only consistent solution reduces to the Hořava linear theory with a cosmological constant.

As a further example, we consider the phantom (super)accelerating expansion. Such a system can be easily expressed in general relativity, where the FRW equation reads as $H^2 = \frac{\kappa^2}{3}\rho_{\text{ph}}$. Here, the subscript “ph” denotes the phantom nature of the fluid, which has an EoS given by $p_{\text{ph}} = w_{\text{ph}}\rho_{\text{ph}}$ with $w_{\text{ph}} < -1$. By using the energy conservation equation, the solution for the Hubble parameter turns out to be

$$H(t) = \frac{H_0}{t_s - t}, \quad (\text{III.212})$$

where $H_0 = -1/3(1 + w_{\text{ph}})$, and t_s is the Rip time. As in the above example, one can rewrite the Hubble parameter as a function of the number of e-foldings; this yields

$$G(N) = H^2(N) = H_0^2 e^{2N/H_0}. \quad (\text{III.213})$$

Then, by using the expression of the scalar curvature, the relation between $\tilde{R}$ and N is given by

$$e^{2N/H_0} = \frac{R}{H_0(AH_0 + 6\mu)}. \quad (\text{III.214})$$

By inserting (III.213) and (III.214) into the differential equation (III.204), we get (without matter)

$$\tilde{R}^2 \frac{d^2 F(\tilde{R})}{d\tilde{R}^2} + k_1 \tilde{R} \frac{dF(\tilde{R})}{d\tilde{R}} + k_0 F(\tilde{R}) = 0, \quad (\text{III.215})$$

where

$$k_1 = -\frac{(AH_0 + 6\mu)((AH_0 + 3\mu))}{6\mu(AH_0 + 12\mu)}, \quad k_0 = \frac{(AH_0 + 6\mu)^2}{12\mu(AH_0 + 12\mu)}. \quad (\text{III.216})$$

Eq. (III.215) is an Euler equation, whose solution is well-known

$$F(R) = C_1 R^{m_+} + C_2 R^{m_-}, \quad \text{where} \quad m_{\pm} = \frac{1 - k_1 \pm \sqrt{(k_1 - 1)^2 - 4k_0}}{2}. \quad (\text{III.217})$$

Such a theory belongs to the class of models with positive and negative powers of the curvature introduced in ref. [2] as is discussed in the second chapter. Thus, an $F(\tilde{R})$ Hořava-Lifshitz gravity has been reconstructed that reproduces the phantom dark epoch with no need of any exotic fluid. In the same way, any given cosmology may be reconstructed, including the unified inflation within dark energy era.

G. Non-minimal Yang-Mills theory

We now develop a reconstruction scheme of the Yang-Mills theory by starting with the following action

$$S = \int dx^4 \sqrt{-g} \left[\frac{R}{2\kappa^2} + \mathcal{F}(F_{\mu\nu}^a F^{a\mu\nu}) \right], \quad (\text{III.218})$$

where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c$, and $\mathcal{F}$ may be assumed to be a continuously differentiable function.

Here, we review the reconstruction of the model (III.218) based on ref. [140].

We concentrate here on the $SU(2)$ case where $f^{abc} = \epsilon^{abc}$. Because

$$\frac{\delta(F_{\mu\nu}^a F^{a\mu\nu})}{\delta A_\beta^h} = -4\epsilon^{hbc} A_\gamma^b F^{c\gamma\beta}, \quad (\text{III.219})$$

$$\frac{\delta(F_{\mu\nu}^a F^{a\mu\nu})}{\delta(\partial_\alpha A_\beta^h)} = 4F^{h\alpha\beta}, \quad (\text{III.220})$$

the equation of motion for the field potential A_μ^a turns into

$$\partial_\nu \left[\frac{\delta S}{\delta(\partial_\nu A_\mu^a)} \right] - \frac{\delta S}{\delta A_\mu^a} = 0, \quad (\text{III.221})$$

and, from here,

$$\partial_\nu [\sqrt{-g} \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) F^{a\nu\mu}] + \sqrt{-g} \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) \epsilon^{abc} A_\nu^b F^{c\nu\mu} = 0. \quad (\text{III.222})$$

The variation of (III.218) with respect to $g^{\mu\nu}$ yields the following equation of motion

$$\frac{1}{2\kappa^2} \left(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) - \frac{1}{2} g_{\mu\nu} \mathcal{F}(F_{\alpha\beta}^a F^{a\alpha\beta}) + 2\mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) F_{\mu\rho}^a F_\nu^{a\rho} = 0, \quad (\text{III.223})$$

where the following relation is used $\frac{\delta(F_{\rho\sigma}^a F^{a\rho\sigma})}{\delta g^{\mu\nu}} = 2\mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) F_{\mu\gamma}^a F_\nu^{a\gamma}$. Considering now a FRW universe (II.6), and the following Ansatz for the gauge field,

$$A_\mu^a = \begin{cases} \bar{\alpha} e^{\lambda(t)} \delta_\mu^a, & \mu = i, \\ 0, & \mu = 0, \end{cases} \quad (\text{III.224})$$

the $\mu = 0$ component of (III.221) becomes an identity, the $\mu = i$ component yields

$$\partial_t \left[a(t) \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) \dot{\lambda}(t) e^{\lambda(t)} \right] + \frac{2\bar{\alpha}^2}{a(t)} \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) e^{3\lambda(t)} = 0, \quad (\text{III.225})$$

and the (t, t) component of (III.223) is

$$\frac{3H^2(t)}{2\kappa^2} + \frac{1}{2} \mathcal{F}(F_{\alpha\beta}^a F^{a\alpha\beta}) + 2\bar{\alpha}^2 \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) \frac{\dot{\lambda}^2(t) e^{2\lambda(t)}}{a^2(t)} = 0, \quad (\text{III.226})$$

and the (i, i) component of (III.223) reduces to

$$\frac{1}{2\kappa^2} \left[2\dot{H}(t) + 3H^2(t) \right] - \frac{1}{2} \mathcal{F}(F_{\alpha\beta}^a F^{a\alpha\beta}) - 2\bar{\alpha}^2 \frac{e^{2\lambda(t)}}{a^2(t)} \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) \left[\dot{\lambda}^2(t) - 2\bar{\alpha}^2 \frac{e^{2\lambda(t)}}{a^2(t)} \right] = 0. \quad (\text{III.227})$$

Adding (III.226) to (III.227), one arrives at

$$a^4(t) \dot{H}(t) - 4\kappa^2 \bar{\alpha}^4 \mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) e^{4\lambda(t)} = 0, \quad (\text{III.228})$$

and then

$$\mathcal{F}'(F_{\alpha\beta}^a F^{a\alpha\beta}) = \frac{a^4(t) \dot{H}(t)}{4\kappa^2 \bar{\alpha}^4} e^{-4\lambda(t)}. \quad (\text{III.229})$$

Using (III.229), Eq. (III.225) reduces to:

$$2\bar{\alpha}^2 \dot{H}(t) e^{2\lambda(t)} + \left[5a(t) \dot{a}(t) \dot{H}(t) + a^2(t) \ddot{H}(t) \right] \dot{\lambda}(t) - 3a^2(t) \dot{H}(t) \dot{\lambda}^2(t) + a^2(t) \dot{H}(t) \ddot{\lambda}(t) = 0, \quad (\text{III.230})$$

which constitutes a differential equation for $\lambda(t)$. Thus, by using Eq. (III.224), once we have the function $\lambda(t)$, given by (III.230), we can obtain the corresponding Yang-Mills theory that reproduces the selected cosmology. The Ansatz considered above actually leads to a mathematical solution of the problem.

As an example, we consider the case of a power-law expansion: $a(t) = \left(\frac{t}{t_1}\right)^{h_1}$, where t_1 and h_1 are constant, and assuming $\lambda(t) = (h_1 - 1) \ln\left(\frac{t}{t_1}\right) + \lambda_1$, where λ_1 is again a constant, Eq. (III.230) reduces to the following algebraic equation

$$h_1(h_1 - 1) + \bar{\alpha}^2 t_1^2 e^{2\lambda_1} = 0, \quad (\text{III.231})$$

thus,

$$\lambda_1 = \frac{1}{2} \ln\left(\frac{h_1(1 - h_1)}{\bar{\alpha}^2 t_1^2}\right), \quad (\text{III.232})$$

and

$$\lambda(t) = (h_1 - 1) \ln\left(\frac{t}{t_1}\right) + \frac{1}{2} \ln\left(\frac{h_1(1 - h_1)}{\bar{\alpha}^2 t_1^2}\right). \quad (\text{III.233})$$

With the help of this reconstruction scheme, the function $\lambda(t)$ given by (III.233) is obtained. Using (III.224), one is able to reproduce the cosmology given by the power-law expansion: $a(t) = \left(\frac{t}{t_1}\right)^{h_1}$.

In summary, a cosmological reconstruction scheme was developed in the current chapter. It was presented in a very general form and was applied to a number of modified gravities introduced in the second chapter. Several examples of important cosmological epochs are considered. The cosmological reconstruction of modified gravities that realize such epochs as solutions was successfully made. Despite the fact that we limited ourselves to only a spatially flat FRW universe, this scheme may be easily used for the generation of other background solutions, i.e., an anisotropic universe, black holes, wormholes, etc.

IV. FINITE-TIME FUTURE SINGULARITIES IN MODIFIED GRAVITY

This chapter is devoted to the study of finite-time future singularities, which often occur in the current dark energy models. It is demonstrated that this is a quite general phenomenon; the effective phantom and some effective quintessence dark energy models including fluid, scalar, and modified models may show such behavior in the finite future. We start from a simple dark fluid universe and demonstrate that coupling with dark matter cannot cure such future singularities. The introduction of a specific form of modified gravity may naturally solve the problem of future singularities. The occurrence of singularities in modified gravity is considered in detail, as is the possible avoidance of the singularities via the addition of extra gravitational terms.

A. Future singularities in a dark, fluid universe: dark matter versus modified gravity

Let us reiterate several simple facts about coupled phantom dark energy. First, we assume that the energy density ρ_{DE} and the pressure p_{DE} of the dark matter satisfy the standard conservation law (II.144). Let the EoS parameter w_{DE} : $w_{\text{DE}} = p_{\text{DE}}/\rho_{\text{DE}} < -1$. The phantom dark energy solution of the FRW equation (II.3) (after replacing p_{matter} and ρ_{matter} with p_{DE} and ρ_{DE} , respectively) is given by

$$H = \frac{\frac{2}{3(1+w_{\text{DE}})}}{t_s - t}, \quad (\text{IV.1})$$

with the famous Big Rip singularity, at $t = t_s$.

Following ref. [47], the model where phantom dark energy couples with dark matter is considered. Then, the conservation law is modified as

$$\dot{\rho}_{\text{DE}} + 3H(1 + w_{\text{DE}})\rho_{\text{DE}} = -Q\rho_{\text{DE}}, \quad \dot{\rho}_{\text{DM}} + 3H\rho_{\text{DM}} = Q\rho_{\text{DE}}. \quad (\text{IV.2})$$

Here, Q is assumed to be a constant. The first equation can be solved as

$$\rho_{\text{DE}} = \rho_{\text{DE}(0)} a^{-3(1+w_{\text{DE}})} e^{-Qt}. \quad (\text{IV.3})$$

Here, $\rho_{\text{DE}(0)}$ is a constant of the integration. The second equation (IV.2) gives

$$\rho_{\text{DM}} = Q a(t)^{-3} \int^t dt' \rho_{\text{DE}(0)} a^{-3w_{\text{DE}}} e^{-Qt'}. \quad (\text{IV.4})$$

Thus, the second FRW equation is

$$-\frac{1}{\kappa^2} (2\dot{H} + 3H^2) = p_{\text{DE}} = w_{\text{DE}} \rho_{\text{DE}} = w \rho_{\text{DE}(0)} a^{-3(1+w_{\text{DE}})} e^{-Qt}. \quad (\text{IV.5})$$

An exact solution of (IV.5) is the de Sitter space

$$a(t) = a_0 e^{-\frac{Q}{3(1+w_{\text{DE}})}t} \quad \left(H = -\frac{Q}{3(1+w_{\text{DE}})} \right), \quad (\text{IV.6})$$

where a_0 is given by

$$-\frac{3}{\kappa^2} \left(\frac{Q}{3(1+w_{\text{DE}})} \right)^2 = w \rho_{\text{DE}(0)} a_0^{-3(1+w_{\text{DE}})}. \quad (\text{IV.7})$$

Note that because we are considering the phantom dark energy with $w_{\text{DE}} < -1$, H (IV.6) is positive and Eq. (IV.7) has a real solution.

Thus, the coupling of the dark matter with the phantom dark energy may give the de Sitter solution instead of the Big Rip solution (IV.1). This does not always mean that the Big Rip singularity can be avoided, but it opens a possibility that the universe could evolve into a de Sitter universe instead of the Big Rip.

Let us show that the above construction may help to solve the coincidence problem. One may identify the Hubble rate H with the present value of the Hubble rate $H = -\frac{Q}{3(1+w_{\text{DE}})} = H_0 \sim 10^{-33} \text{ eV}$. Eq. (IV.3) shows that the dark energy density is a constant

$$\rho_{\text{DE}} = \rho_{\text{DE}(0)} a_0^{-3(1+w_{\text{DE}})}. \quad (\text{IV.8})$$

Eq. (IV.4) can be integrated as

$$\rho_{\text{DM}} = \rho_{\text{DM}0} a^{-3} - (1+w_{\text{DE}}) \rho_{\text{DE}(0)} a_0^{-3(1+w_{\text{DE}})}. \quad (\text{IV.9})$$

Here, $\rho_{\text{DM}0}$ is a constant of integration, but the first FRW equation

$$\frac{3}{\kappa^2} H^2 = \rho_{\text{DM}} + \rho_{\text{DE}}, \quad (\text{IV.10})$$

shows $\rho_{\text{DM}0} = 0$ and, therefore, the dark matter density ρ_{DM} is also constant:

$$\rho_{\text{DM}} = -(1+w_{\text{DE}}) \rho_{\text{DE}(0)} a_0^{-3(1+w_{\text{DE}})} = -(1+w_{\text{DE}}) \rho_{\text{DE}}. \quad (\text{IV.11})$$

Then, if the de Sitter solution (IV.6) is an attractor, by choosing

$$-(1+w_{\text{DE}}) \sim \frac{1}{3}, \quad \text{i.e.} \quad w_{\text{DE}} \sim -\frac{4}{3}, \quad (\text{IV.12})$$

the coincidence problem could be solved. That is, even starting with a wide range of initial conditions, the solution approaches the de Sitter solution, in which the ratio of the dark energy and the dark matter is approximately 1/3 and is almost independent of the initial condition.

To check whether or not the de Sitter solution (IV.6) is an attractor, the following perturbation is considered

$$a = a_0 e^{-\frac{Q}{3(1+w_{\text{DE}})}t + \Delta(t)}. \quad (\text{IV.13})$$

Here, $\Delta(t)$ is assumed to be small. The second FRW equation (IV.5) gives

$$-\frac{1}{\kappa^2} \left(2\ddot{\Delta} - \frac{2Q}{(1+w_{\text{DE}})} \dot{\Delta} \right) = -3(1+w_{\text{DE}}) w \rho_{\text{DE}(0)} a_0^{-3(1+w_{\text{DE}})} \Delta = \frac{3(1+w_{\text{DE}})}{\kappa^2} \left(\frac{Q}{3(1+w_{\text{DE}})} \right)^2 \Delta, \quad (\text{IV.14})$$

which is a very simple linear differential equation with a constant coefficient. Here, Eq. (IV.7) is used. Assuming $\Delta = e^{\lambda t}$, it follows that

$$0 = \lambda^2 - \frac{Q}{1 + w_{\text{DE}}} \lambda + 3(1 + w_{\text{DE}}) \left(\frac{Q}{3(1 + w_{\text{DE}})} \right)^2, \quad (\text{IV.15})$$

that is,

$$\lambda = \lambda_{\pm} \equiv \frac{Q}{2(1 + w_{\text{DE}})} \pm \frac{1}{2} \left\{ \left(\frac{Q}{1 + w_{\text{DE}}} \right)^2 - \frac{4(1 + w_{\text{DE}})}{3} \left(\frac{Q}{(1 + w_{\text{DE}})} \right)^2 \right\}^{\frac{1}{2}}. \quad (\text{IV.16})$$

Then, $\lambda_- < 0$ but $\lambda_+ > 0$ and, therefore, de Sitter solution (IV.6) is not stable. Because $\lambda_- < 0$, however, the solution is saddle point and, therefore, with the appropriate initial condition, there is a solution that approaches the saddle point de Sitter solution (IV.6). Although we need to study the global structure of the space of the solutions, such an initial condition could correspond to the direction toward λ_- . Thus, generally speaking, the coupling of the dark energy with dark matter does not prohibit the existence of the (Big Rip) singular solution. If the singularity corresponds to the stable solution, because the de Sitter space is not completely stable, the solution will finally evolve into the singular solution. Thus, the appropriate choice of initial conditions may help to realize the non-singular de Sitter cosmology, which also solves the coincidence problem. At least, one may expect that the future singularity would occur at more later times compared with the case in which the dark matter does not couple with dark energy.

Let us consider the simple example of a perfect fluid with the following EoS [141]:

$$p = -\rho + A\rho^{\alpha}, \quad (\text{IV.17})$$

with a constant A and α . The EoS (IV.17) is a special case of (II.339). In the spatially flat FRW space-time (II.6), the Hubble rate is found to be

$$H = \begin{cases} \frac{\frac{3}{2}A}{t_s - t}, & \text{when } \alpha = 1, \quad A > 0 \\ \frac{-\frac{3}{2}A}{t_s - t}, & \text{when } \alpha = 1, \quad A < 0 \\ B e^{-\frac{\sqrt{3}A}{2}t}, & \text{when } \alpha = \frac{1}{2}, \quad A < 0 \\ C t^{1/(1-2\alpha)}, & \text{when } \alpha \neq 1, \frac{1}{2} \\ \text{or } \tilde{C} (t_s - t)^{1/(1-2\alpha)} & \text{when } \alpha \neq 1, \frac{1}{2} \end{cases} \quad (\text{IV.18})$$

Here, B , C , and $\tilde{C}$ are positive constants. Now, one can describe the future, finite-time singularities of the universe filled with the above dark fluid for different choices of theory parameters (see ref. [122]). When $\alpha < 0$, a Type II or sudden future singularity occurs. When $0 < \alpha < 1/2$ and $1/(1 - 2\alpha)$ is not an integer, a Type IV singularity occurs. When $\alpha = 0$, there is no singularity. When $1/2 < \alpha < 1$ or $\alpha = 1$ and $A < 0$, a Type I or Big Rip-type singularity occurs (see also the last section of the second chapter). When $\alpha > 1$, a Type III singularity occurs (for the classification of all four types of future singularities, see [34]).

In the case of a Type II singularity, where $\alpha < 0$, H vanishes as $H \sim (t_s - t)^{1/(1-2\alpha)}$ when $t \rightarrow t_s$ and, therefore, ρ vanishes as follows from the FRW equation. Then, near the singularity, the EoS (II.339) is reduced to

$$p \sim A\rho^{\alpha}. \quad (\text{IV.19})$$

On the other hand, in the case of a Type I singularity, where $1/2 < \alpha < 1$ or $\alpha = 1$ and $A < 0$, H and, therefore, ρ diverge when $t \rightarrow t_s$. Then, the EoS (II.339) reduces to

$$p \sim -\rho \text{ or } p \sim -(1 - A)\rho. \quad (\text{IV.20})$$

In the case of a Type III singularity, where $\alpha > 1$, H and ρ diverge when $t \rightarrow t_s$ and, therefore, the EoS (II.339) reduces to

$$p \sim A\rho^{\alpha}. \quad (\text{IV.21})$$

Let us now consider the dark energy (IV.17) coupled with the dark matter as in (IV.2). The conservation law is given by

$$\dot{\rho}_{\text{DE}} + 3H A \rho_{\text{DE}}^{\alpha} = -Q\rho_{\text{DE}}, \quad \dot{\rho}_{\text{DM}} + 3H \rho_{\text{DM}} = Q\rho_{\text{DE}}. \quad (\text{IV.22})$$

The solution of (IV.22) is

$$\rho_{\text{DE}}(t) = e^{-Qt} \left(-3A(1-\alpha) \int^t dt' H(t') e^{(1-\alpha)Qt'} \right)^{\frac{1}{1-\alpha}}, \quad \rho_{\text{DM}} = Qa(t)^{-3} \int^t dt' a(t')^3 \rho_{\text{DE}}(t'). \quad (\text{IV.23})$$

If $A < 0$ (and $Q > 0$), there is a de Sitter solution $H = H_0$ with a constant H_0

$$\frac{3}{\kappa^2} H_0^2 = \left(1 + \frac{Q}{3H_0} \right) \left(-\frac{3AH_0}{Q} \right)^{\frac{1}{1-\alpha}}, \quad (\text{IV.24})$$

and ρ_{DE} and ρ_{DM} are constants

$$\rho_{\text{DE}} = \left(-\frac{3AH_0}{Q} \right)^{\frac{1}{1-\alpha}}, \quad \rho_{\text{DM}} = \frac{Q}{3H_0} \left(-\frac{3AH_0}{Q} \right)^{\frac{1}{1-\alpha}}. \quad (\text{IV.25})$$

This demonstrates that if $H_0 \sim Q$, we find $\rho_{\text{DM}}/\rho_{\text{DE}} \sim 1/3$, and the coincidence problem may be solved. If $H_0 = Q$, Eq. (IV.24) determines the value of A :

$$A = -\frac{1}{3} \left(\frac{9}{4\kappa^2} H_0^2 \right)^{1-\alpha}. \quad (\text{IV.26})$$

One can now investigate the (in)stability of the de Sitter solution $H = H_0$ by putting $H = H_0 + \delta H$. The perturbation of the energy density is

$$\delta\rho_{\text{DE}} = -3A \left(-\frac{3AH_0}{Q} \right)^{\frac{\alpha}{1-\alpha}} e^{-(1-\alpha)Qt} \int^t dt' \delta H(t') e^{(1-\alpha)Qt'}. \quad (\text{IV.27})$$

The second FRW equation becomes

$$-\frac{1}{\kappa^2} (\delta\dot{H} + 6H_0\delta H) e^{(1-\alpha)Qt'} = 3A \left(1 + \frac{\alpha Q}{3H_0} \right) \left(-\frac{3AH_0}{Q} \right)^{\frac{\alpha}{1-\alpha}} \int^t dt' \delta H(t') e^{(1-\alpha)Qt'}. \quad (\text{IV.28})$$

By differentiating both sides of (IV.28), one gets

$$\begin{aligned} 0 &= \delta\ddot{H} + \{6H_0 + (1-\alpha)Q\} \delta\dot{H} + \left\{ 6H_0(1-\alpha)Q + 3A\kappa^2 \left(1 + \frac{\alpha Q}{3H_0} \right) \left(-\frac{3AH_0}{Q} \right)^{\frac{\alpha}{1-\alpha}} \right\} \delta H \\ &= \delta\ddot{H} + \{6H_0 + (1-\alpha)Q\} \delta\dot{H} + 3H_0(1-2\alpha)Q\delta H. \end{aligned} \quad (\text{IV.29})$$

In the second equality, Eq. (IV.24) is used. Assuming $\delta H \propto e^{\lambda t}$, one gets

$$0 = \lambda^2 + \{6H_0 + (1-\alpha)Q\} \lambda + 3H_0(1-2\alpha)Q, \quad (\text{IV.30})$$

whose solution is given by

$$\lambda = \lambda_{\pm} \equiv -\frac{6H_0 + (1-\alpha)Q}{2} \pm \frac{1}{2} \left\{ \{6H_0 + (1-\alpha)Q\}^2 - 12H_0(1-2\alpha)Q \right\}^{\frac{1}{2}}. \quad (\text{IV.31})$$

Because $H_0, Q > 0$, if

$$\alpha < \frac{1}{2}, \quad (\text{IV.32})$$

both $\lambda_{\pm}$ are real and negative if

$$D = (6H_0 + (1-\alpha)Q)^2 - 12H_0(1-2\alpha)Q > 0, \quad (\text{IV.33})$$

or complex, but the real part is negative if

$$D = (6H_0 + (1-\alpha)Q)^2 - 12H_0(1-2\alpha)Q < 0. \quad (\text{IV.34})$$

Thus, as long as $\alpha < \frac{1}{2}$ (IV.32), the de Sitter solution is stable and, therefore, the singularity can be resolved. On the other hand, if

$$6H_0 + (1 - \alpha)Q > 0, \quad 3H_0(1 - 2\alpha)Q < 0, \quad (\text{IV.35})$$

that is,

$$1 + \frac{6H_0}{Q} > \alpha > \frac{1}{2}, \quad (\text{IV.36})$$

we find $\lambda_+ > 0$ and $\lambda_- < 0$, as at the beginning of this section. If

$$6H_0 + (1 - \alpha)Q < 0, \quad 3H_0(1 - 2\alpha)Q < 0, \quad (\text{IV.37})$$

that is,

$$\alpha > 1 + \frac{6H_0}{Q}, \quad (\text{IV.38})$$

we find $\lambda_{\pm} > 0$ and the de Sitter solution is completely unstable.

Thus, the Type II singularity, where $\alpha < 0$, and the Type IV singularity, where $0 < \alpha < 1/2$ and $1/(1 - 2\alpha)$ is not an integer, can be resolved by the coupling of the dark energy with the dark matter. As mentioned above, even with a coupling of the dark matter with the dark energy, there could be a singular solution. The de Sitter solution with $\alpha < \frac{1}{2}$ is, however, at least a local minimum. Then, if the universe started with an appropriate initial condition, the universe would evolve into the de Sitter universe (asymptotically evolving into a de Sitter universe). A Type I (Big Rip) singularity, where $1/2 < \alpha < 1$, and a Type III singularity, where $\alpha > 1$, could not be removed by the coupling of the dark matter with the dark energy. Even if a solution goes near the de Sitter point, the solution could evolve into the singular solution if the singular solution is stable. Nevertheless, the ability to avoid some future singularities due to the coupling of the dark energy with the dark matter looks quite promising.

Because the coupling of the dark energy with the dark matter does not always remove the singularity, we now consider what kind of fluid could cure the future singularity. In the case of the Big Rip singularity, for example, the energy density of the dark energy diverges like $\rho_{\text{DE}} \sim 1/(t_s - t)^2 \sim R$ when $t \rightarrow t_s$. Here, R is the scalar curvature. Then, one becomes interested in a fluid with a positive pressure (and a positive energy density) that grows more rapidly than the dark energy pressure. There is no such fluid with a constant EoS parameter. However, one can consider a pressure that is proportional to a power of the curvature; for example,

$$p_{\text{fluid}} \propto R^{1+\epsilon}, \quad (\text{IV.39})$$

with $\epsilon > 0$. Then, the total EoS parameter becomes greater than -1 for a large curvature, and the Big Rip does not occur.

This kind of inhomogeneous effective fluid [118] could be realized by quantum effects (for instance, taking account of conformal anomaly) or by modified gravity as was explained in the second chapter. As an example, we consider $F(R)$ gravity, where $F(R) = R + f(R)$ behaves as $f(R) \propto R^m$. In the case of $m = 2$, the power-law solution, like the Big Rip singularity is prohibited. It can happen that the (asymptotic) de Sitter solution occurs, instead of the power-law solution and, therefore, the singularity would be resolved. Therefore, the introduction of the special form of $f(R)$ (precisely, the R^2 term) prevents the future singularity in the universal way. Note that as with other dark energies, the modified gravity itself (see next section) may lead to all four possible future singularities, which may again be resolved by the addition of the R^2 term [37]. Thus, curing all future singularities of fluid dark energy may be done via modifying the gravity. This seems to be quite a fundamental reason to start from modified gravity as a gravitational alternative for dark energy.

B. Finite-time singularities in $F(R)$ gravity

Let us show that $F(R)$ gravity generates all four known types of finite-time singularities. Moreover, the resolution of future singularities may be done again by adding the R^2 term [37, 40]. This section is based on results of refs. [36, 39].

As the first example, we consider the case of the Big Rip singularity (IV.1). To find the $F(R)$ gravity that generates the Big Rip-type singularity, it is convenient to use the reconstruction scheme developed in subsection III C 1. Let $h_0 \equiv \frac{2}{3(1+w_{\text{DE}})}$. In the case of (IV.1), the general solution of (III.105) when the matter contribution is neglected, $\rho_{i0} = 0$, is given by

$$P(\phi) = P_+(t_s - \phi)^{\alpha_+} + P_-(t_s - \phi)^{\alpha_-}, \quad \alpha_{\pm} \equiv \frac{-h_0 + 1 \pm \sqrt{h_0^2 - 10h_0 + 1}}{2}, \quad (\text{IV.40})$$

when $h_0 > 5 + 2\sqrt{6}$ or $h_0 < 5 - 2\sqrt{6}$ and

$$P(\phi) = (t_s - \phi)^{-(h_0+1)/2} \left(\hat{A} \cos \left((t_s - \phi) \ln \frac{-h_0^2 + 10h_0 - 1}{2} \right) + \hat{B} \sin \left((t_s - \phi) \ln \frac{-h_0^2 + 10h_0 - 1}{2} \right) \right), \quad (\text{IV.41})$$

when $5 + 2\sqrt{6} > h_0 > 5 - 2\sqrt{6}$. One finds the form of $F(R)$ when R is large as

$$F(R) \propto R^{1-\alpha-/2}, \quad (\text{IV.42})$$

for the $h_0 > 5 + 2\sqrt{6}$ or $h_0 < 5 - 2\sqrt{6}$ case and

$$F(R) \propto R^{(h_0+1)/4} \times (\text{oscillating parts}), \quad (\text{IV.43})$$

for the $5 + 2\sqrt{6} > h_0 > 5 - 2\sqrt{6}$ case.

Let us investigate a more general singularity

$$H \sim h_0 (t_s - t)^{-\beta}. \quad (\text{IV.44})$$

Here, h_0 and β are constants, h_0 is assumed to be positive, and $t < t_s$, as it should be for an expanding universe. Even for a non-integer $\beta < 0$, some derivative of H and, therefore, the curvature become singular. Assume $\beta \neq 1$. Furthermore, because $\beta = 0$ corresponds to the de Sitter space, it is proposed that $\beta \neq 0$. When $\beta > 1$, the scalar curvature R behaves as

$$R \sim 12H^2 \sim 12h_0^2 (t_s - t)^{-2\beta}. \quad (\text{IV.45})$$

On the other hand, when $\beta < 1$, the scalar curvature R behaves as

$$R \sim 6\dot{H} \sim 6h_0\beta (t_s - t)^{-\beta-1}. \quad (\text{IV.46})$$

If we write

$$P(\phi) = e^{-h_0(t_s - \phi)^{-\beta+1}/2(1-\beta)} S(\phi), \quad (\text{IV.47})$$

Eq. (III.105), without the matter contribution, has the following Schrödinger equation like form:

$$0 = \frac{d^2 S(\phi)}{d\phi^2} + \left(\frac{5\beta h_0}{2} (t_s - \phi)^{-\beta-1} - \frac{h_0^2}{4} (t_s - \phi)^{-2\beta} \right) S. \quad (\text{IV.48})$$

When $\phi = t \rightarrow t_s$, in the case that $\beta > 1$, one finds

$$\left| \frac{5\beta h_0}{2} (t_s - \phi)^{-\beta-1} \right| \ll \left| \frac{h_0^2}{4} (t_s - \phi)^{-2\beta} \right|. \quad (\text{IV.49})$$

On the other hand, in the case that $\beta < 1$, it follows that

$$\left| \frac{5\beta h_0}{2} (t_s - \phi)^{-\beta-1} \right| \gg \left| \frac{h_0^2}{4} (t_s - \phi)^{-2\beta} \right|. \quad (\text{IV.50})$$

In either case, Eq. (IV.48) reduces to the following form:

$$0 = \frac{d^2 S(\phi)}{d\phi^2} - V_0 (t_s - \phi)^{-\alpha} S, \quad (\text{IV.51})$$

when $\phi = t \rightarrow t_s$. Here

$$\begin{aligned} V_0 &= -\frac{5\beta h_0}{2}, \quad \alpha = \beta + 1 \quad \text{when } \beta < 1 \\ V_0 &= \frac{h_0^2}{4}, \quad \alpha = 2\beta \quad \text{when } \beta > 1 \end{aligned} \quad (\text{IV.52})$$

With a further redefinition

$$y \equiv (t_s - \phi)^{1-\alpha/2}, \quad S = y^{(\alpha/4)(1-\alpha/2)^{-1}} \varphi, \quad (\text{IV.53})$$

Eq. (IV.51) has the following form:

$$0 = \frac{d^2\varphi}{dy^2} - \left\{ \left(\frac{\alpha^2}{16} - \frac{\alpha}{4} \right) \frac{1}{y^2} + \frac{4V_0}{(2-\alpha)^2} \right\} \varphi. \quad (\text{IV.54})$$

Note that $y \rightarrow 0$ when $\phi \rightarrow t_s$ if $1 - \alpha/2 > 0$, but $y \rightarrow \infty$ when $\phi \rightarrow t_s$ if $1 - \alpha/2 < 0$. Then, if $1 - \alpha/2 > 0$, Eq. (IV.54) reduces to the following form when $\phi \rightarrow t_s$:

$$0 = \frac{d^2\varphi}{dy^2} - \left(\frac{\alpha^2}{16} - \frac{\alpha}{4} \right) \frac{1}{y^2} \varphi, \quad (\text{IV.55})$$

whose general solution is given by

$$\varphi = Ay^{\alpha/4-1} + By^{-\alpha/4}. \quad (\text{IV.56})$$

Here, A and B are constants of integration. On the other hand, if $1 - \alpha/2 < 0$, Eq. (IV.54) reduces to the following form when $\phi \rightarrow t_s$:

$$0 = \frac{d^2\varphi}{dy^2} + \frac{4V_0}{(\alpha-2)^2} \varphi. \quad (\text{IV.57})$$

When $V_0 > 0$, the general solution of (IV.57) is given by

$$y = \tilde{A} \cos(\omega y) + \tilde{B} \sin(\omega y), \quad \omega \equiv \frac{2\sqrt{V_0}}{\alpha-2}. \quad (\text{IV.58})$$

Here, $\tilde{A}$ and $\tilde{B}$ are constants of integration. On the other hand, if $V_0 < 0$, the general solution has the following form

$$y = \hat{A} e^{\hat{\omega} y} + \hat{B} e^{-\hat{\omega} y}, \quad \hat{\omega} \equiv \frac{2\sqrt{-V_0}}{\alpha-2}. \quad (\text{IV.59})$$

From the above analysis, one may obtain the asymptotic solution for P when $\phi \rightarrow t_s$.

- $\beta > 1$ case: From (IV.52), $\alpha = 2\beta > 2$ and, therefore, $1 - \alpha/2 = 1 - \beta < 0$, which corresponds to (IV.57). Because we also find $V_0 > 0$, the solution is given by (IV.58). Combining (IV.44), (IV.47), (IV.53), and (IV.58), we find the following asymptotic expression of $P(\phi)$:

$$P(\phi) \sim e^{(h_0/2(\beta-1))(t_s-\phi)^{-\beta+1}} (t_s - \phi)^{\beta/2} \left(\tilde{A} \cos\left(\omega (t_s - \phi)^{-\beta+1}\right) + \tilde{B} \sin\left(\omega (t_s - \phi)^{-\beta+1}\right) \right),$$

$$\omega \equiv \frac{h_0}{2(\beta-1)}. \quad (\text{IV.60})$$

When $\phi \rightarrow t_s$, $P(\phi)$ tends to vanish very rapidly. For large R , $F(R)$ looks like (at large R)

$$F(R) \propto e^{(h_0/2(\beta-1))\left(\frac{R}{12h_0}\right)^{(\beta-1)/2\beta}} R^{-1/4} \times (\text{oscillating part}). \quad (\text{IV.61})$$

- $1 > \beta > 0$ case: From (IV.52), it follows that $\alpha = \beta + 1$ and, therefore, $1 - \alpha/2 = 1/2 - \beta/2 > 0$, which corresponds to (IV.55). Because

$$\frac{\alpha}{4} - 1 - \left(-\frac{\alpha}{4}\right) = \beta > 0, \quad (\text{IV.62})$$

the second term in (IV.56) dominates when $\phi \rightarrow t_s$ if $B \neq 0$. Then, by combining (IV.44), (IV.47), (IV.53), and (IV.56), the following asymptotic expression of $P(\phi)$ is obtained:

$$P(\phi) \sim B e^{-(h_0/2(1-\beta))(t_s-\phi)^{1-\beta}} (t_s - \phi)^{(\beta+1)/8}. \quad (\text{IV.63})$$

Therefore,

$$F(R) \sim e^{-(h_0/2(1-\beta))(-6\beta h_0 R)^{(\beta-1)/(\beta+1)}} R^{7/8}. \quad (\text{IV.64})$$

Eq. (IV.46) shows that when $\phi = t \rightarrow t_s$, $R \rightarrow \infty$ in the case that $\beta > -1$, but $R \rightarrow 0$ in the case that $\beta < -1$.

- $\beta < 0$ case: As in $1 > \beta > 0$ case, one gets $\alpha = \beta + 1$ and, therefore, $1 - \alpha/2 > 0$; however, because

$$\frac{\alpha}{4} - 1 - \left(-\frac{\alpha}{4}\right) = \beta < 0, \quad (\text{IV.65})$$

the first term in (IV.56) dominates when $\phi \rightarrow t_s$ if $A \neq 0$. Thus, the asymptotic expression of $P(\phi)$ is as follows:

$$P(\phi) \sim A e^{-(h_0/2(1-\beta))(t_s-\phi)^{1-\beta}} (t_s - \phi)^{-(\beta^2-6\beta+1)/8}. \quad (\text{IV.66})$$

Then, $F(R)$ is given by

$$F(R) \sim (-6h_0\beta R)^{(\beta^2+2\beta+9)/8(\beta+1)} e^{-(h_0/2(1-\beta))(-6h_0\beta R)^{(\beta-1)/(\beta+1)}}. \quad (\text{IV.67})$$

Note that $-6h_0\beta R > 0$ when $h_0, R > 0$.

When $\beta > 1$ in (IV.44), R behaves as in (IV.45), and when $\beta < 1$, the scalar curvature R behaves as in (IV.46). Conversely, when R behaves as

$$R \sim 6\dot{H} \sim R_0 (\beta + 1) (t_s - t)^{-\gamma}, \quad (\text{IV.68})$$

if $\gamma > 2$, which corresponds to $\beta = \gamma/2 > 1$, H behaves as

$$H \sim \sqrt{\frac{R_0}{12}} (t_s - t)^{-\gamma/2}. \quad (\text{IV.69})$$

On the other hand, if $2 > \gamma > 1$, which corresponds to $1 > \beta = \gamma - 1 > 0$, H is given by

$$H \sim \frac{R_0}{6(\gamma - 1)} (t_s - t)^{-\gamma+1}, \quad (\text{IV.70})$$

and if $\gamma < 1$, which corresponds to $\beta = \gamma - 1 < 0$, one obtains

$$H \sim H_0 + \frac{R_0}{6(\gamma - 1)} (t_s - t)^{-\gamma+1}. \quad (\text{IV.71})$$

Here, H_0 is an arbitrary constant, which is chosen to vanish in (IV.44). Then, because $H = \dot{a}(t)/a(t)$, if $\gamma > 2$, we find

$$a(t) \propto \exp\left(\left(\frac{2}{\gamma} - 1\right) \sqrt{\frac{R_0}{12}} (t_s - t)^{-\gamma/2+1}\right), \quad (\text{IV.72})$$

when $2 > \gamma > 1$, $a(t)$ behaves as

$$a(t) \propto \exp\left(\frac{R_0}{6\gamma(\gamma - 1)} (t_s - t)^{-\gamma}\right), \quad (\text{IV.73})$$

and if $\gamma < 1$,

$$a(t) \propto \exp\left(H_0 t + \frac{R_0}{6\gamma(\gamma - 1)} (t_s - t)^{-\gamma}\right). \quad (\text{IV.74})$$

In any case, there appears a sudden future singularity at $t = t_s$.

Because the second term in (IV.71) is smaller than the first term, the asymptotic solution of Eq.(III.105) looks as follows:

$$P \sim P_0 \left(1 + \frac{2h_0}{1-\beta} (t_s - \phi)^{1-\beta}\right), \quad (\text{IV.75})$$

with a constant P_0 , which gives

$$F(R) \sim F_0 R + F_1 R^{2\beta/(\beta+1)}. \quad (\text{IV.76})$$

Here, F_0 and F_1 are constants. When $0 > \beta > -1$, we find that $2\beta/(\beta+1) < 0$. On the other hand, when $\beta < -1$, it follows $2\beta/(\beta+1) > 2$. As is seen from (IV.42), $F(R)$ generates the Big Rip singularity when R is large. Thus, even if R is small, the $F(R)$ generates a singularity where higher derivatives of H diverge.

Let us also investigate how the effective EoS parameter w_{eff} for (IV.71) behaves when $t \sim t_s$. In Einstein gravity, the FRW equations are given by

$$\frac{3}{\kappa^2}H^2 = \rho, \quad -\frac{1}{\kappa^2}(2\dot{H} + 3H^2) = p. \quad (\text{IV.77})$$

As was discussed in section A of the second chapter, for $F(R)$ gravity, we may introduce the EoS parameter w_{eff} by

$$w_{\text{eff}} = -1 - \frac{2\dot{H}}{3H^2}. \quad (\text{IV.78})$$

Then, if $\beta = \gamma/2 > 1$, it follows that

$$w_{\text{eff}} \sim -1 - \frac{2\beta}{3h_0}(t_s - t)^{-1+\beta} \rightarrow -1, \quad (\text{IV.79})$$

when $t \rightarrow t_s$. If $1 > \beta = \gamma - 1 > 0$, we find

$$w_{\text{eff}} \sim -\frac{2\beta}{3h_0}(t_s - t)^{-1+\beta} \rightarrow -\infty. \quad (\text{IV.80})$$

Finally, if $\beta = \gamma - 1 < 0$, one gets the expression (IV.80) when $H_0 = 0$. When $H_0 \neq 0$, on the other hand, we obtain

$$w_{\text{eff}} \sim -1 - \frac{2\beta h_0}{3H_0^2}(t_s - t)^{-1-\beta}, \quad (\text{IV.81})$$

when H_0 vanishes. Then, if $-1 < \beta < 0$, $w_{\text{eff}} \rightarrow +\infty$ when $t \rightarrow t_s$. On the other hand, if $\beta < -1$, $w_{\text{eff}} \rightarrow -1$.

Eq. (IV.77) also shows that, even for $F(R)$ gravity, we may define the effective energy density ρ_{eff} and the effective pressure p_{eff} by

$$\rho_{\text{eff}} \equiv \frac{3}{\kappa^2}H^2, \quad p_{\text{eff}} \equiv -\frac{1}{\kappa^2}(2\dot{H} + 3H^2). \quad (\text{IV.82})$$

We now assume that H behaves as (IV.44). Then, if $\beta > 1$, when $t \rightarrow t_s$, $a \sim \exp(h_0(t_s - t)^{1-\beta}/(\beta-1)) \rightarrow \infty$ and $\rho_{\text{eff}}, |p_{\text{eff}}| \rightarrow \infty$. If $\beta = 1$, we find $a \sim (t_s - t)^{-h_0} \rightarrow \infty$ and $\rho_{\text{eff}}, |p_{\text{eff}}| \rightarrow \infty$. If $0 < \beta < 1$, a goes to a constant but $\rho, |p| \rightarrow \infty$. If $-1 < \beta < 0$, we find that a and ρ vanish but $|p_{\text{eff}}| \rightarrow \infty$. When $\beta < 0$, instead of (IV.44), as in (IV.70), one may assume

$$H \sim H_0 + h_0(t_s - t)^{-\beta}. \quad (\text{IV.83})$$

Thus, ρ_{eff} has a finite value $3H_0^2/\kappa^2$ in the limit $t \rightarrow t_s$ when $-1 < \beta < 0$. If $\beta < -1$ but β is not any integer, a is finite and ρ_{eff} and p_{eff} vanish if $H_0 = 0$ or both of ρ_{eff} and p_{eff} are finite if $H_0 \neq 0$ but higher derivatives of H diverge.

In the classification of the singularities in [34], Type I corresponds to the $\beta > 1$ or $\beta = 1$ case, Type II to the $-1 < \beta < 0$ case, Type III to the $0 < \beta < 1$ case, and Type IV to the $\beta < -1$ case (where β is not an integer). Thus, we have constructed $F(R)$ gravity examples that show all four types of the above finite-time singularities. Viable $F(R)$ gravities (see second chapter) may also show (soft) future singular behavior. It is interesting to mention that future singularities manifest themselves in different presentations of $F(R)$ gravity. For instance, if the scalar-tensor version shows the Big Rip-type singularity, then its $F(R)$ counterpart becomes a complex theory precisely at the rip time [142].

Near the future singularity at $t = t_s$, the curvature becomes large in general. As a result, near the singularity, the quantum fields/quantum gravity effects become very important again. All classical considerations are not valid, and all speculations about a future cosmic doomsday cannot restrict the classical theory structure because quantum effects can stop (or shift) the future singularity. Moreover, the quantum corrections usually contain the curvature powers, which become important near the singularity. Thus, any claim about the appearance of the effective phantom/quintessence phase in modified gravity that subsequently enters the future singularity is not justified without quantum effects occurring near the singularity. One may include the massless quantum effects by taking into account the conformal anomaly contribution as a back-reaction near the singularity. The conformal anomaly T_A has the following well-known form:

$$T_A = b\left(\mathcal{F} + \frac{2}{3}\square R\right) + b'\mathcal{G} + b''\square R, \quad (\text{IV.84})$$

where $\mathcal{F}$ is the square of the 4D Weyl tensor, and $\mathcal{G}$ is the Gauss-Bonnet invariant, which are given by

$$\mathcal{F} = \frac{1}{3}R^2 - 2R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \quad \mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}. \quad (\text{IV.85})$$

In general, with N scalars, $N_{1/2}$ spinors, N_1 vector fields, N_2 ($= 0$ or 1) gravitons and N_{HD} higher-derivative conformal scalars, b and b' are given by

$$b = \frac{N + 6N_{1/2} + 12N_1 + 611N_2 - 8N_{\text{HD}}}{120(4\pi)^2}, \quad b' = -\frac{N + 11N_{1/2} + 62N_1 + 1411N_2 - 28N_{\text{HD}}}{360(4\pi)^2}. \quad (\text{IV.86})$$

As is seen, $b > 0$ and $b' < 0$ for the usual matter, except the higher-derivative conformal scalars. Notice that b'' can be shifted by the finite renormalization of the local counterterm R^2 , so b'' can be an arbitrary coefficient.

By including the trace anomaly, the trace of (II.5) is modified as

$$R + 2f(R) - Rf'(R) - 3\Box f'(R) = -\frac{\kappa^2}{2}(T_{\text{matter}} + T_A). \quad (\text{IV.87})$$

For the FRW universe, we find

$$\mathcal{F} = 0, \quad \mathcal{G} = 24(\dot{H}H^2 + H^4). \quad (\text{IV.88})$$

We now assume that H behaves as (IV.44) and neglect the contribution from matter by putting $T_{\text{matter}} = 0$. Then, in the case that $f(R)$ behaves as $f(R) \sim -\frac{\alpha}{R^n}$, Eq. (IV.87) becomes

$$R + 3\Box\left(\frac{\alpha n}{R^{n+1}}\right) = -\frac{\kappa^2}{2}T_A. \quad (\text{IV.89})$$

First, we consider the case that $2b/3 + b'' = 0$ and, therefore, $T_A = \mathcal{G}$. The solution (IV.83) is assumed. If $-1 < \beta < 0$, R behaves as $(t_s - t)^{-\beta-1}$ and $\mathcal{G}$ behaves as $\mathcal{G} \sim 24\dot{H}H^2 \sim (t_s - t)^{-3\beta-1}$. Because $-3\beta - 1 > -\beta - 1$ when $\beta < 0$, T_A is negligible compared with R . Here, $\Box(\alpha n/R^{n+1}) \sim (t_s - t)^{(\beta+1)(n+1)-2}$. Then, the curvature singularity appears in a finite time, and the quantum correction does not prevent the singularity when $2b/3 + b'' = 0$. One may, however, consider the case that $2b/3 + b'' \neq 0$. In this case, T_A behaves as

$$T_A \sim \left(\frac{2}{3}b + b''\right)\Box R \sim (t_s - t)^{-\beta-3}. \quad (\text{IV.90})$$

Because $R \sim (t_s - t)^{-\beta-1}$ and $\Box(\alpha n/R^{n+1}) \sim (t_s - t)^{(\beta+1)(n+1)-2}$, the terms in the l.h.s of (IV.89) are always less singular than T_A because

$$-\beta - 1, (\beta + 1)(n + 1) - 2 > -\beta - 3. \quad (\text{IV.91})$$

This indicates that Eq. (IV.89) does not allow the singular solution and the curvature singularity does not appear. Therefore, in the case that $2b/3 + b'' \neq 0$, the quantum effects prevent the singularity appearance.

In the above analysis, the $\Box R$ term acts against the singularity. The $\Box R$ term is generated by a local term R^2 , which shows that if one modifies $F(R)$ by adding the R^2 term as

$$F(R) \rightarrow F(R) + \gamma R^2, \quad (\text{IV.92})$$

with a constant γ , the curvature singularity cannot be generated. Thus, the addition of R^2 term which is relevant at the very early universe gives the universal tool for resolving the finite-time future singularities. Moreover, as was mentioned in section A of the second chapter, this term is the source of early-time inflation. Thus, removal of the singularity gives the natural prescription for the unification of the inflation with the dark energy epoch in frames of modified gravity.

C. Future singularities in $F(\mathcal{G})$ gravity

Let us show that the occurrence of future singularities is a general phenomenon in modified gravity. In this section, which is based on ref. [41], it is demonstrated that modified Gauss-Bonnet gravity can also drive the universe evolution to the singularity.

We start with the $F(\mathcal{G})$ models, in which the finite-time future singularities can occur when the form of H is taken as Eq. (IV.83). To find such $F(\mathcal{G})$ gravities, the reconstruction method is used in a form which is different from that in subsection III D.

Using proper functions $P(t)$ and $Q(t)$ of a scalar field t which is identified with the cosmic time, the action in Eq. (II.123) can be rewritten as

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} (R + P(t)\mathcal{G} + Q(t)) + \mathcal{L}_{\text{matter}} \right]. \quad (\text{IV.93})$$

The variation with respect to t yields

$$\frac{dP(t)}{dt}\mathcal{G} + \frac{dQ(t)}{dt} = 0, \quad (\text{IV.94})$$

from which one can find $t = t(\mathcal{G})$. Substituting $t = t(\mathcal{G})$ into Eq. (IV.93), the action in terms of $F(\mathcal{G})$ follows

$$F(\mathcal{G}) = P(t)\mathcal{G} + Q(t). \quad (\text{IV.95})$$

The scale factor is presented as

$$a(t) = a_0 \exp(g(t)), \quad (\text{IV.96})$$

where a_0 is a constant and $g(t)$ is a proper function. Using Eqs. (II.126), (II.173), (II.174), (IV.96), the matter conservation law (II.144) and then neglecting the contribution from matter, we get the differential equation

$$2\frac{d}{dt}\left(\dot{g}^2(t)\frac{dP(t)}{dt}\right) - 2\dot{g}^3(t)\frac{dP(t)}{dt} + \ddot{g}(t) = 0. \quad (\text{IV.97})$$

With the help of Eq. (II.126), $Q(t)$ is given by

$$Q(t) = -24\dot{g}^3(t)\frac{dP(t)}{dt} - 6\dot{g}^2(t). \quad (\text{IV.98})$$

First, the Big Rip singularity can be analyzed. If $\beta = 1$ in Eq. (IV.83) with $H_0 = 0$, H and $\mathcal{G}$ are given by

$$H = \frac{h_0}{(t_s - t)}, \quad \mathcal{G} = \frac{24h_0^3}{(t_s - t)^4}(1 + h_0). \quad (\text{IV.99})$$

The most general solution of Eq. (IV.97) is

$$P(t) = \frac{1}{4h_0(h_0 - 1)}(2t_s - t)t + c_1 \frac{(t_s - t)^{3-h_0}}{3 - h_0} + c_2, \quad (\text{IV.100})$$

where c_1 and c_2 are constants. From Eq. (IV.98), one gets

$$Q(t) = -\frac{6h_0^2}{(t_s - t)^2} - \frac{24h_0^3 \left[\frac{(t_s - t)}{2h_0(h_0 - 1)} - c_1(t_s - t)^{2-h_0} \right]}{(t_s - t)^3}. \quad (\text{IV.101})$$

Furthermore, from Eq. (IV.94), we obtain

$$t = \left[\frac{24(h_0^3 + h_0^4)}{\mathcal{G}} \right]^{1/4} + t_s. \quad (\text{IV.102})$$

Solving Eq. (IV.95), we find the most general form of $F(\mathcal{G})$, which realizes the Big Rip singularity

$$F(\mathcal{G}) = \frac{\sqrt{6h_0^3(1 + h_0)}}{h_0(1 - h_0)}\sqrt{\mathcal{G}} + c_1\mathcal{G}^{\frac{h_0+1}{4}} + c_2\mathcal{G}. \quad (\text{IV.103})$$

This is an exact solution in the case of Eq. (IV.99). In general, if, for large values of $\mathcal{G}$, $F(\mathcal{G}) \sim \alpha\mathcal{G}^{1/2}$, where $\alpha(\neq 0)$ is a constant, the Big Rip singularity occurs for any value of $h_0 \neq 1$. In the case of $h_0 = 1$, the solution of $\mathcal{G}(H, \dot{H} \dots)$ is zero for $F(\mathcal{G}) = \alpha\mathcal{G}^{1/2}$. Note that $\alpha\mathcal{G}^{(1+h_0)/4}$ is an invariant with respect to the Big Rip solution.

In the case of $h_0 = 1$, it is possible to find another exact solution for $P(t)$

$$P(t) = \alpha(t_s - t)^q \ln[\gamma(t_s - t)^z], \quad (\text{IV.104})$$

where $\gamma(> 0)$ is a positive constant and q and z are constants. Eq. (IV.97) is satisfied for the case of Eq. (IV.99) if $q = 3 - h_0 = 2$ (and, therefore, $h_0 = 1$) and $z\alpha = -1/4$. From Eq. (IV.98), we have

$$Q(t) = -\frac{12}{(t_s - t)^2} \ln[\gamma(t_s - t)]. \quad (\text{IV.105})$$

The form of $F(\mathcal{G})$ is given by

$$F(\mathcal{G}) = \frac{\sqrt{3}}{2} \sqrt{\mathcal{G}} \ln(\gamma \mathcal{G}). \quad (\text{IV.106})$$

In general, if, for large values of $\mathcal{G}$, $F(\mathcal{G}) \sim \alpha \sqrt{\mathcal{G}} \ln(\gamma \mathcal{G})$ with $\alpha > 0$ and $\gamma > 0$, the Big Rip singularity occurs. The same result is found for $F(\mathcal{G}) \sim \alpha \sqrt{\mathcal{G}} \ln(\gamma \mathcal{G}^z + G_0)$ with $\alpha > 0$, $\gamma > 0$ and $z > 0$, where G_0 is a constant.

Next, we investigate the other types of singularities. If $\beta \neq 1$, Eq. (IV.83) with $H_0 = 0$ implies that the scale factor $a(t)$ behaves as

$$a(t) = \exp \left[\frac{h_0(t_s - t)^{1-\beta}}{\beta - 1} \right]. \quad (\text{IV.107})$$

We consider the case in which H and $\mathcal{G}$ are given by

$$H = \frac{h_0}{(t_s - t)^\beta}, \quad \beta > 1, \quad \mathcal{G} \sim \frac{24h_0^4}{(t_s - t)^{4\beta}}. \quad (\text{IV.108})$$

A solution of Eq. (IV.97) in the limit $t \rightarrow t_s$ is given by

$$P(t) \simeq \frac{\alpha}{(t_s - t)^z} \quad (\text{IV.109})$$

with $z = -2\beta$ and $\alpha = -1/4h_0^2$. The form of $F(\mathcal{G})$ is expressed as

$$F(\mathcal{G}) = -12 \sqrt{\frac{\mathcal{G}}{24}}. \quad (\text{IV.110})$$

Thus, if, for large values of $\mathcal{G}$, $F(\mathcal{G}) \sim -\alpha \sqrt{\mathcal{G}}$ with $\alpha > 0$, a Type I singularity could appear.

When $\beta < 1$, the forms of H and $\mathcal{G}$ are given by

$$H = \frac{h_0}{(t_s - t)^\beta}, \quad 0 < \beta < 1, \quad \mathcal{G} \sim \frac{24h_0^3\beta}{(t_s - t)^{3\beta+1}}. \quad (\text{IV.111})$$

An asymptotic solution of Eq. (IV.97) in the limit $t \rightarrow t_s$ is given by

$$P(t) \simeq \frac{\alpha}{(t_s - t)^z} \quad (\text{IV.112})$$

with $z = -(1 + \beta)$ and $\alpha = 1/2h_0(1 + \beta)$. The form of $F(\mathcal{G})$ becomes

$$F(\mathcal{G}) = \frac{6h_0^2}{(\beta + 1)} (3\beta + 1) \left(\frac{|\mathcal{G}|}{24h_0^3|\beta|} \right)^{2\beta/(3\beta+1)}. \quad (\text{IV.113})$$

Thus, if for large values of $\mathcal{G}$, $F(\mathcal{G})$ has the form

$$F(\mathcal{G}) \sim \alpha |\mathcal{G}|^\gamma, \quad \gamma = \frac{2\beta}{3\beta + 1}, \quad (\text{IV.114})$$

with $\alpha > 0$ and $0 < \gamma < 1/2$, we find that $0 < \beta < 1$ and a Type III singularity could emerge.

If, for $\mathcal{G} \rightarrow -\infty$, $F(\mathcal{G})$ has the form in Eq. (IV.114) with $\alpha > 0$ and $-\infty < \gamma < 0$, we find $-1/3 < \beta < 0$ and a Type II (sudden) singularity could occur. Moreover, if for $\mathcal{G} \rightarrow 0^-$, $F(\mathcal{G})$ has the form in Eq. (IV.114) with $\alpha < 0$ and $1 < \gamma < \infty$, it is obtained $-1 < \beta < -1/3$ and a Type II singularity could occur.

If, for $\mathcal{G} \rightarrow 0^-$, $F(\mathcal{G})$ has the form in Eq. (IV.114) with $\alpha > 0$ and $2/3 < \gamma < 1$, we obtain $-\infty < \beta < -1$ and a Type IV singularity could emerge. It is also required that $\gamma \neq 2n/(3n-1)$, where n is a natural number.

We can generate all of the possible Type II singularities as shown above except in the case of $\beta = -1/3$, i.e., $H = h_0/(t_s - t)^{1/3}$. In this case, the form of $\mathcal{G}$ is:

$$\mathcal{G} = 24h_0^3\beta + 24h_0^4(t_s - t)^{4/3} < 0. \quad (\text{IV.115})$$

To find t in terms of $\mathcal{G}$, one must consider the whole expression of $\mathcal{G}$ by also taking into account the low term of $(t_s - t)$. It follows that

$$F(\mathcal{G}) \simeq \frac{1}{4\sqrt{6}h_0^3}\mathcal{G}(\mathcal{G} + 8h_0^3)^{1/2} + \frac{2}{\sqrt{6}}(\mathcal{G} + 8h_0^3)^{1/2}. \quad (\text{IV.116})$$

As a consequence, the specific model $F(\mathcal{G}) = \sigma_1\mathcal{G}(\mathcal{G} + c_3)^{1/2} + \sigma_2(\mathcal{G} + c_3)^{1/2}$, where σ_1 , σ_2 and c_3 are positive constants, can generate a Type II singularity.

Let us end this section by considering the following realistic model, again for $n > 0$,

$$f_4(\mathcal{G}) = \mathcal{G}^\alpha \frac{a_1\mathcal{G}^n + b_1}{a_2\mathcal{G}^n + b_2}. \quad (\text{IV.117})$$

Because for large $\mathcal{G}$, one has $F_4(\mathcal{G}) \simeq a_1/a_2\mathcal{G}^\alpha$ and for small $\mathcal{G}$, one has $f_4(\mathcal{G}) \simeq b_1/b_2\mathcal{G}^\alpha$, we do not find any type of singularities for

$$\frac{1}{2} < \alpha < \frac{2}{3}. \quad (\text{IV.118})$$

Eq. (IV.117) provides an example of a realistic model free of all possible future singularities when Eq. (IV.118) is satisfied, independently of the coefficients. Moreover, this model suggests the universal scenario for resolving future finite-time singularities. Adding the above model to any singular dark energy results in a combined non-singular model. Thus, unlike convenient dark energy, which may be singular or not, (non-singular) modified gravity may suggest the universal recipe for resolving the future finite-time singularity. In this respect, modified gravity seems to be a more fundamental theory than convenient dark energies.

D. Late-time dynamics in Hořava-Lifshitz gravity

Let us show that future singularities may also occur in Hořava-Lifshitz dark energy models [79]. Assume the behavior of the Hubble rate is the same as in (IV.44). Let us now consider some simple singularity examples in the power-law Hořava-Lifshitz $F(R)$ gravity and compare them with the standard case.

In the case that

$$F(\tilde{R}) = \tilde{R} + \chi\tilde{R}^m, \quad (\text{IV.119})$$

one finds the necessary conditions for the presence of the singularities:

$$\begin{array}{ll} \text{Type I} & \beta > 1, \quad m > \frac{\beta-1}{2\beta} \\ \text{Type II} & m < 0, \quad -1 < \beta < 0 \\ \text{Type III} & 0 < \beta < 1, \quad m \neq 0 \\ \text{Type IV} & \beta \in \mathbb{Q} - \mathbb{Z}, \quad \beta < -1. \end{array} \quad (\text{IV.120})$$

It is important to stress that the conditions (IV.120) are only necessary for the existence of the singularity. In the following, only the case that $\lambda > \frac{1}{3}$ is considered for simplicity. Then, one sees that a solution

$$a = a_0 t^\gamma, \quad \rho = \rho_0 t^{-2}, \quad \gamma = \frac{2(m-1)(2m-1)\mu}{(2m-1)(3\lambda-1)-6(m-1)\mu}, \quad \rho_0 = 0, \quad (\text{IV.121})$$

presents a singularity of Type I for

$$\left\{ \begin{array}{l} 0 < m < \frac{1}{2} \quad \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8} < \mu < \frac{6m\lambda-2m-3\lambda+1}{8m^3-16m^2+16m-8}, \\ \frac{1}{2} < m < 1 \quad \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8} < \mu < \frac{6m\lambda-2m-3\lambda+1}{6m-6}, \\ 1 < m < 2 \quad \frac{6m\lambda-2m-3\lambda+1}{6m-6} < \mu < \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8}, \\ m = 2 \quad \mu > \frac{1}{6}(9\lambda-3), \\ m > 2 \quad \mu < \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8} \quad \mu > \frac{6m\lambda-2m-3\lambda+1}{6m-6}, \end{array} \right. \quad (\text{IV.122})$$

a singularity of Type II for

$$\left\{ \begin{array}{l} m < -1 \quad \frac{6m\lambda-2m-3\lambda+1}{4m^2-4} < \mu < 0, \\ m = -1 \quad \mu < 0, \\ -1 < m < 0 \quad \mu < 0 \quad \mu > \frac{6m\lambda-2m-3\lambda+1}{4m^2-4}, \end{array} \right. \quad (\text{IV.123})$$

and a singularity of Type III for

$$\left\{ \begin{array}{l} m < 0 \quad \mu < 0 \quad \mu > \frac{6m\lambda-2m-3\lambda+1}{6m-6}, \\ 0 < m < \frac{1}{2} \quad 0 < \mu < \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8}, \\ \frac{1}{2} < m < 1 \quad \mu < \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8} \quad \mu > 0, \\ 1 < m < 2 \quad \mu < 0 \quad \mu > \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8}, \\ m = 2 \quad \mu < 0 \\ m > 2 \quad \frac{-6m\lambda+2m+3\lambda-1}{4m^2-12m+8} < \mu < 0. \end{array} \right. \quad (\text{IV.124})$$

This is very different from the standard case, where we have a singularity of Type I only for $m > 2$, and a Type III for $\frac{1}{2} < m < 1$ and $m < 0$.

For the solution

$$\begin{aligned} a &= a_0 t^\gamma, \quad \rho = \rho_0 t^{-2}, \quad \gamma = \frac{2m}{3(1+w)}, \\ \rho_0 &= \frac{\chi [3\mu(m-1)(2m(w+2)-w-1)-m(2m-1)(3\lambda-1)]}{\kappa^2 (m[3\lambda-6\mu-1]+3(w+1)\mu)} \\ &\quad \times \left(\frac{4m^2(-3\lambda+6\mu+1)-12m(w+1)\mu}{(w+1)^2} \right)^m, \end{aligned} \quad (\text{IV.125})$$

we have a singularity of Type III when

$$\left\{ \begin{array}{l} -3 < m \leq -\frac{3}{2} \quad \frac{1}{3}(-2m-3) < w \leq 1, \\ -\frac{3}{2} < m < \frac{1}{2}(1-\sqrt{13}) \quad 0 \leq w \leq 1. \end{array} \right. \quad (\text{IV.126})$$

For the solution

$$a = a_0 t^\gamma, \quad \rho = \rho_0 t^{-2}, \quad \gamma = \frac{2\mu}{-3\lambda+6\mu+1}, \quad \rho_0 = 0, \quad (\text{IV.127})$$

which exists only in the Hořava-Lifshitz version of $F(R)$ gravity, we have instead a singularity of Type I for

$$\left\{ \begin{array}{l} \lambda < \frac{1}{3} \quad \frac{1}{6}(3\lambda-1) < \mu < \frac{1}{8}(3\lambda-1) \quad m > 1, \\ \lambda > \frac{1}{3} \quad \frac{1}{8}(3\lambda-1) < \mu < \frac{1}{6}(3\lambda-1) \quad m > 1, \end{array} \right. \quad (\text{IV.128})$$

and of Type III for

$$\left\{ \begin{array}{l} \lambda < \frac{1}{3} \quad \frac{1}{8}(3\lambda-1) < \mu < 0 \quad m > 1, \\ \lambda > \frac{1}{3} \quad 0 < \mu < \frac{1}{8}(3\lambda-1) \quad m > 1. \end{array} \right. \quad (\text{IV.129})$$

The results above clearly show that the presence of singularities is deeply altered in the Hořava-Lifshitz version of $F(R)$ gravity. In particular, it seems that the additional parameters make it much easier to realize the singularities. The intervals we have presented above for the parameters can then be interpreted as constraints on this type of Hořava-Lifshitz $F(R)$ gravity. Thus, we have demonstrated that modified Hořava-Lifshitz gravity can be incorporated into the phantom like or quintessence like accelerating cosmologies, which might lead to singularities of Type I, II,

or III. Of course, as in the case of conventional theory one can resolve these future singularities by adding a higher-derivative gravitational term, which is relevant at very early times. However, the structure of this term is different from the case of convenient modified gravity. For example, the addition of the R^2 term does not necessarily remove all types of singularities in the theory under discussion.

In summary, in this chapter, we demonstrated that an effective phantom as well as some quintessence dark energy models may drive the universe evolution to future finite-time singularities. This is a universal property, regardless of whether dark energy is a fluid, field model or modified gravity. To resolve such future singularities, one has to add extra gravitational terms to the theory action. In other words, resolving the singularities invites the modification of gravity, even if the initial dark energy was not modified gravity. The additional gravitational term, which is usually relevant at very early times, also helps in the realization of the inflationary phase. In this way, the additional support for unification of early-time inflation with late-time acceleration in modified gravity follows.

V. DISCUSSION

In summary, we reviewed a number of popular modified gravities including $F(R)$ theory, modified Gauss-Bonnet and scalar-Gauss-Bonnet gravities, non-minimal models, non-local gravity, modified $F(R)$ Hořava-Lifshitz theory and (power-counting renormalizable) covariant gravity. General properties and different representations of such theories were considered. Their spatially flat FRW equations, which often have higher-derivative terms and/or non-canonical and non-standard couplings were derived. The accelerating cosmological solutions including the de Sitter universe and an attempt to obtain the unified description of inflation with dark energy were proposed. It was demonstrated that the qualitative possibility of such a unification is a very natural property for this large class of alternative gravities. This is also the case for the theories with broken Lorentz symmetry like Hořava-Lifshitz gravity.

To describe the background evolution of modified gravity, we developed the cosmological reconstruction method (inverse problem) in terms of cosmological time or e-folding. Using the freedom in the choice of scalar potentials and of the modified term function, which depends on geometrical invariants, such as curvature and the Gauss-Bonnet term, we arrived to master differential equations whose solution solves the problem. It was demonstrated that eventually, any given cosmology may be reconstructed in this way. We explicitly considered the reconstruction in scalar-tensor theory, Brans-Dicke gravity, the k-essence model, $F(R)$ theory and the Lagrange multiplier $F(R)$ theory, modified Gauss-Bonnet and scalar-Gauss-Bonnet gravities, Hořava-Lifshitz gravity and non-minimal Yang-Mills theory. As an example of the number of universe evolution eras including Λ CDM one, the effective quintessence or phantom acceleration, the unified early-time inflation with late-time acceleration was shown to be the background solution of the above theories. It should be stressed that even if specific modified gravity cannot capture the entire sequence of the universe evolution (namely, the inflation, radiation and matter dominance, dark energy era) with good accuracy, it may be additionally reconstructed so that it can become a viable theory. Thus, such a partial reconstruction is an extremely powerful tool that may help the theory to pass the known observationally related tests. Of course, the price for that is some extra modifications of the action. Fortunately, such extra modifications may be attributed to the gravitational sector relevant at the very early universe.

Special attention was paid to late-time dynamics of the effective quintessence/phantom dark energy of arbitrary nature: fluid, particle model or modified gravity. It is known that all or some of the energy conditions for such a (phantom or quintessence) dark energy universe are violated. As a result, the universe may end up with its evolution in one of four known types of finite-time singularities. The appearance of all four types of future singularities in coupled fluid dark energy, $F(R)$ theory, modified Gauss-Bonnet gravity and modified $F(R)$ Hořava-Lifshitz gravity was demonstrated. The universal prescription for resolving such singularities that may lead to bad phenomenological consequences was proposed. It consists of the addition of an extra higher-derivative gravitational term relevant in the very early universe and helping in the realization of the inflation. In the case of $F(R)$ theory or standard fluid dark energy, this term is proportional to R^2 at large curvatures.

The advantage of the approach proposed in this review is its very general character. Even if one of the particular gravity models under discussion is outside of mainstream research, its properties and background solutions will remain mathematically and physically correct. Moreover, the developed reconstruction scheme proposes the way to change the properties of any particular theory in the desirable way. Thus, we did not discuss the cosmological perturbation theory or structure formation in modified gravity. Indeed, the standard way to study the cosmological perturbations in modified gravity is based on the close analogy with general relativity. As a result, higher-order differential equations are approximated by second-order equations. Clearly, this is not a consistent approach. To analyze the cosmological perturbations one has to develop a powerful numerical technique that should be applied to higher-derivative differential equations. We also did not discuss the local tests of the gravity (especially, post-Newtonian regime tests) in detail. This is due to existence of a number of observationally related reviews [4] from a point of view. From another point of view, only the small curvature regime is relevant for the purposes of the local tests. Clearly, all modified gravities

under consideration may be parametrized by some effective theory with small deviations from general relativity. In other words, a very large class of models with qualitatively different properties may quite easily pass the local tests.

The next important step in the investigation of modified gravity should be related to a detailed proposal on the description of the sequence of the entire evolution of the universe: inflation, radiation/matter dominance, and dark energy. As we saw in this work, the equations of motion for such theories are extremely complicated, and including all of the universe matter content makes them even more involved. Presumably, such a study can be done only numerically. This is an extremely hard task that should give the final answer to the question: is the evolution of the observable universe governed by modified gravity?

Acknowledgments

We are grateful to all of our friends and collaborators, especially, those listed here: M. C. B. Abdalla, G. Allemandi, K. Bamba, A. Borowiec, K. A. Bronnikov, I. Brevik, F. Briscese, S. Carloni, S. Capozziello, V. Cardone, M. Chaichian, G. Cognola, A. Dev, P. K. S. Dunsby, E. Elizalde, V. Faraoni, M. Francaviglia, C. Q. Geng, O. G. Gorbunova, R. Goswami, Y. Ito, D. Jain, S. Jhingan, M. De Laurentis, J. E. Lidsey, J. Matsumoto, M. Oksanen, D. Saez-Gomez, M. Sami, M. Sasaki, L. Sebastiani, I. Thongkool, A. Toporensky, A. Troisi, P. V. Tretyakov, A. Tureanu, L. Vanzo, P. Wang, H. Yoshioka, S. Zerbini, and Y. I. Zhang for stimulating discussions of different aspects of modified gravity. This research has been supported in part by the MEC (Spain) project FIS2006-02842 and AGAUR (Catalonia) 2009SGR-994 (SDO), by the Global COE Program of Nagoya University (G07) provided by the Ministry of Education, Culture, Sports, Science & Technology and by the JSPS Grant-in-Aid for Scientific Research (S) # 22224003 (SN).

-
- [1] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, *Phys. Rev. Lett.* **91**, 071301 (2003) [arXiv:astro-ph/0302506].
 - [2] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **68**, 123512 (2003) [arXiv:hep-th/0307288].
 - [3] S. Nojiri and S. D. Odintsov, *eConf C0602061*, 06 (2006) [*Int. J. Geom. Meth. Mod. Phys.* **4**, 115 (2007)] [arXiv:hep-th/0601213]; arXiv:0807.0685.
 - [4] S. Capozziello, M. De Laurentis and V. Faraoni, arXiv:0909.4672 [gr-qc];
S. Capozziello and M. Francaviglia, *Gen. Rel. Grav.* **40**, 357 (2008) [arXiv:0706.1146 [astro-ph]];
T. P. Sotiriou and V. Faraoni, *Rev. Mod. Phys.* **82**, 451 (2010) [arXiv:0805.1726 [gr-qc]];
F. S. N. Lobo, arXiv:0807.1640 [gr-qc].
 - [5] W. Nelson, *Phys. Rev. D* **82**, 124044 (2010) [arXiv:1012.3353 [gr-qc]]. H. W. Lee, K. Y. Kim and Y. S. Myung, arXiv:1010.5556 [hep-th];
L. Amendola, K. Enqvist and T. Koivisto, arXiv:1010.4776 [gr-qc];
Y. Zhong, Y. X. Liu and K. Yang, arXiv:1010.3478 [hep-th];
D. S. Gorbunov and A. G. Panin, arXiv:1009.2448 [hep-ph];
C. Corda, *Eur. Phys. J. C* **65**, 257 (2010) [arXiv:1007.4077 [gr-qc]];
L. G. Jaime, L. Patino and M. Salgado, arXiv:1006.5747 [gr-qc];
H. Farajollahi, M. Setare, F. Milani and F. Tayebi, arXiv:1005.2026 [physics.gen-ph];
A. Balcerzak and M. P. Dabrowski, *Phys. Rev. D* **81**, 123527 (2010) [arXiv:1004.0150 [hep-th]];
S. Capozziello, P. Martin-Moruno and C. Rubano, *Phys. Lett. B* **689**, 117 (2010) [arXiv:1003.5394 [gr-qc]];
F. Schmidt, *Phys. Rev. D* **81**, 103002 (2010) [arXiv:1003.0409 [astro-ph.CO]];
T. Saidov and A. Zhuk, *Phys. Rev. D* **81**, 124002 (2010) [arXiv:1002.4138 [hep-th]];
I. Brevik, S. Nojiri, S. D. Odintsov and D. Saez-Gomez, arXiv:1002.1942 [hep-th];
R. Garcia-Salcedo, T. Gonzalez, C. Moreno, Y. Napoles, Y. Leyva and I. Quiros, *JCAP* **1002**, 027 (2010) [arXiv:0912.5048 [gr-qc]];
V. Dzhunushaliev, V. Folomeev, B. Kleihaus and J. Kunz, *JHEP* **1004**, 130 (2010) [arXiv:0912.2812 [gr-qc]];
A. Aghmohammadi, K. Saaïdi, M. R. Abolhassani and A. Vajdi, *Phys. Scripta* **80**, 065008 (2009) [arXiv:0912.2510 [gr-qc]];
T. Multamaki, J. Vainio and I. Vilja, *Phys. Rev. D* **81**, 064025 (2010) [arXiv:0910.5659 [gr-qc]];
R. G. Cai, L. M. Cao, Y. P. Hu and N. Ohta, *Phys. Rev. D* **80**, 104016 (2009) [arXiv:0910.2387 [hep-th]];
K. Bamba, C. Q. Geng, S. Nojiri and S. D. Odintsov, *Europhys. Lett.* **89**, 50003 (2010) [arXiv:0909.4397 [hep-th]];
K. Nozari and T. Azizi, *Phys. Lett. B* **680**, 205 (2009) [arXiv:0909.0351 [gr-qc]];
S. Tsujikawa, R. Gannouji, B. Moraes and D. Polarski, *Phys. Rev. D* **80**, 084044 (2009) [arXiv:0908.2669 [astro-ph.CO]];
S. K. Chakrabarti, E. N. Saridakis and A. A. Sen, arXiv:0908.0293 [astro-ph.CO];
I. Quiros, Y. Leyva and Y. Napoles, *Phys. Rev. D* **80**, 024022 (2009) [arXiv:0906.1190 [gr-qc]];
L. Hui, A. Nicolis and C. Stubbs, *Phys. Rev. D* **80**, 104002 (2009) [arXiv:0905.2966 [astro-ph.CO]];
C. Gao, *Phys. Lett. B* **684**, 85 (2010) [arXiv:0905.0310 [astro-ph.CO]];
B. C. Paul, P. S. Debnath and S. Ghose, *Phys. Rev. D* **79**, 083534 (2009) [arXiv:0904.0345 [astro-ph.CO]];

- S. V. Ketov, *Class. Quant. Grav.* **26**, 135006 (2009) [arXiv:0903.0251 [hep-th]];
 F. Bauer, J. Sola and H. Stefancic, *Phys. Lett. B* **678**, 427 (2009) [arXiv:0902.2215 [hep-th]];
 M. R. Setare, *Int. J. Mod. Phys. D* **17**, 2219 (2008) [arXiv:0901.3252 [hep-th]].
- [6] L. N. Granda, arXiv:0812.1596 [hep-th];
 R. Durrer and R. Maartens, arXiv:0811.4132 [astro-ph];
 T. S. Koivisto, arXiv:0811.1957 [astro-ph];
 V. Faraoni, arXiv:0811.1870 [gr-qc];
 E. Dyer and K. Hinterbichler, *Phys. Rev. D* **79**, 024028 (2009) [arXiv:0809.4033 [gr-qc]];
 D. Saez-Gomez, *Gen. Rel. Grav.* **41**, 1527 (2009) [arXiv:0809.1311 [hep-th]];
 M. Ishak and J. Moldenhauer, *JCAP* **0901**, 024 (2009) [arXiv:0808.0951 [astro-ph]];
 O. M. Lecian and G. Montani, *Class. Quant. Grav.* **26**, 045014 (2009) [arXiv:0807.4428 [gr-qc]];
 H. Stefancic, *Phys. Lett. B* **670**, 246 (2009) [arXiv:0807.3692 [gr-qc]];
 S. Nojiri and S. D. Odintsov, *AIP Conf. Proc.* **1115**, 212 (2009) [arXiv:0810.1557 [hep-th]];
 J. L. Cortes and J. Indurain, *Astropart. Phys.* **31**, 177 (2009) [arXiv:0805.3481 [astro-ph]];
 T. Multamaki, J. Vainio and I. Vilja, *Class. Quant. Grav.* **26**, 075005 (2009) [arXiv:0805.1834 [gr-qc]];
 B. Vakili, *Phys. Lett. B* **664**, 16 (2008) [arXiv:0804.3449 [gr-qc]];
 N. Deruelle, M. Sasaki and Y. Sendouda, *Phys. Rev. D* **77**, 124024 (2008) [arXiv:0803.2742 [gr-qc]]. S. A. Appleby and R. A. Battye, *JCAP* **0805**, 019 (2008) [arXiv:0803.1081 [astro-ph]];
 A. de la Cruz-Dombriz, A. Dobado and A. L. Maroto, *Phys. Rev. D* **77**, 123515 (2008) [arXiv:0802.2999 [astro-ph]];
 S. DeDeo and D. Psaltis, *Phys. Rev. D* **78**, 064013 (2008) [arXiv:0712.3939 [astro-ph]];
 X. Wu and Z. H. Zhu, *Phys. Lett. B* **660**, 293 (2008) [arXiv:0712.3603 [astro-ph]];
 P. F. Machado and F. Saueressig, *Phys. Rev. D* **77**, 124045 (2008) [arXiv:0712.0445 [hep-th]];
 T. Multamaki, A. Putaja, I. Vilja and E. C. Vagenas, *Class. Quant. Grav.* **25**, 075017 (2008) [arXiv:0712.0276 [gr-qc]];
 S. Tsujikawa, K. Uddin and R. Tavakol, *Phys. Rev. D* **77**, 043007 (2008) [arXiv:0712.0082 [astro-ph]];
 J. D. Evans, L. M. H. Hall and P. Caillol, *Phys. Rev. D* **77**, 083514 (2008) [arXiv:0711.3695 [astro-ph]];
 G. Barenboim and J. Lykken, *JCAP* **0803**, 017 (2008) [arXiv:0711.3653 [astro-ph]];
 C. G. Boehmer, T. Harko and F. S. N. Lobo, *JCAP* **0803**, 024 (2008) [arXiv:0710.0966 [gr-qc]];
 R. Saffari and S. Rahvar, *Phys. Rev. D* **77**, 104028 (2008) [arXiv:0708.1482 [astro-ph]];
 A. Iglesias, N. Kaloper, A. Padilla and M. Park, *Phys. Rev. D* **76**, 104001 (2007) [arXiv:0708.1163 [astro-ph]];
 J. Santos, J. S. Alcaniz, M. J. Reboucas and F. C. Carvalho, *Phys. Rev. D* **76**, 083513 (2007) [arXiv:0708.0411 [astro-ph]];
 T. P. Sotiriou, V. Faraoni and S. Liberati, *Int. J. Mod. Phys. D* **17**, 399 (2008) [arXiv:0707.2748 [gr-qc]];
 F. Schmidt, M. Liguori and S. Dodelson, *Phys. Rev. D* **76**, 083518 (2007) [arXiv:0706.1775 [astro-ph]];
 J. C. C. de Souza and V. Faraoni, *Class. Quant. Grav.* **24**, 3637 (2007) [arXiv:0706.1223 [gr-qc]];
 S. Rahvar and Y. Sobouti, *Mod. Phys. Lett. A* **23**, 1929 (2008) [arXiv:0704.0680 [astro-ph]];
 V. Faraoni, *Phys. Rev. D* **75**, 067302 (2007) [arXiv:gr-qc/0703044];
 L. M. Sokolowski, *Class. Quant. Grav.* **24**, 3391 (2007) [arXiv:gr-qc/0702097];
 T. Rador, *Phys. Lett. B* **652**, 228 (2007) [arXiv:hep-th/0702081];
 S. Capozziello and R. Garattini, *Class. Quant. Grav.* **24**, 1627 (2007) [arXiv:gr-qc/0702075];
 B. Li and J. D. Barrow, *Phys. Rev. D* **75**, 084010 (2007) [arXiv:gr-qc/0701111].
- [7] T. P. Sotiriou, *Phys. Lett. B* **645**, 389 (2007) [arXiv:gr-qc/0611107];
 N. J. Poplawski, *Phys. Rev. D* **74**, 084032 (2006) [arXiv:gr-qc/0607124];
 A. Borowiec, W. Godlowski and M. Szydlowski, *eConf C0602061*, 09 (2006) [*Int. J. Geom. Meth. Mod. Phys.* **4**, 183 (2007)] [arXiv:astro-ph/0607639];
 A. de la Cruz-Dombriz and A. Dobado, *Phys. Rev. D* **74**, 087501 (2006) [arXiv:gr-qc/0607118];
 S. A. Bludman, arXiv:astro-ph/0605198;
 T. P. Sotiriou and S. Liberati, *Annals Phys.* **322**, 935 (2007) [arXiv:gr-qc/0604006];
 T. G. Rizzo, arXiv:hep-ph/0603242;
 S. K. Srivastava, *Int. J. Mod. Phys. D* **17**, 755 (2008) [arXiv:astro-ph/0602116];
 T. Koivisto, *Phys. Rev. D* **73**, 083517 (2006) [arXiv:astro-ph/0602031];
 K. Atazadeh and H. R. Sepangi, *Int. J. Mod. Phys. D* **16**, 687 (2007) [arXiv:gr-qc/0602028];
 H. J. Schmidt, *eConf C0602061*, 12 (2006) [*Int. J. Geom. Meth. Mod. Phys.* **4**, 209 (2007)] [arXiv:gr-qc/0602017];
 I. H. Brevik, *Int. J. Mod. Phys. D* **15**, 767 (2006) [arXiv:gr-qc/0601100];
 T. P. Sotiriou, *Phys. Rev. D* **73**, 063515 (2006) [arXiv:gr-qc/0509029];
 S. Capozziello, V. F. Cardone, E. Piedipalumbo and C. Rubano, *Class. Quant. Grav.* **23**, 1205 (2006) [arXiv:astro-ph/0507438];
 Y. Ezawa, H. Iwasaki, Y. Ohkuwa, S. Watanabe, N. Yamada and T. Yano, *Class. Quant. Grav.* **23**, 3205 (2006) [arXiv:gr-qc/0507060];
 T. Koivisto, *Class. Quant. Grav.* **23**, 4289 (2006) [arXiv:gr-qc/0505128];
 D. A. Easson, *Int. J. Mod. Phys. A* **19**, 5343 (2004) [arXiv:astro-ph/0411209];
 X. H. Meng and P. Wang, *Class. Quant. Grav.* **22**, 23 (2005) [arXiv:gr-qc/0411007];
 S. M. Carroll, A. De Felice, V. Duvvuri, D. A. Easson, M. Trodden and M. S. Turner, *Phys. Rev. D* **71**, 063513 (2005) [arXiv:astro-ph/0410031];
 G. Allemandi, A. Borowiec and M. Francaviglia, *Phys. Rev. D* **70**, 043524 (2004) [arXiv:hep-th/0403264];
 S. Nojiri, *TSPU Vestnik* **44N7**, 49 (2004) [arXiv:hep-th/0407099];

- I. H. Brevik, S. Nojiri, S. D. Odintsov and L. Vanzo, *Phys. Rev. D* **70**, 043520 (2004) [arXiv:hep-th/0401073];
D. N. Vollick, *Class. Quant. Grav.* **21**, 3813 (2004) [arXiv:gr-qc/0312041];
G. J. Olmo and W. Komp, arXiv:gr-qc/0403092;
E. E. Flanagan, *Class. Quant. Grav.* **21**, 417 (2003) [arXiv:gr-qc/0309015];
S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **576**, 5 (2003) [arXiv:hep-th/0307071].
- [8] G. Leon and E. N. Saridakis, arXiv:1007.3956 [gr-qc];
S. Carloni, R. Goswami and P. K. S. Dunsby, arXiv:1005.1840 [gr-qc];
H. Farajollahi and F. Milani, *Mod. Phys. Lett. A* **25**, 2349 (2010) [arXiv:1004.3512 [gr-qc]];
J. Santos, M. J. Reboucas and T. B. R. Oliveira, *Phys. Rev. D* **81**, 123017 (2010) [arXiv:1004.2501 [astro-ph.CO]];
K. Saaidi, A. Vaji and A. Aghamohammadi, *Gen. Rel. Grav.* **42**, 2421 (2010) [arXiv:1001.4149 [gr-qc]];
S. Capozziello, M. De Laurentis and A. Stabile, *Class. Quant. Grav.* **27**, 165008 (2010) [arXiv:0912.5286 [gr-qc]];
S. Nojiri, S. D. Odintsov, A. Toporensky and P. Tretyakov, *Gen. Rel. Grav.* **42**, 1997 (2010) [arXiv:0912.2488 [hep-th]];
V. Reijonen, arXiv:0912.0825 [gr-qc];
M. Sharif and M. F. Shamir, *Class. Quant. Grav.* **26**, 235020 (2009) [arXiv:0910.5787 [gr-qc]];
F. S. N. Lobo and M. A. Oliveira, *Phys. Rev. D* **80**, 104012 (2009) [arXiv:0909.5539 [gr-qc]];
V. Faraoni, *Class. Quant. Grav.* **26**, 195013 (2009) [arXiv:0909.0514 [gr-qc]];
A. M. Nzioki, S. Carloni, R. Goswami and P. K. S. Dunsby, *Phys. Rev. D* **81**, 084028 (2010) [arXiv:0908.3333 [gr-qc]];
A. de la Cruz-Dombriz, A. Dobado and A. L. Maroto, *Phys. Rev. D* **80**, 124011 (2009) [arXiv:0907.3872 [gr-qc]];
N. Goheer, J. Larena and P. K. S. Dunsby, *Phys. Rev. D* **80**, 061301 (2009) [arXiv:0906.3860 [gr-qc]];
M. F. Figueiro and A. Saa, *Phys. Rev. D* **80**, 063504 (2009) [arXiv:0906.2588 [gr-qc]];
D. Momeni, *Int. J. Mod. Phys. D* **18**, 1719 (2009) [arXiv:0903.0067 [gr-qc]];
G. Cognola, E. Elizalde, S. D. Odintsov, P. Tretyakov and S. Zerbini, *Phys. Rev. D* **79**, 044001 (2009) [arXiv:0810.4989 [gr-qc]];
T. Clifton, *Phys. Rev. D* **78**, 083501 (2008) [arXiv:0807.4682 [gr-qc]];
H. Oyaizu, M. Lima and W. Hu, *Phys. Rev. D* **78**, 123524 (2008) [arXiv:0807.2462 [astro-ph]];
L. Hollenstein and F. S. N. Lobo, *Phys. Rev. D* **78**, 124007 (2008) [arXiv:0807.2325 [gr-qc]];
C. S. J. Pun, Z. Kovacs and T. Harko, *Phys. Rev. D* **78**, 024043 (2008) [arXiv:0806.0679 [gr-qc]];
R. Goswami, N. Goheer and P. K. S. Dunsby, *Phys. Rev. D* **78**, 044011 (2008) [arXiv:0804.3528 [gr-qc]];
D. N. Vollick, *Phys. Rev. D* **76**, 124001 (2007) [arXiv:0710.1859 [gr-qc]];
N. Lanahan-Tremblay and V. Faraoni, *Class. Quant. Grav.* **24**, 5667 (2007) [arXiv:0709.4414 [gr-qc]];
S. Capozziello, A. Stabile and A. Troisi, *Class. Quant. Grav.* **25**, 085004 (2008) [arXiv:0709.0891 [gr-qc]];
C. G. Boehmer, L. Hollenstein and F. S. N. Lobo, *Phys. Rev. D* **76**, 084005 (2007) [arXiv:0706.1663 [gr-qc]];
K. Henttunen, T. Multamaki and I. Vilja, *Phys. Rev. D* **77**, 024040 (2008) [arXiv:0705.2683 [astro-ph]];
K. Kainulainen, J. Piilonen, V. Reijonen and D. Sunhede, *Phys. Rev. D* **76**, 024020 (2007) [arXiv:0704.2729 [gr-qc]];
T. Clifton, *Class. Quant. Grav.* **24**, 5073 (2007) [arXiv:gr-qc/0703126];
M. D. Seifert, *Phys. Rev. D* **76**, 064002 (2007) [arXiv:gr-qc/0703060];
I. Sawicki and W. Hu, *Phys. Rev. D* **75**, 127502 (2007) [arXiv:astro-ph/0702278];
G. Cognola, M. Gastaldi and S. Zerbini, *Int. J. Theor. Phys.* **47**, 898 (2008) [arXiv:gr-qc/0701138];
D. Bazeia, B. Carneiro da Cunha, R. Menezes and A. Y. Petrov, *Phys. Lett. B* **649**, 445 (2007) [arXiv:hep-th/0701106];
S. Baghram, M. Farhang and S. Rahvar, *Phys. Rev. D* **75**, 044024 (2007) [arXiv:astro-ph/0701013];
T. Multamaki and I. Vilja, *Phys. Rev. D* **76**, 064021 (2007) [arXiv:astro-ph/0612775];
J. A. Leach, S. Carloni and P. K. S. Dunsby, *Class. Quant. Grav.* **23**, 4915 (2006) [arXiv:gr-qc/0603012];
T. Clifton and J. D. Barrow, *Class. Quant. Grav.* **23**, 2951 (2006) [arXiv:gr-qc/0601118];
V. Faraoni, *Phys. Rev. D* **72**, 124005 (2005) [arXiv:gr-qc/0511094];
T. Clifton and J. D. Barrow, *Phys. Rev. D* **72**, 103005 (2005) [arXiv:gr-qc/0509059];
N. Furey and A. DeBenedictis, *Class. Quant. Grav.* **22**, 313 (2005) [arXiv:gr-qc/0410088];
S. Carloni, P. K. S. Dunsby, S. Capozziello and A. Troisi, *Class. Quant. Grav.* **22**, 4839 (2005) [arXiv:gr-qc/0410046];
U. Gunther, A. Zhuk, V. B. Bezerra and C. Romero, *Class. Quant. Grav.* **22**, 3135 (2005) [arXiv:hep-th/0409112];
H. Kleinert and H. J. Schmidt, *Gen. Rel. Grav.* **34**, 1295 (2002) [arXiv:gr-qc/0006074].
- [9] S. Capozziello, R. Cianci, M. De Laurentis and S. Vignolo, arXiv:1007.3670 [gr-qc];
K. Saaidi and A. Aghamohammadi, arXiv:1006.1842 [gr-qc];
X. Fu, P. Wu and H. Yu, *Eur. Phys. J. C* **68**, 271 (2010);
B. Jain and J. Khoury, *Annals Phys.* **325**, 1479 (2010) [arXiv:1004.3294 [astro-ph.CO]];
L. Lombriser, A. Slosar, U. Seljak and W. Hu, arXiv:1003.3009 [astro-ph.CO];
W. Cui, P. Zhang and X. Yang, *Phys. Rev. D* **81**, 103528 (2010) [arXiv:1001.5184 [astro-ph.CO]];
T. Narikawa and K. Yamamoto, *Phys. Rev. D* **81**, 043528 (2010) [Erratum-ibid. *D* **81**, 129903 (2010)] [arXiv:0912.1445 [astro-ph.CO]];
F. Schmidt, A. Vikhlinin and W. Hu, *Phys. Rev. D* **80**, 083505 (2009) [arXiv:0908.2457 [astro-ph.CO]];
Y. Bisabr, *Phys. Lett. B* **683**, 96 (2010) [arXiv:0907.3838 [gr-qc]];
S. Nesseris and A. Mazumdar, *Phys. Rev. D* **79**, 104006 (2009) [arXiv:0902.1185 [astro-ph.CO]];
S. Bellucci, S. Capozziello, M. De Laurentis and V. Faraoni, *Phys. Rev. D* **79**, 104004 (2009) [arXiv:0812.1348 [gr-qc]];
F. Schmidt, M. V. Lima, H. Oyaizu and W. Hu, *Phys. Rev. D* **79**, 083518 (2009) [arXiv:0812.0545 [astro-ph]];
A. Borisov and B. Jain, *Phys. Rev. D* **79**, 103506 (2009) [arXiv:0812.0013 [astro-ph]];
G. Lambiase, S. Mohanty and G. Scarpetta, *JCAP* **0807**, 019 (2008);

- S. Capozziello, M. De Laurentis, S. Nojiri and S. D. Odintsov, *Gen. Rel. Grav.* **41**, 2313 (2009) [arXiv:0808.1335 [hep-th]]; arXiv:0911.2139;
- A. Dev, D. Jain, S. Jhingan, S. Nojiri, M. Sami and I. Thongkool, *Phys. Rev. D* **78**, 083515 (2008) [arXiv:0807.3445 [hep-th]];
- J. U. Kang and G. Panotopoulos, *Phys. Lett. B* **677**, 6 (2009) [arXiv:0806.1493 [astro-ph]];
- F. C. Carvalho, E. M. Santos, J. S. Alcaniz and J. Santos, *JCAP* **0809**, 008 (2008) [arXiv:0804.2878 [astro-ph]];
- S. Capozziello, V. F. Cardone and V. Salzano, *Phys. Rev. D* **78**, 063504 (2008) [arXiv:0802.1583 [astro-ph]];
- L. Iorio and M. L. Ruggiero, *Schol. Res. Exch.* **2008**, 968393 (2008) [arXiv:0711.0256 [gr-qc]];
- V. Faraoni, *Phys. Rev. D* **76**, 127501 (2007) [arXiv:0710.1291 [gr-qc]];
- T. Multamaki and I. Vilja, *Phys. Lett. B* **659**, 843 (2008) [arXiv:0709.3422 [astro-ph]];
- K. N. Ananda, S. Carloni and P. K. S. Dunsby, *Phys. Rev. D* **77**, 024033 (2008) [arXiv:0708.2258 [gr-qc]];
- W. Hu and I. Sawicki, *Phys. Rev. D* **76**, 104043 (2007) [arXiv:0708.1190 [astro-ph]];
- S. Capozziello, A. Stabile and A. Troisi, *Phys. Rev. D* **76**, 104019 (2007) [arXiv:0708.0723 [gr-qc]];
- Y. S. Song, H. Peiris and W. Hu, *Phys. Rev. D* **76**, 063517 (2007) [arXiv:0706.2399 [astro-ph]];
- S. Fay, S. Nesseris and L. Perivolaropoulos, *Phys. Rev. D* **76**, 063504 (2007) [arXiv:gr-qc/0703006];
- P. J. Zhang, *Phys. Rev. D* **76**, 024007 (2007) [arXiv:astro-ph/0701662];
- S. Fay, R. Tavakol and S. Tsujikawa, *Phys. Rev. D* **75**, 063509 (2007) [arXiv:astro-ph/0701479];
- L. Amendola, R. Gannouji, D. Polarski and S. Tsujikawa, *Phys. Rev. D* **75**, 083504 (2007) [arXiv:gr-qc/0612180];
- G. J. Olmo, *Phys. Rev. Lett.* **98**, 061101 (2007) [arXiv:gr-qc/0612002];
- A. J. Bustelo and D. E. Barraco, *Class. Quant. Grav.* **24**, 2333 (2007) [arXiv:gr-qc/0611149];
- X. H. Jin, D. J. Liu and X. Z. Li, arXiv:astro-ph/0610854;
- N. J. Poplawski, *Class. Quant. Grav.* **24**, 3013 (2007) [arXiv:gr-qc/0610133];
- A. L. Erickcek, T. L. Smith and M. Kamionkowski, *Phys. Rev. D* **74**, 121501 (2006) [arXiv:astro-ph/0610483];
- B. Li, K. C. Chan and M. C. Chu, *Phys. Rev. D* **76**, 024002 (2007) [arXiv:astro-ph/0610794];
- G. Allemandi and M. L. Ruggiero, *Gen. Rel. Grav.* **39**, 1381 (2007) [arXiv:astro-ph/0610661];
- Y. S. Song, W. Hu and I. Sawicki, *Phys. Rev. D* **75**, 044004 (2007) [arXiv:astro-ph/0610532];
- S. Mendoza and Y. M. Rosas-Guevara, *Astron. Astrophys.* **472**, 367 (2007) [arXiv:astro-ph/0610390];
- G. Lambiase and G. Scarpetta, *Phys. Rev. D* **74**, 087504 (2006) [arXiv:astro-ph/0610367];
- D. Huterer and E. V. Linder, *Phys. Rev. D* **75**, 023519 (2007) [arXiv:astro-ph/0608681];
- A. W. Brookfield, C. van de Bruck and L. M. H. Hall, *Phys. Rev. D* **74**, 064028 (2006) [arXiv:hep-th/0608015];
- M. L. Ruggiero and L. Iorio, *JCAP* **0701**, 010 (2007) [arXiv:gr-qc/0607093];
- V. Faraoni, *Phys. Rev. D* **74**, 023529 (2006) [arXiv:gr-qc/0607016];
- O. Mena, J. Santiago and J. Weller, *Phys. Rev. Lett.* **96**, 041103 (2006) [arXiv:astro-ph/0510453].
- [10] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, *JCAP* **0502**, 010 (2005) [arXiv:hep-th/0501096].
- [11] S. Nojiri and S. D. Odintsov, *AIP Conf. Proc.* **1241**, 1094 (2010) [arXiv:0910.1464 [hep-th]].
- [12] A. De Felice and S. Tsujikawa, *Living Rev. Rel.* **13**, 3 (2010) [arXiv:1002.4928 [gr-qc]].
- [13] S. Carloni, P. K. S. Dunsby and A. Troisi, *Phys. Rev. D* **77**, 024024 (2008) [arXiv:0707.0106 [gr-qc]].
- [14] S. Carloni, E. Elizalde and S. Odintsov, *Gen. Rel. Grav.* **42**, 1667 (2010) [arXiv:0907.3941 [gr-qc]].
- [15] W. Hu and I. Sawicki, *Phys. Rev. D* **76**, 064004 (2007) [arXiv:0705.1158 [astro-ph]].
- [16] S. Capozziello, S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **634**, 93 (2006) [arXiv:hep-th/0512118].
- [17] R. R. Caldwell, *Phys. Lett. B* **545**, 23 (2002) [arXiv:astro-ph/9908168].
- [18] S. Capozziello, S. Nojiri, S. D. Odintsov and A. Troisi, *Phys. Lett. B* **639**, 135 (2006) [arXiv:astro-ph/0604431].
- [19] A. D. Dolgov and M. Kawasaki, *Phys. Lett. B* **573**, 1 (2003) [arXiv:astro-ph/0307285].
- [20] S. Capozziello, *Int. J. Mod. Phys. D* **11**, 483 (2002) [arXiv:gr-qc/0201033].
- [21] S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, *Phys. Rev. D* **70**, 043528 (2004) [arXiv:astro-ph/0306438].
- [22] S. Capozziello, S. Carloni and A. Troisi, *Recent Res. Dev. Astron. Astrophys.* **1**, 625 (2003) [arXiv:astro-ph/0303041].
- [23] S. Capozziello, arXiv:gr-qc/0412088;
- X. H. Meng and P. Wang, *Gen. Rel. Grav.* **36**, 1947 (2004) [arXiv:gr-qc/0311019];
- A. E. Dominguez and D. E. Barraco, *Phys. Rev. D* **70**, 043505 (2004) [arXiv:gr-qc/0408069];
- G. Allemandi, M. Francaviglia, M. L. Ruggiero and A. Tartaglia, *Gen. Rel. Grav.* **37**, 1891 (2005) [arXiv:gr-qc/0506123];
- T. Koivisto, *Class. Quant. Grav.* **23**, 4289 (2006) [arXiv:gr-qc/0505128];
- G. J. Olmo, *Phys. Rev. D* **72**, 083505 (2005);
- I. Navarro and K. Van Acoleyen, *Phys. Lett. B* **622**, 1 (2005) [arXiv:gr-qc/0506096];
- J. A. R. Cembranos, *Phys. Rev. D* **73**, 064029 (2006) [arXiv:gr-qc/0507039]. T. P. Sotiriou, *Gen. Rel. Grav.* **38**, 1407 (2006) [arXiv:gr-qc/0507027];
- M. Amarzguoui, O. Elgaroy, D. F. Mota and T. Multamaki, *Astron. Astrophys.* **454**, 707 (2006) [arXiv:astro-ph/0510519];
- C. G. Shao, R. G. Cai, B. Wang and R. K. Su, *Phys. Lett. B* **633**, 164 (2006) [arXiv:gr-qc/0511034];
- K. A. Bronnikov, S. A. Kononogov and V. N. Melnikov, *Gen. Rel. Grav.* **38**, 1215 (2006) [arXiv:gr-qc/0601114];
- K. Atazadeh and H. R. Sepangi, *Int. J. Mod. Phys. D* **16**, 687 (2007) [arXiv:gr-qc/0602028].
- [24] M. E. Soussa and R. P. Woodard, *Gen. Rel. Grav.* **36**, 855 (2004) [arXiv:astro-ph/0308114].
- [25] R. P. Woodard, *Lect. Notes Phys.* **720**, 403 (2007) [arXiv:astro-ph/0601672].
- [26] S. Nojiri and S. D. Odintsov, *Mod. Phys. Lett. A* **19**, 627 (2004) [arXiv:hep-th/0310045].
- [27] T. Chiba, *Phys. Lett. B* **575**, 1 (2003) [arXiv:astro-ph/0307338].
- [28] S. Nojiri and S. D. Odintsov, *Gen. Rel. Grav.* **36**, 1765 (2004) [arXiv:hep-th/0308176].

- [29] X. H. Meng and P. Wang, Phys. Lett. B **584**, 1 (2004) [arXiv:hep-th/0309062].
- [30] S. Nojiri and S. D. Odintsov, Phys. Lett. B **657**, 238 (2007) [arXiv:0707.1941 [hep-th]].
- [31] S. Nojiri and S. D. Odintsov, Phys. Rev. D **77**, 026007 (2008) [arXiv:0710.1738 [hep-th]].
- [32] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani and S. Zerbini, Phys. Rev. D **77**, 046009 (2008) [arXiv:0712.4017 [hep-th]].
- [33] B. McInnes, JHEP **0208** (2002) 029 [arXiv:hep-th/0112066];
 S. Nojiri and S. D. Odintsov, Phys. Lett. B **562**, 147 (2003) [arXiv:hep-th/0303117];
 S. Nojiri and S. D. Odintsov, Phys. Lett. B **571**, 1 (2003) [arXiv:hep-th/0306212];
 V. Faraoni, Int. J. Mod. Phys. D **11**, 471 (2002) [arXiv:astro-ph/0110067];
 P. F. Gonzalez-Diaz, Phys. Lett. B **586**, 1 (2004) [arXiv:astro-ph/0312579];
 P. F. Gonzalez-Diaz, TSPU Vestnik **44N7**, 36 (2004) [arXiv:hep-th/0408225];
 B. McInnes, Nucl. Phys. B **718**, 55 (2005) [arXiv:hep-th/0502209];
 P. Singh, M. Sami and N. Dadhich, Phys. Rev. D **68**, 023522 (2003) [arXiv:hep-th/0305110];
 C. Csaki, N. Kaloper and J. Terning, Annals Phys. **317**, 410 (2005) [arXiv:astro-ph/0409596]. P. X. Wu and H. W. Yu, Nucl. Phys. B **727**, 355 (2005) [arXiv:astro-ph/0407424]. S. Nesseris and L. Perivolaropoulos, Phys. Rev. D **70**, 123529 (2004) [arXiv:astro-ph/0410309]. M. Sami and A. Toporensky, Mod. Phys. Lett. A **19**, 1509 (2004) [arXiv:gr-qc/0312009];
 H. Stefancic, Phys. Lett. B **586**, 5 (2004) [arXiv:astro-ph/0310904];
 L. P. Chimento and R. Lazkoz, Phys. Rev. Lett. **91**, 211301 (2003) [arXiv:gr-qc/0307111];
 Mod. Phys. Lett. A **19**, 2479 (2004) [arXiv:gr-qc/0405020];
 J. G. Hao and X. Z. Li, Phys. Lett. B **606**, 7 (2005) [arXiv:astro-ph/0404154];
 E. Babichev, V. Dokuchaev and Yu. Eroshenko, Class. Quant. Grav. **22**, 143 (2005) [arXiv:astro-ph/0407190];
 X. F. Zhang, H. Li, Y. S. Piao and X. M. Zhang, Mod. Phys. Lett. A **21**, 231 (2006) [arXiv:astro-ph/0501652];
 E. Elizalde, S. Nojiri, S. D. Odintsov and P. Wang, Phys. Rev. D **71**, 103504 (2005) [arXiv:hep-th/0502082];
 M. P. Dabrowski and T. Stachowiak, Annals Phys. **321**, 771 (2006) [arXiv:hep-th/0411199];
 F. S. N. Lobo, Phys. Rev. D **71**, 084011 (2005) [arXiv:gr-qc/0502099];
 R. G. Cai, H. S. Zhang and A. Wang, Commun. Theor. Phys. **44**, 948 (2005) [arXiv:hep-th/0505186];
 I. Y. Aref'eva, A. S. Koshelev and S. Y. Vernov, Theor. Math. Phys. **148**, 895 (2006) [Teor. Mat. Fiz. **148**, 23 (2006)] [arXiv:astro-ph/0412619];
 Phys. Rev. D **72**, 064017 (2005) [arXiv:astro-ph/0507067];
 H. Q. Lu, Z. G. Huang and W. Fang, arXiv:hep-th/0504038;
 W. Godlowski and M. Szydlowski, Phys. Lett. B **623**, 10 (2005) [arXiv:astro-ph/0507322];
 J. Sola and H. Stefancic, Phys. Lett. B **624**, 147 (2005) [arXiv:astro-ph/0505133];
 B. Guberina, R. Horvat and H. Nikolic, Phys. Rev. D **72**, 125011 (2005) [arXiv:astro-ph/0507666];
 M. P. Dabrowski, C. Kiefer and B. Sandhofer, Phys. Rev. D **74**, 044022 (2006) [arXiv:hep-th/0605229];
 E. M. Barboza and N. A. Lemos, arXiv:gr-qc/0606084.
- [34] S. Nojiri, S. D. Odintsov and S. Tsujikawa, Phys. Rev. D **71**, 063004 (2005) [arXiv:hep-th/0501025].
- [35] J. D. Barrow, Class. Quant. Grav. **21**, L79 (2004) [arXiv:gr-qc/0403084];
 S. Cotsakis and I. Klaoudatou, J. Geom. Phys. **55**, 306 (2005) [arXiv:gr-qc/0409022];
 S. Nojiri and S. D. Odintsov, Phys. Lett. B **595**, 1 (2004) [arXiv:hep-th/0405078];
 J. D. Barrow, Class. Quant. Grav. **21**, 5619 (2004) [arXiv:gr-qc/0409062];
 V. Sahni and Y. Shtanov, JCAP **0311**, 014 (2003) [arXiv:astro-ph/0202346];
 K. Lake, Class. Quant. Grav. **21**, L129 (2004) [arXiv:gr-qc/0407107];
 J. D. Barrow and C. G. Tsagas, Class. Quant. Grav. **22**, 1563 (2005) [arXiv:gr-qc/0411045];
 M. P. Dabrowski, Phys. Rev. D **71**, 103505 (2005) [arXiv:gr-qc/0410033];
 Phys. Lett. B **625**, 184 (2005); A. Balcerzak and M. P. Dabrowski, Phys. Rev. D **73**, 101301 (2006) [arXiv:hep-th/0604034];
 L. Fernandez-Jambrina and R. Lazkoz, Phys. Rev. D **70**, 121503 (2004) [arXiv:gr-qc/0410124]; Phys. Rev. D **74**, 064030 (2006) [arXiv:gr-qc/0607073]; arXiv:0805.2284 [gr-qc];
 P. Tretyakov, A. Toporensky, Y. Shtanov and V. Sahni, Class. Quant. Grav. **23**, 3259 (2006) [arXiv:gr-qc/0510104];
 H. Stefancic, Phys. Rev. D **71**, 084024 (2005) [arXiv:astro-ph/0411630];
 A. V. Yurov, A. V. Astashenok and P. F. Gonzalez-Diaz, Grav. Cosmol. **14**, 205 (2008) [arXiv:0705.4108 [astro-ph]];
 I. Brevik and O. Gorbunova, Eur. Phys. J. C **56**, 425 (2008) [arXiv:0806.1399 [gr-qc]];
 M. Bouhmadi-Lopez, P. F. Gonzalez-Diaz and P. Martin-Moruno, Phys. Lett. B **659**, 1 (2008) [arXiv:gr-qc/0612135]; arXiv:0707.2390 [gr-qc];
 M. Sami, P. Singh and S. Tsujikawa, Phys. Rev. D **74**, 043514 (2006) [arXiv:gr-qc/0605113];
 C. Cattoen and M. Visser, Class. Quant. Grav. **22**, 4913 (2005) [arXiv:gr-qc/0508045];
 J. D. Barrow and S. Z. W. Lip, arXiv:0901.1626 [gr-qc];
 M. Bouhmadi-Lopez, Y. Tavakoli and P. V. Moniz, arXiv:0911.1428 [gr-qc].
- [36] S. Nojiri and S. D. Odintsov, Phys. Rev. D **78**, 046006 (2008) [arXiv:0804.3519 [hep-th]].
- [37] M. C. B. Abdalla, S. Nojiri and S. D. Odintsov, Class. Quant. Grav. **22**, L35 (2005) [arXiv:hep-th/0409177].
- [38] S. A. Appleby and R. A. Battye, Phys. Lett. B **654**, 7 (2007) [arXiv:0705.3199 [astro-ph]];
 A. A. Starobinsky, JETP Lett. **86**, 157 (2007) [arXiv:0706.2041 [astro-ph]];
 S. Nojiri and S. D. Odintsov, Phys. Lett. B **652**, 343 (2007) [arXiv:0706.1378 [hep-th]];
 L. Pogosian and A. Silvestri, Phys. Rev. D **77**, 023503 (2008) [Erratum-ibid. D **81**, 049901 (2010)] [arXiv:0709.0296 [astro-ph]];

- S. Capozziello and S. Tsujikawa, Phys. Rev. D **77**, 107501 (2008) [arXiv:0712.2268 [gr-qc]].
- [39] K. Bamba, S. Nojiri and S. D. Odintsov, JCAP **0810**, 045 (2008) [arXiv:0807.2575 [hep-th]].
- [40] S. Capozziello, M. De Laurentis, S. Nojiri and S. D. Odintsov, Phys. Rev. D **79**, 124007 (2009) [arXiv:0903.2753 [hep-th]].
- [41] K. Bamba, S. D. Odintsov, L. Sebastiani and S. Zerbini, Eur. Phys. J. C **67**, 295 (2010) [arXiv:0911.4390 [hep-th]].
- [42] T. Kobayashi and K. i. Maeda, Phys. Rev. D **79**, 024009 (2009) [arXiv:0810.5664 [astro-ph]].
- [43] M. Sami, arXiv:0904.3445 [hep-th].
- [44] I. Thongkool, M. Sami, R. Gannouji and S. Jhingan, Phys. Rev. D **80**, 043523 (2009) [arXiv:0906.2460 [hep-th]].
- [45] E. Babichev and D. Langlois, Phys. Rev. D **81**, 124051 (2010) [arXiv:0911.1297 [gr-qc]].
- [46] S. A. Appleby and J. Weller, arXiv:1008.2693 [astro-ph.CO].
- [47] S. Nojiri and S. D. Odintsov, Phys. Lett. B **686**, 44 (2010) [arXiv:0911.2781 [hep-th]].
- [48] K. Bamba, C. Q. Geng and C. C. Lee, JCAP **1008**, 021 (2010) [arXiv:1005.4574 [astro-ph.CO]];
E. V. Linder, Phys. Rev. D **80**, 123528 (2009) [arXiv:0905.2962 [astro-ph.CO]];
L. Yang, C. C. Lee, L. W. Luo and C. Q. Geng, arXiv:1010.2058 [astro-ph.CO].
- [49] S. Nojiri and S. D. Odintsov, arXiv:1008.4275 [hep-th].
- [50] J. Khoury and A. Weltman, Phys. Rev. D **69**, 044026 (2004) [arXiv:astro-ph/0309411].
- [51] P. Brax, C. van de Bruck, A. C. Davis, J. Khoury and A. Weltman, Phys. Rev. D **70** (2004) 123518 [arXiv:astro-ph/0408415].
- [52] S. Nojiri and S. D. Odintsov, Phys. Lett. B **631**, 1 (2005) [arXiv:hep-th/0508049].
- [53] S. Nojiri, S. D. Odintsov and O. G. Gorbunova, J. Phys. A **39**, 6627 (2006) [arXiv:hep-th/0510183].
- [54] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, Phys. Rev. D **73**, 084007 (2006) [arXiv:hep-th/0601008].
- [55] S. Nojiri, S. D. Odintsov and M. Sasaki, Phys. Rev. D **71**, 123509 (2005) [arXiv:hep-th/0504052].
- [56] M. Sami, A. Toporensky, P. V. Tretyakov and S. Tsujikawa, Phys. Lett. B **619**, 193 (2005) [arXiv:hep-th/0504154].
- [57] G. Calcagni, S. Tsujikawa and M. Sami, Class. Quant. Grav. **22**, 3977 (2005) [arXiv:hep-th/0505193].
- [58] H. M. Sadjadi, arXiv:1009.2941 [gr-qc];
R. Myrzakulov, D. Saez-Gomez and A. Tureanu, arXiv:1009.0902 [gr-qc];
L. Sebastiani, arXiv:1008.3041 [gr-qc];
J. Middleton, arXiv:1007.4669 [gr-qc];
A. De Felice and T. Tanaka, Prog. Theor. Phys. **124**, 503 (2010) [arXiv:1006.4399 [astro-ph.CO]];
O. Gorbunova and L. Sebastiani, arXiv:1004.1505 [gr-qc];
J. Moldenhauer, M. Ishak, J. Thompson and D. A. Easson, Phys. Rev. D **81**, 063514 (2010) [arXiv:1004.2459 [astro-ph.CO]];
I. D. Saltas and M. Hindmarsh, arXiv:1002.1710 [gr-qc];
E. Elizalde, R. Myrzakulov, V. V. Obukhov and D. Saez-Gomez, Class. Quant. Grav. **27**, 095007 (2010) [arXiv:1001.3636 [gr-qc]];
R. Di Criscienzo and S. Zerbini, J. Math. Phys. **50**, 103517 (2009) [Erratum-ibid. **51**, 059901 (2010)] [arXiv:0907.4265 [hep-th]];
A. De Felice and S. Tsujikawa, Phys. Rev. D **80**, 063516 (2009) [arXiv:0907.1830 [hep-th]];
N. Goheer, R. Goswami, P. K. S. Dunsby and K. Ananda, Phys. Rev. D **79**, 121301 (2009) [arXiv:0904.2559 [gr-qc]];
S. Y. Zhou, E. J. Copeland and P. M. Saffin, JCAP **0907**, 009 (2009) [arXiv:0903.4610 [gr-qc]];
K. Uddin, J. E. Lidsey and R. Tavakol, Gen. Rel. Grav. **41**, 2725 (2009) [arXiv:0903.0270 [gr-qc]];
C. G. Boehmer and F. S. N. Lobo, Phys. Rev. D **79**, 067504 (2009) [arXiv:0902.2982 [gr-qc]];
M. Alimohammadi and A. Ghalee, Phys. Rev. D **79**, 063006 (2009) [arXiv:0811.1286 [gr-qc]];
S. C. Davis, arXiv:0709.4453 [hep-th];
B. Li, J. D. Barrow and D. F. Mota, Phys. Rev. D **76**, 044027 (2007) [arXiv:0705.3795 [gr-qc]];
I. H. Brevik and J. Quiroga, Int. J. Mod. Phys. D **16**, 817 (2007) [arXiv:gr-qc/0610044].
- [59] D. G. Boulware and S. Deser, Phys. Lett. B **175**, 409 (1986).
- [60] S. Nojiri, S. D. Odintsov and M. Sami, Phys. Rev. D **74**, 046004 (2006) [arXiv:hep-th/0605039].
- [61] M. Satoh, arXiv:1008.2724 [astro-ph.CO];
Z. K. Guo and D. J. Schwarz, Phys. Rev. D **81**, 123520 (2010) [arXiv:1001.1897 [hep-th]];
K. Nozari, T. Azizi and M. R. Setare, JCAP **0910**, 022 (2009) [arXiv:0910.0611 [gr-qc]];
Z. K. Guo and D. J. Schwarz, Phys. Rev. D **80**, 063523 (2009) [arXiv:0907.0427 [hep-th]];
J. Sadeghi, M. R. Setare, A. Banijamali and F. Milani, Phys. Rev. D **79**, 123003 (2009);
T. S. Koivisto, D. F. Mota and C. Pitrou, JHEP **0909**, 092 (2009) [arXiv:0903.4158 [astro-ph.CO]];
M. Gurses, Gen. Rel. Grav. **40**, 1825 (2008);
M. R. Setare and E. N. Saridakis, Phys. Lett. B **670**, 1 (2008) [arXiv:0810.3296 [hep-th]];
B. C. Paul and S. Ghose, Gen. Rel. Grav. **42**, 795 (2010) [arXiv:0809.4131 [hep-th]];
M. Satoh and J. Soda, JCAP **0809**, 019 (2008) [arXiv:0806.4594 [astro-ph]];
R. G. Cai, B. Hu and S. Koh, Phys. Lett. B **671**, 181 (2009) [arXiv:0806.2508 [hep-th]];
D. Bazeia, R. Menezes and A. Y. Petrov, Eur. Phys. J. C **58**, 171 (2008) [arXiv:0806.2299 [hep-th]];
R. Chingangbam, M. Sami, P. V. Tretyakov and A. V. Toporensky, Phys. Lett. B **661**, 162 (2008) [arXiv:0711.2122 [hep-th]];
E. Elizalde, J. Q. Hurtado and H. I. Arcos, Int. J. Mod. Phys. D **17**, 2159 (2008) [arXiv:0708.0591 [gr-qc]];
K. Bamba, Z. K. Guo and N. Ohta, Prog. Theor. Phys. **118**, 879 (2007) [arXiv:0707.4334 [hep-th]];
S. Nojiri, S. D. Odintsov and P. V. Tretyakov, Phys. Lett. B **651**, 224 (2007) [arXiv:0704.2520 [hep-th]];

- B. M. Leith and I. P. Neupane, JCAP **0705**, 019 (2007) [arXiv:hep-th/0702002];
 G. Cognola, E. Elizalde, S. Nojiri, S. Odintsov and S. Zerbini, Phys. Rev. D **75**, 086002 (2007) [arXiv:hep-th/0611198];
 T. P. Sotiriou and E. Barausse, Phys. Rev. D **75**, 084007 (2007) [arXiv:gr-qc/0612065];
 T. Koivisto and D. F. Mota, Phys. Rev. D **75**, 023518 (2007) [arXiv:hep-th/0609155];
 S. Tsujikawa and M. Sami, JCAP **0701**, 006 (2007) [arXiv:hep-th/0608178];
 A. K. Sanyal, Phys. Lett. B **645**, 1 (2007) [arXiv:astro-ph/0608104];
 T. Koivisto and D. F. Mota, Phys. Lett. B **644**, 104 (2007) [arXiv:astro-ph/0606078];
 B. M. N. Carter and I. P. Neupane, JCAP **0606**, 004 (2006) [arXiv:hep-th/0512262];
 B. M. N. Carter and I. P. Neupane, Phys. Lett. B **638**, 94 (2006) [arXiv:hep-th/0510109].
- [62] E. Elizalde, S. Jhingan, S. Nojiri, S. D. Odintsov, M. Sami and I. Thongkool, Eur. Phys. J. C **53**, 447 (2008) [arXiv:0705.1211 [hep-th]].
- [63] S. Deser and R. P. Woodard, Phys. Rev. Lett. **99**, 111301 (2007) [arXiv:0706.2151 [astro-ph]];
 C. Deffayet and R. P. Woodard, JCAP **0908**, 023 (2009) [arXiv:0904.0961 [gr-qc]].
- [64] S. Nojiri and S. D. Odintsov, Phys. Lett. B **659**, 821 (2008) [arXiv:0708.0924 [hep-th]].
- [65] S. Jhingan, S. Nojiri, S. D. Odintsov, M. Sami, I. Thongkool and S. Zerbini, Phys. Lett. B **663**, 424 (2008) [arXiv:0803.2613 [hep-th]].
- [66] S. Nojiri, S. D. Odintsov, M. Sasaki and Y. I. Zhang, arXiv:1010.5375 [gr-qc];
 T. Koivisto, Phys. Rev. D **77**, 123513 (2008) [arXiv:0803.3399 [gr-qc]];
 T. S. Koivisto, Phys. Rev. D **78**, 123505 (2008) [arXiv:0807.3778 [gr-qc]];
 N. A. Koshelev, Grav. Cosmol. **15**, 220 (2009) [arXiv:0809.4927 [gr-qc]];
 S. Nesseris and A. Mazumdar, Phys. Rev. D **79**, 104006 (2009) [arXiv:0902.1185 [astro-ph.CO]];
 K. A. Bronnikov and E. Elizalde, Phys. Rev. D **81**, 044032 (2010) [arXiv:0910.3929 [hep-th]];
 G. Calcagni and G. Nardelli, arXiv:1004.5144 [hep-th].
- [67] S. Capozziello, E. Elizalde, S. Nojiri and S. D. Odintsov, Phys. Lett. B **671** (2009) 193 [arXiv:0809.1535 [hep-th]].
- [68] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, Eur. Phys. J. C **64**, 483 (2009) [arXiv:0905.0543 [gr-qc]].
- [69] S. Nojiri and S. D. Odintsov, Phys. Lett. B **599**, 137 (2004) [arXiv:astro-ph/0403622].
- [70] G. Allemandi, A. Borowiec, M. Francaviglia and S. D. Odintsov, Phys. Rev. D **72**, 063505 (2005) [arXiv:gr-qc/0504057].
- [71] S. Capozziello, G. Lambiase and H. J. Schmidt, Annalen Phys. **9**, 39 (2000) [arXiv:gr-qc/9906051].
- [72] T. Harko, Phys. Lett. B **669**, 376 (2008) [arXiv:0810.0742 [gr-qc]];
 T. Harko, Phys. Rev. D **81**, 044021 (2010) [arXiv:1001.5349 [gr-qc]];
 L. N. Granda and W. Cardona, JCAP **1007**, 021 (2010) [arXiv:1005.2716 [hep-th]];
 T. Harko and F. S. N. Lobo, arXiv:1008.4193 [gr-qc];
 T. Harko, F. S. N. Lobo, S. Nojiri and S. D. Odintsov, arXiv:1104.2669 [gr-qc].
- [73] V. Faraoni, Phys. Rev. D **80**, 124040 (2009) [arXiv:0912.1249 [astro-ph.GA]];
 O. Bertolami, C. G. Boehmer, T. Harko and F. S. N. Lobo, Phys. Rev. D **75**, 104016 (2007) [arXiv:0704.1733 [gr-qc]].
- [74] T. Harko and F. S. N. Lobo, arXiv:1007.4415 [gr-qc];
 C. Deffayet, O. Pujolas, I. Sawicki and A. Vikman, arXiv:1008.0048 [hep-th];
 M. Mohseni, Phys. Lett. B **682**, 89 (2009) [arXiv:0911.2754 [hep-th]];
 L. N. Granda, JCAP **1007**, 006 (2010) [arXiv:0911.3702 [hep-th]];
 C. Gao, JCAP **1006**, 023 (2010) [arXiv:1002.4035 [gr-qc]];
 G. Cognola, E. Elizalde, S. Nojiri and S. D. Odintsov, Open Astron. J. **3**, 20 (2010) [arXiv:0909.2747 [gr-qc]];
 J. Sadeghi, M. R. Setare and A. Banijamali, Eur. Phys. J. C **64**, 433 (2009) [arXiv:0906.0713 [hep-th]];
 A. Borowiec, arXiv:0812.4383 [gr-qc];
 S. Nesseris, Phys. Rev. D **79**, 044015 (2009) [arXiv:0811.4292 [astro-ph]];
 T. S. Koivisto, arXiv:0811.1957 [astro-ph];
 D. Puetzfeld and Y. N. Obukhov, Phys. Rev. D **78**, 121501 (2008) [arXiv:0811.0913 [astro-ph]];
 T. P. Sotiriou and V. Faraoni, Class. Quant. Grav. **25**, 205002 (2008) [arXiv:0805.1249 [gr-qc]];
 T. P. Sotiriou, Phys. Lett. B **664**, 225 (2008) [arXiv:0805.1160 [gr-qc]];
 S. M. Barr, S. P. Ng and R. J. Scherrer, Phys. Rev. D **73**, 063530 (2006) [arXiv:hep-ph/0601053];
 T. Inagaki, S. Nojiri and S. D. Odintsov, JCAP **0506**, 010 (2005) [arXiv:gr-qc/0504054];
 O. Bertolami, J. Paramos, T. Harko and F. S. N. Lobo, arXiv:0811.2876 [gr-qc];
 O. Bertolami and J. Paramos, Class. Quant. Grav. **25**, 245017 (2008) [arXiv:0805.1241 [gr-qc]].
- [75] K. Bamba and S. D. Odintsov, JCAP **0804**, 024 (2008) [arXiv:0801.0954 [astro-ph]].
- [76] A. B. Balakin and W. T. Ni, Class. Quant. Grav. **27**, 055003 (2010) [arXiv:0911.2946 [gr-qc]];
 H. J. Mosquera Cuesta and G. Lambiase, Phys. Rev. D **80**, 023013 (2009) [arXiv:0907.3678 [astro-ph.CO]];
 A. B. Balakin, V. V. Bochkarev and J. P. S. Lemos, Phys. Rev. D **77**, 084013 (2008) [arXiv:0712.4066 [gr-qc]];
 A. B. Balakin and J. P. S. Lemos, Class. Quant. Grav. **22**, 1867 (2005) [arXiv:gr-qc/0503076];
 J. B. Jimenez and A. L. Maroto, Phys. Lett. B **686**, 175 (2010) [arXiv:0903.4672 [astro-ph.CO]].
- [77] K. Bamba, S. Nojiri and S. D. Odintsov, Phys. Rev. D **77**, 123532 (2008) [arXiv:0803.3384 [hep-th]].
- [78] W. Zhao, Y. Zhang and M. L. Tong, arXiv:0909.3874 [astro-ph.CO];
 B. Himmetoglu, C. R. Contaldi and M. Peloso, Phys. Rev. D **80**, 123530 (2009) [arXiv:0909.3524 [astro-ph.CO]];
 A. B. Balakin, H. Dehnen and A. E. Zayats, Annals Phys. **323**, 2183 (2008) [arXiv:0804.2196 [gr-qc]];
 A. B. Balakin, H. Dehnen and A. E. Zayats, Int. J. Mod. Phys. D **17**, 1255 (2008) [arXiv:0710.4992 [gr-qc]];
 E. Elizalde, J. E. Lidsey, S. Nojiri and S. D. Odintsov, Phys. Lett. B **574**, 1 (2003) [arXiv:hep-th/0307177].

- [79] M. Chaichian, S. Nojiri, S. D. Odintsov, M. Oksanen and A. Tureanu, *Class. Quant. Grav.* **27**, 185021 (2010) [arXiv:1001.4102 [hep-th]];
S. Carloni, M. Chaichian, S. Nojiri, S. D. Odintsov, M. Oksanen and A. Tureanu, arXiv:1003.3925 [hep-th].
- [80] R. L. Arnowitt, S. Deser and C. W. Misner, arXiv:gr-qc/0405109, originally “Gravitation: An Introduction to Current Research”, L. Witten ed., Wiley, New York, 1962.
- [81] C. Gao, arXiv:0905.0310 [astro-ph.CO].
- [82] R. M. Wald, *General Relativity*, University of Chicago Press, 1984, Chicago and London.
- [83] É.ourgoulhon, arXiv:gr-qc/0703035.
- [84] P. Hořava, *Phys. Rev. D* **79**, 084008 (2009) [arXiv:0901.3775 [hep-th]].
- [85] J. Kluson, arXiv:0910.5852 [hep-th];
J. Kluson, *JHEP* **0911**, 078 (2009) [arXiv:0907.3566 [hep-th]].
- [86] M. Henneaux, A. Kleinschmidt and G. L. Gomez, arXiv:0912.0399 [hep-th].
- [87] M. Li and Y. Pang, *JHEP* **0908**, 015 (2009) [arXiv:0905.2751 [hep-th]].
- [88] S. Mukohyama, *Phys. Rev. D* **80**, 064005 (2009) [arXiv:0905.3563 [hep-th]].
- [89] E. Elizalde, S. Nojiri, S. D. Odintsov and D. Saez-Gomez, arXiv:1006.3387 [hep-th].
- [90] T. Takahashi and J. Soda, *Phys. Rev. Lett.* **102**, 231301 (2009) [arXiv:0904.0554 [hep-th]].
- [91] E. Kiritsis and G. Kofinas, *Nucl. Phys. B* **821**, 467 (2009) [arXiv:0904.1334 [hep-th]].
- [92] R. Brandenberger, *Phys. Rev. D* **80**, 043516 (2009) [arXiv:0904.2835 [hep-th]].
- [93] S. Mukohyama, K. Nakayama, F. Takahashi and S. Yokoyama, *Phys. Lett. B* **679**, 6 (2009) [arXiv:0905.0055 [hep-th]].
- [94] T. P. Sotiriou, M. Visser and S. Weinfurtner, *JHEP* **0910**, 033 (2009) [arXiv:0905.2798 [hep-th]].
- [95] E. N. Saridakis, *Eur. Phys. J. C* **67**, 229 (2010) [arXiv:0905.3532 [hep-th]].
- [96] M. Minamitsuji, *Phys. Lett. B* **684**, 194 (2010) [arXiv:0905.3892 [astro-ph.CO]].
- [97] G. Calcagni, *Phys. Rev. D* **81**, 044006 (2010) [arXiv:0905.3740 [hep-th]].
- [98] A. Wang and Y. Wu, *JCAP* **0907**, 012 (2009) [arXiv:0905.4117 [hep-th]].
- [99] M. i. Park, *JHEP* **0909**, 123 (2009) [arXiv:0905.4480 [hep-th]].
- [100] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **81**, 043001 (2010) [arXiv:0905.4213 [hep-th]].
- [101] M. Jamil, E. N. Saridakis and M. R. Setare, *Phys. Lett. B* **679**, 172 (2009) [arXiv:0906.2847 [hep-th]].
- [102] C. Bogdanos and E. N. Saridakis, *Class. Quant. Grav.* **27**, 075005 (2010) [arXiv:0907.1636 [hep-th]].
- [103] C. G. Boehmer and F. S. N. Lobo, arXiv:0909.3986 [gr-qc].
- [104] I. Bakas, F. Bourliot, D. Lust and M. Petropoulos, *Class. Quant. Grav.* **27**, 045013 (2010) [arXiv:0911.2665 [hep-th]].
- [105] G. Calcagni, *JHEP* **0909**, 112 (2009) [arXiv:0904.0829 [hep-th]].
- [106] S. Carloni, E. Elizalde and P. J. Silva, *Class. Quant. Grav.* **27**, 045004 (2010) [arXiv:0909.2219 [hep-th]].
- [107] X. Gao, Y. Wang, W. Xue and R. Brandenberger, *JCAP* **1002**, 020 (2010) [arXiv:0911.3196 [hep-th]].
- [108] Y. S. Myung, Y. W. Kim, W. S. Son and Y. J. Park, arXiv:0911.2525 [gr-qc].
- [109] E. J. Son and W. Kim, arXiv:1003.3055 [hep-th].
- [110] A. Wang, arXiv:1003.5152 [hep-th].
- [111] A. Ali, S. Dutta, E. N. Saridakis and A. A. Sen, arXiv:1004.2474 [astro-ph.CO].
- [112] J. O. Gong, S. Koh and M. Sasaki, *Phys. Rev. D* **81**, 084053 (2010) [arXiv:1002.1429 [hep-th]].
- [113] E. Elizalde and D. Saez-Gomez, *Phys. Rev. D* **80**, 044030 (2009) [arXiv:0903.2732 [hep-th]].
- [114] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **691**, 60 (2010) [arXiv:1004.3613 [hep-th]].
- [115] S. Nojiri and S. D. Odintsov, arXiv:1007.4856 [hep-th].
- [116] J. Kluson, S. Nojiri and S. D. Odintsov, arXiv:1104.4286 [hep-th].
- [117] M. Chaichian and A. Demichev, *Path Integrals in Physics*, Vol. I and II, Institute of Physics Publishing, 2001, Bristol and Philadelphia;
M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princeton University Press, 1994, Princeton, New Jersey.
- [118] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **72**, 023003 (2005) [arXiv:hep-th/0505215];
S. Capozziello, V. F. Cardone, E. Elizalde, S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **73**, 043512 (2006) [arXiv:astro-ph/0508350].
- [119] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **70**, 103522 (2004) [arXiv:hep-th/0408170].
- [120] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **74**, 086005 (2006) [arXiv:hep-th/0608008].
- [121] S. Nojiri and S. D. Odintsov, *J. Phys. Conf. Ser.* **66**, 012005 (2007) [arXiv:hep-th/0611071]; hep-th/0610164;
- [122] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **676**, 94 (2009) [arXiv:0903.5231 [hep-th]];
S. Nojiri, S. D. Odintsov and H. Stefancic, *Phys. Rev. D* **74**, 086009 (2006) [arXiv:hep-th/0608168].
- [123] S. Nojiri and S. D. Odintsov, *Gen. Rel. Grav.* **38**, 1285 (2006) [arXiv:hep-th/0506212].
- [124] S. Capozziello, S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **632**, 597 (2006) [arXiv:hep-th/0507182].
- [125] A. Vikman, *Phys. Rev. D* **71**, 023515 (2005) [arXiv:astro-ph/0407107].
- [126] E. Komatsu *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **180**, 330 (2009) [arXiv:0803.0547 [astro-ph]].
- [127] D. N. Spergel *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **148**, 175 (2003) [arXiv:astro-ph/0302209].
- [128] H. V. Peiris *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **148**, 213 (2003) [arXiv:astro-ph/0302225].
- [129] D. N. Spergel *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **170**, 377 (2007) [arXiv:astro-ph/0603449].
- [130] C. Brans and R. H. Dicke, *Phys. Rev.* **124**, 925 (1961).
- [131] E. Elizalde, S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **70**, 043539 (2004) [arXiv:hep-th/0405034].

- [132] S. Nojiri, H. Yoshioka, unpublished.
- [133] E. Elizalde, S. Nojiri, S. D. Odintsov, D. Saez-Gomez and V. Faraoni, Phys. Rev. D **77** (2008) 106005 [arXiv:0803.1311 [hep-th]].
- [134] J. Matsumoto and S. Nojiri, Phys. Lett. B **687**, 236 (2010) [arXiv:1001.0220 [hep-th]].
- [135] K. Bamba, C. Q. Geng, S. Nojiri and S. D. Odintsov, Phys. Rev. D **79**, 083014 (2009) [arXiv:0810.4296 [hep-th]].
- [136] S. Capozziello, J. Matsumoto, S. Nojiri and S. D. Odintsov, Phys. Lett. B **693**, 198 (2010) [arXiv:1004.3691 [hep-th]].
- [137] E. A. Lim, I. Sawicki and A. Vikman, JCAP **1005**, 012 (2010) [arXiv:1003.5751 [astro-ph.CO]];
C. Gao, Y. Gong, X. Wang and X. Chen, arXiv:1003.6056 [astro-ph.CO];
Y. F. Cai and E. N. Saridakis, arXiv:1007.3204 [astro-ph.CO];
J. Kluson, arXiv:1009.6067 [hep-th].
- [138] S. Nojiri, S. D. Odintsov and D. Saez-Gomez, Phys. Lett. B **681**, 74 (2009) [arXiv:0908.1269 [hep-th]].
- [139] P. K. S. Dunsby, E. Elizalde, R. Goswami, S. Odintsov and D. S. Gomez, arXiv:1005.2205 [gr-qc].
- [140] E. Elizalde and A. J. Lopez-Revelles, Phys. Rev. D **82**, 063504 (2010) [arXiv:1004.5021 [hep-th]].
- [141] J. D. Barrow, Phys. Lett. B **235**, 40 (1990).
- [142] F. Briscese, E. Elizalde, S. Nojiri and S. D. Odintsov, Phys. Lett. B **646**, 105 (2007) [arXiv:hep-th/0612220].